



An Introduction to Instantons in Higher Dimensions

Tathagata Ghosh

School of Mathematics Postgraduate Research Conference,
University of Leeds, June 15–17, 2022

School of Mathematics
University of Leeds

1. Introduction
2. Introduction2
3. Differential Forms
 - 3.1. Hodge Star Operator
4. Maxwell's Equations
5. Yang–Mills Equations
6. Finally, Instantons (in 4-dimensions)
7. Instantons in higher Dimensions
8. Why would we care?

Introduction

Introduction2

Introduction


Instantons are solutions of certain equations called instanton equations (duh!). They also solve the Yang–Mills equations.

Introduction

Instantons are solutions of certain equations called instanton equations (duh!). They also solve the Yang–Mills equations. In 1862 James Clerk Maxwell came up with the electromagnetism equations, which unified the electric fields and magnetic fields.

Introduction

Instantons are solutions of certain equations called instanton equations (duh!). They also solve the Yang–Mills equations. In 1862 James Clerk Maxwell came up with the electromagnetism equations, which unified the electric fields and magnetic fields.



James Clerk Maxwell

Introduction

In 1954 Yang–Mills equations appeared as a generalization of Maxwell's theory.

Introduction

In 1954 Yang–Mills equations appeared as a generalization of Maxwell's theory.



Yang Chen-Ning and Robert Mills

Introduction

In 1954 Yang–Mills equations appeared as a generalization of Maxwell's theory.



Yang Chen-Ning and Robert Mills

Since then, the Yang–Mills theory in 4-dimensions have been studied extensively by physicists and mathematicians alike.

Differential Forms

Differential Forms

To describe the Yang–Mills equations and Instantons, we start with the notion of differential forms.

Differential Forms

To describe the Yang–Mills equations and Instantons, we start with the notion of differential forms.

Suppose we are in the Euclidean space \mathbb{R}^3 . Denote a point by (x, y, z) (or, (x_1, x_2, x_3)). We call x, y, z coordinate functions. Now, we consider the ‘line elements’ dx, dy, dz .

Differential Forms

To describe the Yang–Mills equations and Instantons, we start with the notion of differential forms.

Suppose we are in the Euclidean space \mathbb{R}^3 . Denote a point by (x, y, z) (or, (x_1, x_2, x_3)). We call x, y, z coordinate functions. Now, we consider the ‘line elements’ dx, dy, dz .

This should be familiar to you, since in one dimension \mathbb{R} , you have integrated a function $f(x)$ over some interval $[a, b]$ as $\int_a^b f(x) dx$.

Differential Forms

To describe the Yang–Mills equations and Instantons, we start with the notion of differential forms.

Suppose we are in the Euclidean space \mathbb{R}^3 . Denote a point by (x, y, z) (or, (x_1, x_2, x_3)). We call x, y, z coordinate functions. Now, we consider the ‘line elements’ dx, dy, dz .

This should be familiar to you, since in one dimension \mathbb{R} , you have integrated a function $f(x)$ over some interval $[a, b]$ as $\int_a^b f(x) dx$.

In two dimensions \mathbb{R}^2 , you have integrated a function $f(x, y)$ over some region $R \subset \mathbb{R}^2$:

$$\iint_R f(x, y) dx dy$$


We called $dx dy$ the ‘area element’.

Differential Forms

Now, let's think of dx, dy being vectors, having directions. If we take dx first, then dy , the area element $dx dy$ has normal vector (which is $dx \times dy$) upwards, by thumb rule.


Differential Forms

Now, let's think of dx, dy being vectors, having directions. If we take dx first, then dy , the area element $dx dy$ has normal vector (which is $dx \times dy$) upwards, by thumb rule.



Differential Forms

Now, let's think of dx, dy being vectors, having directions. If we take dx first, then dy , the area element $dx dy$ has normal vector (which is $dx \times dy$) upwards, by thumb rule.



Now, if we take dy first then dx , the area element $dy dx$ has a normal vector ($dy \times dx$) is downwards, opposite to the previous one.

To incorporate this skew symmetry of the area element $dx dy$ (which is due to the issue of 'orientation'), we denote $dx dy$ by

$$dx \wedge dy$$

To incorporate this skew symmetry of the area element $dx dy$ (which is due to the issue of 'orientation'), we denote $dx dy$ by

$$dx \wedge dy$$

\wedge is called wedge product.

To incorporate this skew symmetry of the area element $dx dy$ (which is due to the issue of 'orientation'), we denote $dx dy$ by

$$dx \wedge dy$$

\wedge is called wedge product.

Clearly, $dx \wedge dy = -dy \wedge dx$.

To incorporate this skew symmetry of the area element $dx dy$ (which is due to the issue of 'orientation'), we denote $dx dy$ by

$$dx \wedge dy$$

\wedge is called wedge product.

Clearly, $dx \wedge dy = -dy \wedge dx$.

Similarly, we have the volume element $dx \wedge dy \wedge dz$.

Differential Forms

Let we are still in \mathbb{R}^3 , we call x, y, z or any real valued function a 0-form. We call dx, dy, dz 1-forms, $dx \wedge dy, dy \wedge dz, dz \wedge dx$ 2-forms, and $dx \wedge dy \wedge dz$ a 3-form. Clearly you can generalize this to any dimensions, for \mathbb{R}^n .

Differential Forms

Let we are still in \mathbb{R}^3 , we call x, y, z or any real valued function a 0-form. We call dx, dy, dz 1-forms, $dx \wedge dy, dy \wedge dz, dz \wedge dx$ 2-forms, and $dx \wedge dy \wedge dz$ a 3-form. Clearly you can generalize this to any dimensions, for \mathbb{R}^n . These forms are linearly independent and spans vector spaces (Actually modules, being over (non-commutative) ring of all smooth real valued functions, $\Omega^0(\mathbb{R}^3)$).

Differential Forms

Let we are still in \mathbb{R}^3 , we call x, y, z or any real valued function a 0-form. We call dx, dy, dz 1-forms, $dx \wedge dy, dy \wedge dz, dz \wedge dx$ 2-forms, and $dx \wedge dy \wedge dz$ a 3-form. Clearly you can generalize this to any dimensions, for \mathbb{R}^n . These forms are linearly independent and spans vector spaces (Actually modules, being over (non-commutative) ring of all smooth real valued functions, $\Omega^0(\mathbb{R}^3)$). dx, dy, dz spans $\Omega^1(\mathbb{R}^3) \cong \mathbb{R}^3$, the space of all 1-forms.

Differential Forms

Let we are still in \mathbb{R}^3 , we call x, y, z or any real valued function a 0-form. We call dx, dy, dz 1-forms, $dx \wedge dy, dy \wedge dz, dz \wedge dx$ 2-forms, and $dx \wedge dy \wedge dz$ a 3-form. Clearly you can generalize this to any dimensions, for \mathbb{R}^n . These forms are linearly independent and spans vector spaces (Actually modules, being over (non-commutative) ring of all smooth real valued functions, $\Omega^0(\mathbb{R}^3)$). dx, dy, dz spans $\Omega^1(\mathbb{R}^3) \cong \mathbb{R}^3$, the space of all 1-forms. $dx \wedge dy, dy \wedge dz, dz \wedge dx$ spans $\Omega^2(\mathbb{R}^3)$, the space of all 2-forms.

Differential Forms

Let we are still in \mathbb{R}^3 , we call x, y, z or any real valued function a 0-form. We call dx, dy, dz 1-forms, $dx \wedge dy, dy \wedge dz, dz \wedge dx$ 2-forms, and $dx \wedge dy \wedge dz$ a 3-form. Clearly you can generalize this to any dimensions, for \mathbb{R}^n .

These forms are linearly independent and spans vector spaces (Actually modules, being over (non-commutative) ring of all smooth real valued functions, $\Omega^0(\mathbb{R}^3)$).

dx, dy, dz spans $\Omega^1(\mathbb{R}^3) \cong \mathbb{R}^3$, the space of all 1-forms.

$dx \wedge dy, dy \wedge dz, dz \wedge dx$ spans $\Omega^2(\mathbb{R}^3)$, the space of all 2-forms.

$dx \wedge dy \wedge dz$ spans $\Omega^3(\mathbb{R}^3)$, the space of all 3-forms.

Differential Forms

Let we are still in \mathbb{R}^3 , we call x, y, z or any real valued function a 0-form. We call dx, dy, dz 1-forms, $dx \wedge dy, dy \wedge dz, dz \wedge dx$ 2-forms, and $dx \wedge dy \wedge dz$ a 3-form. Clearly you can generalize this to any dimensions, for \mathbb{R}^n .

These forms are linearly independent and spans vector spaces (Actually modules, being over (non-commutative) ring of all smooth real valued functions, $\Omega^0(\mathbb{R}^3)$).

dx, dy, dz spans $\Omega^1(\mathbb{R}^3) \cong \mathbb{R}^3$, the space of all 1-forms.

$dx \wedge dy, dy \wedge dz, dz \wedge dx$ spans $\Omega^2(\mathbb{R}^3)$, the space of all 2-forms.

$dx \wedge dy \wedge dz$ spans $\Omega^3(\mathbb{R}^3)$, the space of all 3-forms.

Note that in dimensions 3, any 4-form is zero (just like the volume of a plane object is zero).

Any 1-form on \mathbb{R}^3 can be written as $fdx + gdy + hdz$, for real valued functions $f, g, h \in \Omega^0$. any 2 form can be written as $fdx \wedge dy + gdy \wedge dz + hdz \wedge dx$ etc.

Any 1-form on \mathbb{R}^3 can be written as $fdx + gdy + hdz$, for real valued functions $f, g, h \in \Omega^0$. any 2 form can be written as $fdx \wedge dy + gdy \wedge dz + hdz \wedge dx$ etc.

Before moving forward, we should mention that the 'differential' d of a k -form makes it a $k + 1$ -form.

Differential Forms

Any 1-form on \mathbb{R}^3 can be written as $fdx + gdy + hdz$, for real valued functions $f, g, h \in \Omega^0$. any 2 form can be written as $fdx \wedge dy + gdy \wedge dz + hdz \wedge dx$ etc.

Before moving forward, we should mention that the 'differential' d of a k -form makes it a $k + 1$ -form.

Simple example, x is a 0-form whereas dx is 1-form.

Differential Forms

Any 1-form on \mathbb{R}^3 can be written as $fdx + gdy + hdz$, for real valued functions $f, g, h \in \Omega^0$. any 2 form can be written as $fdx \wedge dy + gdy \wedge dz + hdx \wedge dz$ etc.

Before moving forward, we should mention that the 'differential' d of a k -form makes it a $k + 1$ -form.

Simple example, x is a 0-form whereas dx is 1-form.

Another one: $xdy \wedge dz$ is 2-form but $d(xdy \wedge dz) = dx \wedge dy \wedge dz$ is 3-form.

Differential Forms

Any 1-form on \mathbb{R}^3 can be written as $fdx + gdy + hdz$, for real valued functions $f, g, h \in \Omega^0$. any 2 form can be written as $fdx \wedge dy + gdy \wedge dz + hdx \wedge dz$ etc.

Before moving forward, we should mention that the 'differential' d of a k -form makes it a $k + 1$ -form.

Simple example, x is a 0-form whereas dx is 1-form.

Another one: $xdy \wedge dz$ is 2-form but $d(xdy \wedge dz) = dx \wedge dy \wedge dz$ is 3-form.

Important note: $d^2 = 0$.

1. Introduction
2. Introduction2
3. Differential Forms
 - 3.1. Hodge Star Operator
4. Maxwell's Equations
5. Yang–Mills Equations
6. Finally, Instantons (in 4-dimensions)
7. Instantons in higher Dimensions
8. Why would we care?

Hodge Star Operator

We are still in \mathbb{R}^3 . Recall the volume form $dx \wedge dy \wedge dz$. Hodge star of a form is the rest of the terms in the volume form. So,

$$*dx = dy \wedge dz$$

Since, $dx \wedge dy \wedge dz = -dy \wedge dx \wedge dz$,

$$*dy = -dx \wedge dz = dz \wedge dx$$

$$*(dx \wedge dy) = dz \text{ etc.}$$

.

Hodge Star Operator

We are still in \mathbb{R}^3 . Recall the volume form $dx \wedge dy \wedge dz$. Hodge star of a form is the rest of the terms in the volume form. So,

$$*dx = dy \wedge dz$$

Since, $dx \wedge dy \wedge dz = -dy \wedge dx \wedge dz$,

$$*dy = -dx \wedge dz = dz \wedge dx$$

$$*(dx \wedge dy) = dz \text{ etc.}$$

.

Using the Hodge star, we define the 'codifferential' d^* which decreases forms. Define

$$d^* = *d*$$

Hodge Star Operator

Then, in 3-dimensions, for a 2-form F ,

$$F \in \Omega^2$$

$$*F \in \Omega^{3-2} = \Omega^1$$

$$d * F \in \Omega^{3-1} = \Omega^2$$

$$*d * F \in \Omega^{3-2} = \Omega^1$$

Hodge Star Operator

Then, in 3-dimensions, for a 2-form F ,

$$F \in \Omega^2$$

$$*F \in \Omega^{3-2} = \Omega^1$$

$$d * F \in \Omega^{3-1} = \Omega^2$$

$$*d * F \in \Omega^{3-2} = \Omega^1$$

So, $d^* : \Omega^2 \rightarrow \Omega^1$ whereas $d : \Omega^1 \rightarrow \Omega^2$.

Hodge Star Operator

Then, in 3-dimensions, for a 2-form F ,

$$F \in \Omega^2$$

$$*F \in \Omega^{3-2} = \Omega^1$$

$$d * F \in \Omega^{3-1} = \Omega^2$$

$$*d * F \in \Omega^{3-2} = \Omega^1$$

So, $d^* : \Omega^2 \rightarrow \Omega^1$ whereas $d : \Omega^1 \rightarrow \Omega^2$.

Historically, Hodge introduced the star operator precisely to write down the Maxwell's equations in easier manner. So, let's do that now.

Maxwell's Equations

Maxwell's Equations

In \mathbb{R}^3 , the electric field $\mathbf{E} \in \Omega^1(\mathbb{R}^3)$ is a 1-form and the magnetic field $\mathbf{B} \in \Omega^2(\mathbb{R}^3)$ is a 2-form. Let dx^1, dx^2, dx^3 spans $\Omega^1(\mathbb{R}^3)$. Introduce the 4th coordinate $t = x^0$, and so, $dt(= dx^0), dx^1, dx^2, dx^3$ spans $\Omega^1(\mathbb{R}^4)$.

Maxwell's Equations

In \mathbb{R}^3 , the electric field $\mathbf{E} \in \Omega^1(\mathbb{R}^3)$ is a 1-form and the magnetic field $\mathbf{B} \in \Omega^2(\mathbb{R}^3)$ is a 2-form. Let dx^1, dx^2, dx^3 spans $\Omega^1(\mathbb{R}^3)$. Introduce the 4th coordinate $t = x^0$, and so, $dt(= dx^0), dx^1, dx^2, dx^3$ spans $\Omega^1(\mathbb{R}^4)$.

The Electromagnetic field (strength) is a 2-form on space-time \mathbb{R}^4 ,

$$\mathbf{F} = \mathbf{B} + dt \wedge \mathbf{E} \in \Omega^2(\mathbb{R}^4)$$

We can write

$$\mathbf{F} = \sum_{0 \leq \mu, \nu \leq 3} F_{\mu\nu} dx^\mu \wedge dx^\nu.$$

Maxwell's Equations

In \mathbb{R}^3 , the electric field $\mathbf{E} \in \Omega^1(\mathbb{R}^3)$ is a 1-form and the magnetic field $\mathbf{B} \in \Omega^2(\mathbb{R}^3)$ is a 2-form. Let dx^1, dx^2, dx^3 spans $\Omega^1(\mathbb{R}^3)$. Introduce the 4th coordinate $t = x^0$, and so, $dt(= dx^0), dx^1, dx^2, dx^3$ spans $\Omega^1(\mathbb{R}^4)$.

The Electromagnetic field (strength) is a 2-form on space-time \mathbb{R}^4 ,

$$\mathbf{F} = \mathbf{B} + dt \wedge \mathbf{E} \in \Omega^2(\mathbb{R}^4)$$

We can write

$$\mathbf{F} = \sum_{0 \leq \mu, \nu \leq 3} F_{\mu\nu} dx^\mu \wedge dx^\nu.$$

Now, we have

$$d\mathbf{F} = 0$$

which is Bianchi identity, it always holds.

Maxwell's Equations

Since $d^2 = 0$, we have an 1-form $\mathbf{A} \in \Omega^1(\mathbb{R}^4)$ such that

$$\mathbf{F} = d\mathbf{A}$$

called the electromagnetic potential in space-time.

Maxwell's Equations

Since $d^2 = 0$, we have an 1-form $\mathbf{A} \in \Omega^1(\mathbb{R}^4)$ such that

$$\mathbf{F} = d\mathbf{A}$$

called the electromagnetic potential in space-time.

Mathematicians usually call \mathbf{A} the connection 1-form and \mathbf{F} the curvature 2-form.

Maxwell's Equations

Since $d^2 = 0$, we have an 1-form $\mathbf{A} \in \Omega^1(\mathbb{R}^4)$ such that

$$\mathbf{F} = d\mathbf{A}$$

called the electromagnetic potential in space-time.

Mathematicians usually call \mathbf{A} the connection 1-form and \mathbf{F} the curvature 2-form.

The Maxwell's equations (in vacuum) in space-time are

$$d\mathbf{F} = 0$$

$$d^*\mathbf{F} = 0.$$

Yang–Mills Equations

Yang–Mills Equations

The connection 1-form (or, electromagnetic potential) is real valued (we can think of it as a function $A : \mathbb{R}^4 \rightarrow \mathbb{R}$). So, for $x, y \in \mathbb{R}^4$,

$$[A, A](x, y) := A(x)A(y) - A(y)A(x) = 0$$

since for real numbers a, b we have $ab = ba$ (i.e, commutativity).

Yang–Mills Equations

The connection 1-form (or, electromagnetic potential) is real valued (we can think of it as a function $A : \mathbb{R}^4 \rightarrow \mathbb{R}$). So, for $x, y \in \mathbb{R}^4$,

$$[A, A](x, y) := A(x)A(y) - A(y)A(x) = 0$$

since for real numbers a, b we have $ab = ba$ (i.e, commutativity).

We can think of Yang–Mills equations as generalisations of Maxwell's equations where we consider the potential to 2×2 trace-less skew-Hermitian matrix valued function (rather than real i.e., 1×1 matrix-valued). Now, because of non-commutativity of matrices, in general,

$$[A, A](x, y) := A(x)A(y) - A(y)A(x) \neq 0$$

Yang–Mills Equations

In this general setup, the curvature takes the form

$$F_A = dA + [A, A]$$

Yang–Mills Equations

In this general setup, the curvature takes the form

$$F_A = dA + [A, A]$$

Define the covariant derivative

$$d_A = d + [A, \cdot]$$

Yang–Mills Equations

In this general setup, the curvature takes the form

$$F_A = dA + [A, A]$$

Define the covariant derivative

$$d_A = d + [A, \cdot]$$

Then,

$$F_A = d_A A$$

(We should note that because of the extra term, d_A^2 is not in general zero.)

Yang–Mills Equations

In this general setup, the curvature takes the form

$$F_A = dA + [A, A]$$

Define the covariant derivative

$$d_A = d + [A, \cdot]$$

Then,

$$F_A = d_A A$$

(We should note that because of the extra term, d_A^2 is not in general zero.)

Fianlly, we have the equations

$$d_A F_A = 0 \quad (\text{Bianchi identity})$$

$$d_A^* F_A = 0 \quad (\text{Yang–Mills equation})$$

Finally, Instantons (in 4-dimensions)

Instantons in 4-dimensions

We are in 4-dimensional Euclidean space. Let's look at two particular equations

$$*F_A = F_A \text{ (Self-dual Instanton equation)}$$

$$*F_A = -F_A \text{ (Anti-self-dual Instanton equation)}$$

Instantons in 4-dimensions

We are in 4-dimensional Euclidean space. Let's look at two particular equations

$$*F_A = F_A \quad (\text{Self-dual Instanton equation})$$

$$*F_A = -F_A \quad (\text{Anti-self-dual Instanton equation})$$

Dimensionally it makes sense, since

$$F_A \in \Omega^2 \Rightarrow *F_A \in \Omega^{4-2} = \Omega^2 \ni \pm F_A.$$

Instantons in 4-dimensions

We are in 4-dimensional Euclidean space. Let's look at two particular equations

$$*F_A = F_A \quad (\text{Self-dual Instanton equation})$$

$$*F_A = -F_A \quad (\text{Anti-self-dual Instanton equation})$$

Dimensionally it makes sense, since

$$F_A \in \Omega^2 \Rightarrow *F_A \in \Omega^{4-2} = \Omega^2 \ni \pm F_A.$$

The solutions for these equations are the connections for which the curvature satisfies these equations. These connections are called Instantons.

Now,

$$\begin{aligned}d_A^* F_A &= * d_A * f_A \\&= d_A(\pm F_A) \quad (\text{using instanton equations}) \\&= \pm * d_A F_A \\&= 0 \quad (\text{using Bianchi identity})\end{aligned}$$

Instantons in 4-dimensions

Now,

$$\begin{aligned}d_A^* F_A &= * d_A * f_A \\&= d_A(\pm F_A) \quad (\text{using instanton equations}) \\&= \pm * d_A F_A \\&= 0 \quad (\text{using Bianchi identity})\end{aligned}$$

So, we see that instantons are indeed solutions of the Yang–Mills equations.

Instantons in higher Dimensions

Instantons in higher Dimensions

Now, we want to do the same thing on \mathbb{R}^7 . Here

$F_A \in \Omega^2(\mathbb{R}^7)$. But then,

$$F_A \in \Omega^2 \Rightarrow *F_A \in \Omega^{7-2} = \Omega^5 \not\subset \pm F_A.$$

Instantons in higher Dimensions

Now, we want to do the same thing on \mathbb{R}^7 . Here

$F_A \in \Omega^2(\mathbb{R}^7)$. But then,

$$F_A \in \Omega^2 \Rightarrow *F_A \in \Omega^{7-2} = \Omega^5 \not\subset \pm F_A.$$

So, consider a 3-form $\phi \in \Omega^3$,

$$\phi = dx^1 \wedge dx^2 \wedge dx^7 + dx^3 \wedge dx^4 \wedge dx^7 + \dots$$

Instantons in higher Dimensions

Now, we want to do the same thing on \mathbb{R}^7 . Here

$F_A \in \Omega^2(\mathbb{R}^7)$. But then,

$$F_A \in \Omega^2 \Rightarrow *F_A \in \Omega^{7-2} = \Omega^5 \not\subset \pm F_A.$$

So, consider a 3-form $\phi \in \Omega^3$,

$$\phi = dx^1 \wedge dx^2 \wedge dx^7 + dx^3 \wedge dx^4 \wedge dx^7 + \dots$$

Indeed,

$$F_A \in \Omega^2$$

$$\phi \wedge F_A \in \Omega^{3+2} = \Omega^5$$

$$*(\phi \wedge F_A) \in \Omega^{7-5} = \Omega^2 \ni \pm F_A$$

Instantons in higher Dimensions

Now, we want to do the same thing on \mathbb{R}^7 . Here

$F_A \in \Omega^2(\mathbb{R}^7)$. But then,

$$F_A \in \Omega^2 \Rightarrow *F_A \in \Omega^{7-2} = \Omega^5 \not\subset \pm F_A.$$

So, consider a 3-form $\phi \in \Omega^3$,

$$\phi = dx^1 \wedge dx^2 \wedge dx^7 + dx^3 \wedge dx^4 \wedge dx^7 + \dots$$

Indeed,

$$F_A \in \Omega^2$$

$$\phi \wedge F_A \in \Omega^{3+2} = \Omega^5$$

$$*(\phi \wedge F_A) \in \Omega^{7-5} = \Omega^2 \ni \pm F_A$$

we have the instanton equations in 7-dimensions:

$$*(\phi \wedge F_A) = \pm F_A$$

Instantons in higher Dimensions

Now, we want to do the same thing on \mathbb{R}^7 . Here

$F_A \in \Omega^2(\mathbb{R}^7)$. But then,

$$F_A \in \Omega^2 \Rightarrow *F_A \in \Omega^{7-2} = \Omega^5 \not\subset \pm F_A.$$

So, consider a 3-form $\phi \in \Omega^3$,

$$\phi = dx^1 \wedge dx^2 \wedge dx^7 + dx^3 \wedge dx^4 \wedge dx^7 + \dots$$

Indeed,

$$F_A \in \Omega^2$$

$$\phi \wedge F_A \in \Omega^{3+2} = \Omega^5$$

$$*(\phi \wedge F_A) \in \Omega^{7-5} = \Omega^2 \ni \pm F_A$$

we have the instanton equations in 7-dimensions:

$$*(\phi \wedge F_A) = \pm F_A$$

Similarly, in dimensions 8, we use a 4-form $\Phi \in \Omega^4(\mathbb{R}^8)$.




Why would we care?

Why would we care?

Although I described the Yang–Mills equations and Instantons on Euclidean space \mathbb{R}^n , things become more interesting when we consider ‘smooth manifolds’. Euclidean spaces are also manifolds, so are the Earth, a football, a rugby ball, a doughnut etc. And then, there are these complicated manifolds:

Why would we care?

Although I described the Yang–Mills equations and Instantons on Euclidean space \mathbb{R}^n , things become more interesting when we consider ‘smooth manifolds’. Euclidean spaces are also manifolds, so are the Earth, a football, a rugby ball, a doughnut etc. And then, there are these complicated manifolds:




Why would we care?

A manifold is smooth means it dose not have a pointy part:


Why would we care?

A manifold is smooth means it dose not have a pointy part:



Why would we care?


A manifold is smooth means it dose not have a pointy part:



Manifolds are also 'locally Euclidean', just like the football field is flat, but the whole Earth isn't.

Why would we care?

A manifold is smooth means it dose not have a pointy part:



Manifolds are also 'locally Euclidean', just like the football field is flat, but the whole Earth isn't.

When we define Yang–Mills / Instanton equations on these manifolds, the space of all solutions of these equations (we call them moduli spaces) contain a lot of topological / geometric information about the manifolds, and hence extremely lucrative to mathematicians. For similar reasons, the higher dimensional theory is extensively studied by physicists and mathematicians working on string theory and Quantum field theory.

Thank you!