

RELIABLE QUASI-MONTE CARLO  
WITH CONTROL VARIATES

BY  
DA LI

Submitted in partial fulfillment of the  
requirements for the degree of  
Master of Science in Applied Mathematics  
in the Graduate College of the  
Illinois Institute of Technology

Approved \_\_\_\_\_  
Advisor

Chicago, Illinois  
May 2016



# TABLE OF CONTENTS

	Page
LIST OF TABLES . . . . .	iv
LIST OF FIGURES . . . . .	v
LIST OF SYMBOLS . . . . .	vi
ABSTRACT . . . . .	vii
1. INTRODUCTION . . . . .	1
1.1. What are we going to do? . . . . .	1
1.2. Why this is a good idea? . . . . .	1
1.3. What's the challenge? . . . . .	1
1.4. Outline . . . . .	1
CHAPTER	
2. BACKGROUND . . . . .	3
2.1. Problem Setup . . . . .	3
2.2. Sobol Sequence . . . . .	4
2.3. Control Variates . . . . .	4
2.4. Reliable Adaptive QMC with digital sequence . . . . .	5
3. RELIABLE ADAPTIVE QMC SOBOL WITH CV . . . . .	8
3.1. Idea to add control variates to QMC with Digital Sequence	8
3.2. The problem of C.V. with QMC . . . . .	8
3.3. A new way to find $\beta$ . . . . .	9
3.4. The problem with $\theta$ . . . . .	11
3.5. The modified method . . . . .	12
3.6. The Algorithm . . . . .	13
4. NUMERICAL EXPERIMENT . . . . .	15
4.1. Asian Option . . . . .	15
4.2. Barrier Option . . . . .	16
5. CONCLUSION . . . . .	18
5.1. Discussion . . . . .	18
5.2. Future work . . . . .	18
BIBLIOGRAPHY . . . . .	19

# LIST OF TABLES

Table		Page
4.1	Parameter Setup for Up and In Barrier Call Option . . . . .	15
4.2	Results of cubSobol, cv_old and cv_new with Asian Option . . . . .	16
4.3	Parameter Setup for Up and In Barrier Call Option . . . . .	16
4.4	Results of cubSobol, cv_old and cv_new with Barrier Option . . . . .	17

# LIST OF FIGURES

Figure		Page
2.1	Cone condition for reliable adpative QMC algorithm . . . . .	7
4.1	Walsh coefficients of $f$ . . . . .	17

## LIST OF SYMBOLS

Symbol	Definition
$\mathbb{N}$	Positive Integers
$\mathbb{N}_0$	Nonnegative Integers
$\mathbb{Z}$	Integers
$\mathbb{R}$	Real numbers
$\oplus$	Digital addition

## ABSTRACT

With Quasi-Monte Carlo(QMC) methods being implemented in a guaranteed adaptive algorithm recently, possibility of combining that with traditional proficiency improvement techniques for Monte Carlo(MC) such as control variates is brought to table.

The problem for adding control variates to QMC is that optimal control variate coefficient for QMC is generally not the same as MC. Here we propose a method for computing the optimal control variate coefficients with guaranteed adaptive QMC algorithm. One merit of control variate is that there is always a good solution as good as using no control variates. And our method is implemented in an efficient way that even in that case the extra cost for control variates is not significant.

As for examples we will include two 16 dimensional integration on financial problem. One is pricing arithmetic mean Asian option with geometric mean Asian option as control variates and the other is barrier option with European option as control variates. Our results show that with good control variates, the cost of adaptive QMC is greatly reduced compared with no control variates cases.

## CHAPTER 1

### INTRODUCTION

#### 1.1 What are we going to do?

Recently there are some great results from construction of Quasi-Monte Carlo(QMC) methods that can adaptively choose a sample size for given error tolerances [1]. Our work is trying to combine reliable Quasi-Monte Carlo methods with control variates. We will justify the theory behind it, construct a practical algorithm which can be implemented and tested through high dimensional integration examples.

#### 1.2 Why this is a good idea?

Control Variates(CV) is known to be a variance reduction technique for simple Monte-Carlo(MC) methods [2]. While QMC, which can be viewed as a deterministic version of MC, outperforms MC for many integrals. Naturally we are wondering if QMC can also benefit the CV technique. If that is possible, it can be especially useful for problems where we can easily find good control variates.

#### 1.3 What's the challenge?

The challenge is that the optimal control variate coefficient for QMC is generally not the same as for simple Monte Carlo, as explained by Hickernell, Lemieux, and Owen [3]. This requires us to figure out a right way to get the optimal coefficients for control variates with Quasi-Monte Carlo.

#### 1.4 Outline

In chapter 2 we first will briefly introduce Quasi-Monte Carlo rule and its difference between Monte-Carlo. Then we will briefly talk about digital sequence and layout several concepts which will be used later. Chapter 3 will show the derivations and theories of our methods along with the corresponding algorithm. In chapter 4



we will demonstrate results from several numerical experiments. We choose several option pricing problems for our target. For the final chapter we will discuss the results and future extension of the method.

## CHAPTER 2

### BACKGROUND

#### 2.1 Problem Setup

Numerical integration problems is involved in fields such as physics, , mathematical finance, biology, computer graphics, and many others fields. It usually happens when it is hard to solve some integral analytically. Therefore one has to use numerical methods for such problems. And Monte-Carlo method is the general way to solve problems in such case [4]. The method can be simply explained in the following way.

Suppose we have the following standard integration approximation problem whose format is

$$I = \int_{[0,1]^d} f(x) dx \quad (2.1)$$

Then we will take sample of  $n$  points  $\{\mathbf{x}_0, \dots, \mathbf{x}_n\} \in [0, 1]^d$  follow the uniform distribution randomly, then form a estimator in form of

$$\hat{I}(f) = \frac{1}{n} \sum_{i=1}^n f(X_i)$$

However, there are several problem with simple Monte-Carlo method [5]. First, it is difficult to generate truly random samples. Second, error bound for MC works only probabilistic sense. And last, in many applications the convergence rate of MC is considered not fast enough.

Hence QMC method were introduced to address these problems. For QMC method the estimator is almost the same with MC. The difference is that the sample points are taken from low discrepancy sequence, which is deterministically chosen instead of random. We will briefly review one method for constructing such sequence that we used for our application in the next section.

## 2.2 Sobol Sequence

## 2.3 Control Variates

Control variates has been know as variance reduction technique used in Monte Carlo methods. In this section we will brief review the ideas and some results of this methods.

Suppose we want to solve the integration problem (2.1) showed earlier, now we have a known function  $h$  and its value on the interval  $\int_{[0,1]^d} h(x)dx = \theta$ . We then construct a new estimator as the following

$$\hat{I}_{cv}(f) = \frac{1}{n} \sum_{i=1}^n \left[ f(X_i) - \beta_{mc}[h(X_i) - \theta] \right] \quad s.t. \ X_i \sim \mathcal{U}[0, 1), \text{ i.i.d.}$$

We can easily see it's an unbiased estimator, i.e.  $\mathbb{E}(\hat{I}_{cv}) = I$ . Now we want to pick the right  $\beta_{mc}$  such that make the estimation more efficient. For simplicity, we consider single control variate in this case. Base on previous MC error estimating formula (??), we know it's achievable by minimizing the variance of the estimator, which is

$$\begin{aligned} \text{Var}_{mc}(\hat{I}_{cv}) &= \text{Var}\left(\frac{1}{n} \sum_{i=1}^n [f(X_i) - \beta_{mc}[h(X_i) - \theta]]\right) \\ &= \frac{1}{n} \text{Var}\left(f(X_i) - \beta_{mc}[h(X_i) - \theta]\right) \quad \text{by } X_i \text{ i.i.d} \\ &= \frac{1}{n} \mathbb{E}\left([f(X_i) - \beta_{mc}[h(X_i) - \theta] - I]^2\right) \\ &= \frac{1}{n} \mathbb{E}\left([f(X_i) - I] - \beta_{mc}[h(X_i) - \theta]\right)^2 \\ &= \frac{1}{n} \mathbb{E}\left([f(X_i) - I]^2 - 2\beta_{mc}[f(X_i) - I][h(X_i) - \theta] + \beta_{mc}^2[h(X_i) - \theta]^2\right) \\ &= \frac{1}{n} \left( \text{Var}[f(X_i)] - 2\beta_{mc}\text{Cov}[f(X_i), h(X_i)] + \beta_{mc}^2 \text{Var}[h(X_i)] \right) \\ &= \frac{1}{n} \left( \text{Var}[h(X_i)] \left( \beta_{mc} - \frac{\text{Cov}[f(X_i), h(X_i)]}{\text{Var}[h(X_i)]} \right)^2 + \text{Var}[f(X_i)] - \frac{\text{Cov}^2[f(X_i), h(X_i)]}{\text{Var}[h(X_i)]} \right) \end{aligned}$$

Then the optimal  $\beta_{mc}$  is given by

$$\beta_{mc}^* = \frac{\text{Cov}[f(X_i), h(X_i)]}{\text{Var}[h(X_i)]} \quad (2.2)$$

In which case the variance become

$$\text{Var}_{mc}(\hat{I}_{cv}) = \frac{\text{Var}[f(X_i)]}{n} (1 - \text{corr}^2[f(X_i), h(X_i)])$$

and we always have

$$\text{Var}_{mc}(\hat{I}_{cv}) \leq \frac{\text{Var}[f(X_i)]}{n} = \text{Var}_{mc}(\hat{I})$$

Now we can see the merit of control variates as a variance reduction method. In worst case, we get a completely uncorrelated  $g$  that leads correlation to zero, and we have variance exactly the same as not using control variates. On the other hand, the more correlated our control variates is to the target function, the more variance we can get rid of by using the method.

## 2.4 Reliable Adaptive QMC with digital sequence

### 2.4.1 Idea of adaptive cubature algorithm.

One practical problem for QMC method is that how to get the sample size big enough for a required error tolerance. The idea in work of Hickernell and Jimnez Rugama(2014) [1] is to construct a QMC algorithm with reliable error estimation on digital sequence. Here we briefly summarize their results.

The error of QMC method on digital sequence can be expressed in terms of

Walsh coefficients of the integrand on certain cone conditions.

$$\begin{aligned}
 & \text{if } f \in \mathcal{C} \text{ then } \left| \int_{[0,1]^d} f(x) dx - \hat{I}_m(f) \right| \leq a(r, m) \sum_{[2^{m-r-1}]^{2^{m-r}-1}} |\tilde{f}_{m,k}| \quad (2.3) \\
 & \hat{I}_m(f) := \frac{1}{b^m} \sum_{i=0}^{b^m-1} f(z_i \oplus \Delta) \\
 & \tilde{f}_{m,k} = \text{discrete Walsh coefficients of } f \\
 & a(r, m) = \text{inflation factor that depends on } \mathcal{C}
 \end{aligned}$$

Here is the definition of the cone condition.

$$\begin{aligned}
 \mathcal{C} &:= \left\{ f \in L^2[0,1]^d : \bigcirc \leq \hat{\omega}(m-l)\diamond, l \leq m; \quad \diamond \leq \hat{\omega}(m-l)\square, l^* \leq l \leq m \right\} \\
 \bigcirc &:= \sum_{\kappa=\lfloor b^{l-1} \rfloor}^{b^l-1} \sum_{\lambda=1}^{\infty} |\hat{f}_{\kappa+\lambda b^m}|, \quad \square := \sum_{\kappa=b^{l-1}}^{b^l-1} |\hat{f}_{\kappa}|, \quad \diamond := \sum_{\kappa=b^m}^{\infty} |\hat{f}_{\kappa}| \quad (2.4) \\
 & l^* \in \mathbb{N} \text{ be fixed ; } \forall m \in \mathbb{N}, \hat{\omega}(m), \hat{\omega}(m) \geq 0, \text{ and } \lim_{m \rightarrow \infty} \hat{\omega}(m) = 0, \lim_{m \rightarrow \infty} \hat{\omega}(m) = 0
 \end{aligned}$$

The first inequality( $\bigcirc \leq \diamond$ ) means the sum of the larger indexed Walsh coefficients bounds a partial sum of the same coefficients. Take  $l = 0, m = 12$  for example, in Figure 2.1 the the sum of circles should be bounded by some factor times the sum of diamonds. The second inequality( $\diamond \leq \square$ ) requires the sum of the larger Walsh coefficients be bounded by the sum of smaller indexed Walsh coefficients. Take  $l = 8$  at this time, which means in Figure 2.1 the sum of diamonds should be bounded by some relax factor times the squares.

The cone give some meanings for the functions about how they should behave to get the err bound formula (2.3). This means that  $|\hat{f}_{\kappa}|$  does not dramatically bounce back as  $\kappa$  goes to infinity. Note that in Figure 2.1 we call circles the err bound, this is proven to be true and under the cone conditions we can estimate it using discrete Walsh coefficients instead of true Walsh coefficients.




Figure 2.1. Cone condition for reliable adaptive QMC algorithm

## CHAPTER 3

### RELIABLE ADAPTIVE QMC SOBOL WITH CV

#### 3.1 Idea to add control variates to QMC with Digital Sequence

The idea is similar to traditional control variates technique for Monte-Carlo. If we know the integration of a function  $\mathbf{h} = (h_1, \dots, h_J)$  on the interval same as our  $f$ , say  $\int_{[0,1]^d} h_j dx = \theta_j$ , then we can define a new function  $g$

$$g := f - (\mathbf{h} - \boldsymbol{\theta})\boldsymbol{\beta} \quad (3.1)$$

$$\text{s.t. } \boldsymbol{\theta} = (\theta_1, \dots, \theta_J), \boldsymbol{\beta} = (\beta_1, \dots, \beta_J)^T$$

Then easily we can find that if we replace  $f$  with  $g$ , the integration stays the same.

$$\int_{[0,1]^d} g dx = \int_{[0,1]^d} f - (\mathbf{h} - \boldsymbol{\theta})\boldsymbol{\beta} dx = \int_{[0,1]^d} f dx$$

Now we are wondering if we can still use the same method as MC, the answer is no and the reason is in the next section.

#### 3.2 The problem of C.V. with QMC

As we pointed out earlier, QMC use a different way for generating  $X_i$ , they are still identical (from same distribution) but not independent anymore, which caused the problem for control variates.

Suppose  $X_1, \dots, X_n$  are generated by QMC rule, the estimator stays the same

$$\hat{I}_{cv}(f) = \frac{1}{n} \sum_{i=1}^n \left[ f(X_i) - \beta_{qmc}[h(X_i) - \theta] \right] \quad X_i \in \mathcal{U}(0, 1)$$

We can easily prove it is still unbiased

$$\mathbb{E}(\hat{I}_{cv}) = \mathbb{E}\left(\frac{1}{n} \sum_{i=1}^n \left[ f(X_i) - \beta_{mc}[h(X_i) - \theta] \right]\right) = I$$

However, it's not the same case as MC like we presented before, because we do not have i.i.d for  $X_i$  this time

$$\text{Var}_{qmc}(\hat{I}_{cv}) \neq \frac{1}{n} \text{Var}\left(f(X_i) - \beta_{mc}[h(X_i) - \theta]\right)$$

Instead the variance become

$$\begin{aligned} \text{Var}(\hat{I}_{cv}) &= \text{Var}\left(\hat{I} - \beta_{qmc}\hat{H}\right) \quad s.t. \quad \hat{I} = \sum_{i=1}^n f(X_i), \quad \hat{H} = \sum_{i=1}^n [h(X_i) - \theta] \\ &= \text{Var}(\hat{I}) - 2\beta_{qmc}\text{Cov}(\hat{I}, \hat{H}) + \beta_{qmc}^2 \text{Var}(\hat{H}) \\ &= \text{Var}(\hat{H}) \left( \beta_{qmc} - \frac{\text{Cov}(\hat{I}, \hat{H})}{\text{Var}(\hat{H})} \right)^2 + \text{Var}(\hat{I}) - \frac{\text{Cov}(\hat{I}, \hat{H})^2}{\text{Var}(\hat{H})} \end{aligned}$$

The optimal  $\beta_{qmc}$  is

$$\beta_{qmc}^* = \text{Var}(\hat{H})^{-1} \text{Cov}(\hat{I}, \hat{H}) \quad (3.2)$$

which leave the variance to be

$$\text{Var}_{qmc}(\hat{I}_{cv}) = \text{Var}(\hat{I}) (1 - \text{corr}^2[\hat{I}, \hat{H}])$$

Now we are interested that if our previous formula for  $\hat{\beta}_{mc}$  could be an estimation for  $\hat{\beta}_{qmc}$ . The fact is that they are generally not the same. Let's take the covariance part of formula (3.2) and (2.2) to see the difference.

$$\begin{aligned} \text{Cov}(\hat{I}, \hat{H}) &= \int [f(X_1) + \dots + f(X_n)][h(X_1) + \dots + h(X_n)] d\mathbf{X} \\ &= \int \left[ \sum_{i=1}^n f(X_i)h(X_i) + \sum_{i,j=1}^{i \neq j} f(X_i)h(X_j) \right] d\mathbf{X} \\ &\neq \int f(X_i)h(X_i) dX_i \\ &= \text{Cov}x[(f(X_i), h(X_i))] \end{aligned}$$

There is also a very good example from Hicknell and Lemieux(2005) [3]'s paper, showing that  $\beta_{mc}$  and  $\beta_{qmc}$  can make a quite different results in some cases.

### 3.3 A new way to find $\beta$



As we stated in previous section, we can not find optimal  $\beta$  by minimizing variance of estimator like Monte-Carlo. However, if using the guaranteed adaptive QMC method introduced in chapter 2, we may have another way to find  $\beta$ .

Recall equation (2.3) the error bound for new estimator of  $g$  still holds

$$\left| \int_{[0,1]^d} g dx - \hat{I}_m(g) \right| \leq a(r, m) \sum_{\lfloor 2^{m-r-1} \rfloor}^{2^{m-r-1}-1} |\tilde{g}_{m,k}| \quad (3.3)$$

Naturally, the new estimator become

$$\hat{I}_m(g) := \frac{1}{b^m} \sum_{i=0}^{b^m-1} g(z_i + \Delta) \quad (3.4)$$

From (3.3) it is clear that the optimal  $\beta$  is the one that minimize the error term.

$$\beta^* = \min_{\beta} \sum_{\kappa=b^{m-r-1}}^{b^{m-r}-1} |\hat{g}_{\kappa}| \quad (3.5)$$

$$= \min_{\beta} \sum_{\kappa=b^{m-r-1}}^{b^{m-r}-1} |\hat{f}_{\kappa} - (\hat{\mathbf{h}}_{\kappa} - \hat{\boldsymbol{\theta}})\beta| \quad \hat{\mathbf{h}}_{\kappa} = (\hat{h}_{\kappa,1}, \dots, \hat{h}_{\kappa,J}), \hat{\boldsymbol{\theta}} = (\hat{\theta}_{\kappa,1}, \dots, \hat{\theta}_{\kappa,J}) \quad (3.6)$$

$$= \min_{\beta} \|\hat{\mathbf{f}} - \hat{\mathbf{H}}\beta\|_1 \quad \hat{\mathbf{f}} = (\hat{f}_{b^{m-r-1}}, \dots, \hat{f}_{b^m-1})^T \quad (3.7)$$

$$\approx \min_{\beta} \|\hat{\mathbf{f}} - \hat{\mathbf{H}}\beta\|_2 \quad \hat{\mathbf{H}} = (\hat{\mathbf{H}}_1, \dots, \hat{\mathbf{H}}_J) \quad (3.8)$$

$$\hat{\mathbf{H}}_j = (\hat{h}_{b^{m-r-1},j} - \hat{\theta}_j, \dots, \hat{h}_{b^m-1,j} - \hat{\theta}_j)^T$$

The second equivalence is not hard to get, but the third one may not be so obvious. Let's consider it backwards. Suppose we have a vector  $A$  and it's  $\mathcal{L}_1$ -norm.

$$A = \begin{pmatrix} f_1 - z_1 \\ f_2 - z_2 \\ \dots \\ f_n - z_n \end{pmatrix}, \quad \|A\|_1 = \sum_{i=1}^n |f_i - z_i|, \quad z_i := (\mathbf{h}_i - \boldsymbol{\theta})$$

If we replace the index,  $A$  is exactly what's inside the  $\mathcal{L}_1$ -norm in (3.7). Hence we justified the third equivalence. The reason we use an approximation instead, i.e. the  $\mathcal{L}_1$ -norm, is because there is no efficient way to solve it compared to existing least square methods.

### 3.4 The problem with $\theta$

We noticed a problem in solution for optimal  $\beta$ , which is we do a lot of subtractions with  $\theta$ . This could be a large cost when we have difficult functions which means  $b^{m-r}$  could be very large number. Therefore we present a way to avoid that part.

The idea is form a observation that Walsh transform of  $\theta$  in (??) is actually zero, since  $\hat{h}_\theta = \theta \delta_{\kappa,0}$  and the summation is not start from  $\kappa = 0$ .

This simplifies (3.6) to the following. Note that we only need the information of function  $f$  and  $h$  to calculate  $\beta^*$ ,  $\theta$  has been get rid of the optimization process.

$$\begin{aligned}
 \beta^* &= \min_{\beta} \sum_{\kappa=b^{m-r-1}}^{b^{m-r}-1} |\hat{f}_\kappa - (\hat{h}_\kappa - \hat{\theta})\beta| \\
 &= \min_{\beta} \sum_{\kappa=b^{m-r-1}}^{b^{m-r}-1} |\hat{f}_\kappa - (\hat{h}_\kappa - \theta \delta_{\kappa,0})\beta| \\
 &= \min_{\beta} \sum_{\kappa=b^{m-r-1}}^{b^{m-r}-1} |\hat{f}_\kappa - \hat{h}_\kappa \beta| & \hat{\mathbf{f}} &= (\hat{f}_{b^{m-r-1}}, \dots, \hat{f}_{b^{m-r}-1})^T \\
 &= \min_{\beta} \|\hat{\mathbf{f}} - \hat{\mathbf{H}}\beta\|_1 & \hat{\mathbf{H}} &= (\hat{\mathbf{H}}_1, \dots, \hat{\mathbf{H}}_J) \\
 &\approx \min_{\beta} \|\hat{\mathbf{f}} - \hat{\mathbf{H}}\beta\|_2 & \hat{\mathbf{H}}_j &= (\hat{h}_{b^{m-r-1},j}, \dots, \hat{h}_{b^{m-r}-1,j})^T
 \end{aligned} \tag{3.9}$$

The same problem happened with the estimator (3.4). We have the similar

solution for that.

$$\begin{aligned}
\hat{I}_m(g) &= \frac{1}{b^m} \sum_{i=0}^{b^m-1} g(z_i + \Delta) \\
&= \frac{1}{b^m} \sum_{i=0}^{b^m-1} f(z_i + \Delta) - (\mathbf{h}(z_i + \Delta) - \boldsymbol{\theta})\boldsymbol{\beta} \\
&= \frac{1}{b^m} \sum_{i=0}^{b^m-1} [f(z_i + \Delta) - \mathbf{h}(z_i + \Delta)\boldsymbol{\beta}] + \boldsymbol{\theta}\boldsymbol{\beta}
\end{aligned} \tag{3.10}$$

After organize it the in format of (3.10),  $\theta$  is eliminated from the summation part. From these two parts of work on  $\theta$  we managed to save  $\frac{b-1}{b}b^{m-r} + b^m$  operations of subtraction.

### 3.5 The modified method

Now we make the following changes

$$\begin{aligned}
g &= f - \beta h \\
\hat{I}_m(g) &= \frac{1}{b^m} \sum_{i=0}^{b^m-1} g(z_i + \Delta)
\end{aligned}$$

And we have the following equivalence

$$\begin{aligned}
\int_{[0,1]^d} f dx &= \int_{[0,1]^d} g dx + \theta \beta \\
\hat{I}_m(f) &= \hat{I}_m(g) + \theta \beta
\end{aligned}$$

So the estimation error becomes

$$\left| \int_{[0,1]^d} f dx - \hat{I}_m(f) \right| = \left| \int_{[0,1]^d} g dx - \hat{I}_m(g) \right|$$

Here if our  $g$  is in the cone we introduced earlier (2.4), then we can use the results from Hickernell and Jimnez Rugama(2014) [1], the error is bounded by

$$\left| \int_{[0,1]^d} g dx - \hat{I}_m(g) \right| \leq a(r, m) \sum_{\lfloor 2^{m-r-1} \rfloor}^{2^{m-r}-1} |\tilde{g}_{m,k}|$$

This leads to the same algorithm suggested by Hickernell and Jimnez Rugama(2014) [1], but since we are using control variates, several modifications have to be made.

### 3.6 The Algorithm

We now give the algorithm for reliable adaptive QMC with control variates using digital sequence.

---

**Algorithm 1:** Reliable Adaptive QMC with control variates

---

**Data:** function  $f$  and  $\mathbf{H}$ ; value of  $\int_{[0,1]^d} h_j dx = \theta_j$ ; tolerance  $\varepsilon$

**Result:**  $\hat{I}(f)$ ; samples size; optimal  $\beta$

**begin**

```

1    $m, r =$  start numbers,  $x = 2^m$  sobolset points
2   get kappa map( $\tilde{\kappa}$ ) and Walsh coefficients( $\tilde{f}, \tilde{\mathbf{H}}$ ) using algorithm 2
3    $\beta = \tilde{H}\{\tilde{\kappa}[x(a : b)]\} \setminus \tilde{f}\{\tilde{\kappa}[x(a : b)]\}, (a : b) = (2^{m-r-1} : 2^{m-r} - 1)$ 
4    $g = f - \mathbf{H}\beta$ , repeat step 2 on  $g$ 
5    $\tilde{S}_{m-r,m}(g) = \sum_a^b |\tilde{g}\{\tilde{\kappa}[x(a : b)]\}|$ 
6   check whether  $g$  is in the cone
7   if  $a(m, r)\tilde{S}_{m-r,m}(g) \leq \varepsilon$  then
    |   return  $\hat{I}_m(g) = \sum_{i=0}^{2^m-1} f[x(i)] + \theta\beta$ 
    |   return  $\beta, n = 2^m$ 
8   for  $m = m + 1 : mmax$  do
    |    $xnext =$  next  $2^{m-1}$  sobolset points
    |   repeat step 2 on  $[x, xnext]$ 
    |   repeat step 5, 6, 7

```

---

Note that for generating kappa map, i.e. step ??in Algorithm 1, we used an explicit way to generate it.

Another important point need to be mentioned is that in our algorithm, we used an iterative way, which may require recalculating  $\beta$  each time  $m$  increment.

---

**Algorithm 2:** kappa map and discrete Walsh coefficients

---

**Data:** function  $f$ ;  $Y_v^{(m)}$ ;  $m \in \mathbb{N}_0$

**Result:**  $\tilde{\kappa}$ ;  $\tilde{S}_{m-r,m}(f)$

**begin**

**if**  $m = 0$  **then**

$\hat{\mathbf{v}}(0) = 0$

**if**  $m \geq 1$  **then**

**for**  $m : 1 : -1$  **do**

$\hat{\mathbf{v}}_m(\mathbf{k}) = \hat{\mathbf{v}}_{m-1}(\mathbf{k})$

$\hat{\mathbf{v}}_m(\mathbf{k}) = \mathbf{k}, \mathbf{k} = b^{m-1}, \dots, b^m - 1$

**for**  $l = m - 1 : \max(1, m - r) : -1$  **do**

**for**  $k = 1 : b^l - 1$  **do**

$\forall a \in \mathbb{F}_b$ , find  $a$  s.t.  $|Y_{\hat{\mathbf{v}}(k+a*b^l)}^{(m)}| \geq |Y_{\hat{\mathbf{v}}(k+ab^l)}^{(m)}|$

---

## CHAPTER 4

### NUMERICAL EXPERIMENT

Option Pricing has always been a challenging topic in financial mathematics. Here we are going to demonstrate several examples of pricing different options with ccontrol variates. All the options in this chapter are under geometric brownian motion pricing model.

#### 4.1 Asian Option

There are two types of asian options, depends on which types of mean you want to use. For this example we take arithmetic mean asian call option as our target function, whose payoff function is

$$C_T^{Amean} = \max\left(\frac{1}{d} \sum_{j=1}^d S(jT/d) - K, 0\right)$$

Here we use geometric mean asian payoff as the control variates. The payoff function for which is

$$C_T^{gmean} = \max\left(\left[\prod_{j=1}^d S(jT/d)\right]^{\frac{1}{d}} - K, 0\right)$$

And the exact price of geometric mean asian call option under geometric brownian motion is

Table 4.1. Parameter Setup for Up and In Barrier Call Option

S0	K	TimeVector	r	volatility	abstol	reitol
120	130	1/52:1/52:16/52	0.01	0.5	1e-3	0

Figure 4.1 shows decrease rate the walsh coefficients for the target function and control variates in this example.

Table 4.2. Results of cubSobol, cv\_old and cv\_new with Asian Option

Sample Size			Time Cost		
cubSobol	cv_old	cv_new	cubSobol	cv_old	cv_new
65535	8192	9011	0.2783	0.1034	0.0673

## 4.2 Barrier Option

We will take up and in barrier call option as an example. Here is the payoff function for up and in barrier call option.

$$C_T^{U\&I} = (S_T - K)^+ 1_{\{\max S_t \geq \text{Barrier}\}}$$

From the payoff function it is naturally to consider european call option as control variates. Since if we take the barrier same as strike price, then this is just an european call option. Table ?? shows our setup for the barrier option. Payoff function for European option is

$$C_T = \max(S_T - K, 0)$$

Table 4.3. Parameter Setup for Up and In Barrier Call Option

S0	K	TimeVector	r	volatility	abstol	reitol
120	130	1/52:1/52:16/52	0.01	0.5	1e-3	0

We then took three different barrier as listed in table 4.2, then we compared both original cubSobol algorithm and the one with our modification as described in Chapter 4. We can see from the results in table 4.2 that new CV method takes less time than the old one, and both of them are much faster than QMC without CV.


Figure 4.1. Walsh coefficients of  $f$ 

Table 4.4. Results of cubSobol, cv\_old and cv\_new with Barrier Option

Barrier	Sample Size			Time Cost		
	cubSobol	cv_old	cv_new	cubSobol	cv_old	cv_new
140	524288	78643	65535	1.874	0.5016	0.2743
135	524288	5802	6963	1.959	0.0781	0.0519
130	524288	1024	1024	1.876	0.0270	0.0199



## CHAPTER 5

### CONCLUSION

#### 5.1 Discussion

So far there are only few Quasi-Monte Carlo algorithms that can adaptively determine the sample size needed based on integrand values. This is because the estimation of error for QMC is hard. Several studies show that if using Quasi standard error there will be some serious drawbacks [6]. There is also a way using internal replications of i.i.d. Randomized QMC rules, but the number of replications are not known [3].

For control variates the research progress is also limited since its hard to estimate the value of *beta* as we stated in chapter 3.

In our two numerical examples, using the proper control variates gave great results. Also our modified algorithm for the  $\theta$  problem works as expected.

#### 5.2 Future work

Currently we only have method implemented in Sobol sequences. One further possible work is that we can extend this methods on rank-1 lattices [7]. The idea for getting CV coefficients is the same, but due to the different structure for digital sequences, some effort has to be done for adaption of the method.

## BIBLIOGRAPHY

- [1] F. J. Hickernell and L. A. J. Rugama, “Reliable adaptive cubature using digital sequences,” *arXiv preprint arXiv:1410.8615*, 2014.
- [2] A. N. Avramidis and J. R. Wilson, “Integrated variance reduction strategies for simulation,” *Operations Research*, vol. 44, no. 2, pp. 327–346, 1996.
- [3] F. J. Hickernell, C. Lemieux, A. B. Owen, *et al.*, “Control variates for quasi-monte carlo,” *Statistical Science*, vol. 20, no. 1, pp. 1–31, 2005.
- [4] G. Fishman, *Monte Carlo: concepts, algorithms, and applications*. Springer Science & Business Media, 2013.
- [5] H. Niederreiter, *Quasi-Monte Carlo Methods*. Wiley Online Library, 2010.
- [6] A. B. Owen, “On the warnock-halton quasi-standard error,” *Monte Carlo Methods and Applications mcma*, vol. 12, no. 1, pp. 47–54, 2006.
- [7] L. A. J. Rugama and F. J. Hickernell, “Adaptive multidimensional integration based on rank-1 lattices,” *arXiv preprint arXiv:1411.1966*, 2014.