



Digital Systems and Computer Architecture

Session 2.4


Module 2

Mouli Sankaran

Theorems in Boolean Algebra


Session 2.4: Focus

- Introduction to Boolean Algebra
- Ordinary Algebra Vs Boolean Algebra
- Boolean Expressions
 - Variables, literals and terms
 - Complement and Dual
- DeMorgan's Theorems



Introduction to Boolean Algebra

George Boole




- **George Boole** (2 November 1815 – 8 December 1864)
- He was an English mathematician, educator, philosopher and logician.
- He worked in the fields of differential equations and algebraic logic, and is best known as the author of **The Laws of Thought (1854)** which contains **Boolean algebra**.
- Boolean logic is credited with laying the foundations for the information age.

Boolean Algebra

- Founded by **George Boole**, in **1854**.
- Boolean algebra is **mathematics of logic**, a systematic way of expressing and analyzing the operations of **logic circuits**
- It is one of the **most basic tools** available to the **logic designer**
 - It can be effectively **used** for **simplification** of **complex logic expressions**
- Now, let us have a closer look at the different **postulates** and **theorems** of **Boolean algebra**
- Their applications in **minimizing Boolean expressions**

Introduction to Boolean Algebra

- **Boolean algebra**, quite interestingly, is **simpler** than **ordinary algebra**.
- It is also composed of a set of **symbols** and a set of **rules to manipulate** these symbols



Ordinary Algebra Vs Boolean Algebra


Ordinary Algebra Vs Boolean Algebra

[illegible]

Ordinary Algebra Vs Boolean Algebra

Repeated again for the pdf copy

Ordinary Algebra	Boolean Algebra
Letter symbols can take on any number of values including infinity	Letter symbols can take on either of two values , that is, 0 and 1
Values assigned to a variable have a numerical significance	Values assigned to a variable have a logical significance
‘.’ and ‘+’ are respectively the signs of multiplication and addition	‘.’ means an AND operation and ‘+’ means an OR operation
A+B is read as A plus B	A+B is read as A OR B Basic logic operations are AND, OR and NOT
	Captures both logic operations and set operations such as intersection, union and complement



Boolean Expressions

Variables, Literals and Terms

- **Variables** are different symbols in a **Boolean expression**
- They may take on the values '0' or '1'
- **Complement** of a variable is **not** considered as a **separate variable**
- Each occurrence of a **variable** or its **complement** is called a **literal**
- A **term** is the **expression** formed by **literals** and **operations** at **one level**.
- Each **term** requires a **gate** and each **variable** within the term designates an **input** to the **gate**

$$\overline{A} + A.B$$

Quiz:

Variables : 2

Literals : 3


Terms : 2

Quiz 1: Variables, Literals and Terms

- How many **Variables**, **literals** and **Terms** are there in these **two Boolean expressions**?
- How many **gates** are needed to construct them?

Boolean Expressions	Variables	Literals	Terms	Gates
$F_2 = x'y'z + x'yz + xy'$	3	8	3	NOT : 2 AND : 3 OR : 1
$F_2 = xy' + x'z$	3	4	2	NOT : 2 AND : 2 OR : 1

Note: Assume both 2-input and 3-input gates are available to construct these expressions, without minimization.



Equivalent and Complement of Boolean Expressions


Equivalent and Complement of Boolean Expressions

- **Two given Boolean expressions** are said to be **equivalent**
 - If **one** of them **equals '1'** only when the **other equals '1'** and also **one equals '0'** only when the **other equals '0'**
- They are said to be the **complement** of each other
 - If **one** expression **equals '1'** only when the **other equals '0'**, and **vice versa**
- The **complement** of a given **Boolean expression** is obtained by
 - **Complementing each literal,**
 - **Changing** all **'.'** to **'+'** and all **'+'** to **'.'**, all **0s** to **1s** and all **1s** to **0s**.

Quiz 2: Find Complements

- Find the **complement** of below **Boolean expressions**:

Boolean Expression	Complement
$\overline{A}.B + A.\overline{B}$	$(A + \overline{B}).(\overline{A} + B)$
$(A + B).(\overline{A} + \overline{B})$	$\overline{A}.\overline{B} + A.B$
$[(A.\overline{B} + \overline{C}).D + \overline{E}].F$	$[(\overline{A} + B).C + \overline{D}].E + \overline{F}$



Dual of Boolean Expressions

Dual of Boolean Expressions

The **dual** of a Boolean expression is obtained by **replacing**

- All **‘.’** operations with **‘+’** operations
- All **‘+’** operations with **‘.’** operations,
- All **0s** with **1s** and all **1s** with **0s** and
- Leaving all **literals unchanged**

Quiz 3: Find Dual

- Find the **dual** of below **Boolean expressions**:

Boolean Expression	Dual
$\overline{A}.B + A.\overline{B}$	$(\overline{A} + B).(A + \overline{B})$
$(A + B).(\overline{A} + \overline{B})$	$A.B + \overline{A}.\overline{B}$
$A.\overline{B} + B.\overline{C} + C.\overline{D}$	$(A + \overline{B}).(B + \overline{C}).(C + \overline{D})$

Quiz 4: Find Dual and Complement

- Find the **dual** and **Complement** of below **Boolean expression**:

Boolean Expression	$[(A.\overline{B} + \overline{C}).D + \overline{E}].F$
Dual	$[(A + \overline{B}).\overline{C} + D].\overline{E} + F$
Complement	$[(\overline{A} + B).C + \overline{D}].E + \overline{F}$

What is Dual of a Boolean Expression?

The principle of **duality** pronounces that given an **expression** which is always **valid** in boolean algebra, the **dual expression** is **also** always **valid**

- For example:
 - $A(B + C) = A \cdot B + A \cdot C$
 - It's **dual** is also **valid**
 - $A + (B \cdot C) = (A + B) \cdot (A + C)$

Rules of Boolean Algebra

- **A, B or C** can represent a single variable or a combination of variables

$$1. A + 0 = A$$

$$2. A + 1 = 1$$

$$3. A \cdot 0 = 0$$

$$4. A \cdot 1 = A$$

$$5. A + A = A$$

$$6. A + \bar{A} = 1$$

$$7. A \cdot A = A$$

$$8. A \cdot \bar{A} = 0$$

$$9. \bar{\bar{A}} = A$$

$$10. A + AB = A$$


$$11. A + \bar{A}B = A + B$$

$$12. (A + B)(A + C) = A + BC$$

- These **12 rules** are useful in **simplifying** Boolean expressions


Rule 1

- $A + 0 = A$




Rule 2

- $A + 1 = 1$




Rule 3

- $A \cdot 0 = 0$




Rule 4

- $A \cdot 1 = A$



Rule 5


- $A + A = A$ (Idempotent or Identity Law)



- Valid for any number of the same input:
- $A + A + A + + + A = A$


Rule 6

- $A + \bar{A} = 1$



Rule 7


- **$A \cdot A = A$ (Idempotent or Identity Law)**



- Valid for **any number** of the **same input**:
- **$A \cdot A \cdot A \dots A = A$**


Rule 8

- $A \cdot \bar{A} = 0$



Rule 9

- $\overline{\overline{A}} = A$ (Involution law)




Rule 10

- $A + AB = A$ (Absorption or Redundancy Law)
- $A + AB = A(1 + B)$: Factoring (Distributive law)
- $= A \cdot 1$: By Rule 2, $(1 + B) = 1$
- $= A$: By Rule 4, $A \cdot 1 = A$
- $A + AB = A$

<i>A</i>	<i>B</i>	<i>AB</i>	<i>A + AB</i>
0	0	0	0
0	1	0	0
1	0	0	1
1	1	1	1

↑ equal ↑



Rule 11


- $A + \bar{A}B = A + B$

$$A + \bar{A}B = (A + AB) + \bar{A}B$$

Rule 10: $A = A + AB$

A	B	$\bar{A}B$	$A + \bar{A}B$	$A + B$
0	0	0	0	0
0	1	1	1	1
1	0	0	1	1
1	1	0	1	1

↑ equal ↑



Rule 11

- $A + \bar{A}B = A + B$**

$$\begin{aligned}
 A + \bar{A}B &= (A + AB) + \bar{A}B \\
 &= (AA + AB) + \bar{A}B \\
 &= AA + AB + A\bar{A} + \bar{A}B \\
 &= (A + \bar{A})(A + B) \\
 &= 1 \cdot (A + B) \\
 &= A + B
 \end{aligned}$$

Rule 10: $A = A + AB$

Rule 7: $A = AA$

Rule 8: adding $A\bar{A} = 0$


Factoring

Rule 6: $A + \bar{A} = 1$

Rule 4: drop the 1

A	B	$\bar{A}B$	$A + \bar{A}B$	$A + B$
0	0	0	0	0
0	1	1	1	1
1	0	0	1	1
1	1	0	1	1

↑ equal ↑



Rule 12

- $(A + B) \cdot (A + C) = A + BC$

$$(A + B)(A + C) = AA + AC + AB + BC \quad \text{Distributive law}$$

A	B	C	A + B	A + C	(A + B)(A + C)	BC	A + BC
0	0	0	0	0	0	0	0
0	0	1	0	1	0	0	0
0	1	0	1	0	0	0	0
0	1	1	1	1	1	1	1
1	0	0	1	1	1	0	1
1	0	1	1	1	1	0	1
1	1	0	1	1	1	0	1
1	1	1	1	1	1	1	1

↑ equal ↑

Rule 12

Repeated again for the pdf copy

- $(A + B) \cdot (A + C) = A + BC$

$$(A + B)(A + C) = AA + AC + AB + BC \quad \text{Distributive law}$$

$$= A + AC + AB + BC \quad \text{Rule 7: } AA = A$$

$$= A(1 + C) + AB + BC \quad \text{Factoring (distributive law)}$$

$$= A \cdot 1 + AB + BC \quad \text{Rule 2: } 1 + C = 1$$


$$= A(1 + B) + BC \quad \text{Factoring (distributive law)}$$


$$= A \cdot 1 + BC \quad \text{Rule 2: } 1 + B = 1$$

$$= A + BC \quad \text{Rule 4: } A \cdot 1 = A$$

A	B	C	A + B	A + C	(A + B)(A + C)	BC	A + BC
0	0	0	0	0	0	0	0
0	0	1	0	1	0	0	0
0	1	0	1	0	0	0	0
0	1	1	1	1	1	1	1
1	0	0	1	1	1	0	1
1	0	1	1	1	1	0	1
1	1	0	1	1	1	0	1
1	1	1	1	1	1	1	1


↑ equal ↑





DeMorgon's Theorems

Augustus De Morgan



Augustus De Morgan,
(born June 27, 1806, Madura, India)
English mathematician and logician
whose major **contributions** to the
study of logic include the formulation
of **De Morgan's laws**

DeMorgan, a mathematician who knew Boole,
proposed two important theorems of Boolean Algebra

DeMorgan's Theorem 1


- **Theorem 1:**
- The **complement** of a **sum of variables** is **equal** to the **product of the complements** of the variables

$$\overline{X + Y} = \overline{X} \overline{Y}$$

- It can also be stated as:
- The **complement** of two or more **ORed variables** is **equivalent** to the **AND** of the **complements** of the **individual variables**.
- Valid for any number of variables:

$$\overline{[X_1 + X_2 + X_3 + \dots + X_n]} = \overline{X_1} \cdot \overline{X_2} \cdot \overline{X_3} \cdot \dots \cdot \overline{X_n}$$

DeMorgan's Theorem 1



Inputs		Output	
X	Y	$\overline{X + Y}$	$\overline{\overline{X} \overline{Y}}$
0	0	1	1
0	1	0	0
1	0	0	0
1	1	0	0

DeMorgan's Theorem 2


- **Theorem 2:**
- The **complement** of a **product of variables** is **equal** to the **sum of the complements** of the variables

$$\overline{XY} = \overline{X} + \overline{Y}$$

- It can also be stated as:
- The **complement** of two or more **ANDed variables** is **equivalent** to the **OR** of the **complements** of the **individual variables**.
- Valid for any number of variables:

$$\overline{[X_1 \cdot X_2 \cdot X_3 \cdot \dots \cdot X_n]} = [\overline{X_1} + \overline{X_2} + \overline{X_3} + \dots + \overline{X_n}]$$

DeMorgon's Theorem 2



Inputs		Output	
X	Y	\overline{XY}	$\overline{X} + \overline{Y}$
0	0	1	1
0	1	1	1
1	0	1	1
1	1	0	0

Home work:
Proof for Theorem 2

Session 2.4: Summary

- Introduction to Boolean Algebra
- Ordinary Algebra Vs Boolean Algebra
- Boolean Expressions
 - Variables, literals and terms
 - Complement and Dual
- DeMorgan's Theorems