

AI5002 - Assignment 1

Tuhin Dutta
ai21mtech02002

Download code and LaTeX from below hyperlinks

1. [Code](#)
2. [LaTeX](#)

Problem 6.1.3

Plot the CDF and PDF of

$$A = \sqrt{V} \quad (0.0.1)$$

Solution

We can write $A = \sqrt{V}$ as

$$A = \sqrt{X_1^2 + X_2^2} \quad (0.0.2)$$

since it is given $X_1 \sim N(0, 1)$, $X_2 \sim N(0, 1)$ and $V = X_1^2 + X_2^2$.

Since joint distribution is not mentioned so we assume X_1 and X_2 to be independent otherwise the distribution of A would be unknown.

By definition, the distribution of A is Chi with two degrees of freedom or Rayleigh.

The CDF and PDF plot is as shown in Fig 1.1 and Fig 1.2.




Fig 1.1: Cumulative distribution function




Fig 1.2: Probability density function

Problem 6.1.4

Find an expression for $F_A(x)$ using the definition. Plot this expression and compare with the result of problem 6.1.3

Solution

Given $X_1 \sim N(0, 1)$, $X_2 \sim N(0, 1)$, and $V = X_1^2 + X_2^2$.

So we can write A as

$$A = \sqrt{V}$$

$$\text{or, } A = \sqrt{X_1^2 + X_2^2}$$

Since joint distribution is not mentioned so we assume X_1 and X_2 to be independent otherwise the distribution of A would be unknown.

By definition, the distribution of A is Chi with two degrees of freedom or Rayleigh.

The CDF given by $F_A(x)$ can be written as

$$= P(A \leq x)$$

$$= P(\sqrt{X_1^2 + X_2^2} \leq x)$$

$$= \iint_{\sqrt{X_1^2 + X_2^2} \leq x} F(x_1, x_2) dx_1 dx_2$$

$$= \iint_{\sqrt{X_1^2 + X_2^2} \leq x} F(x_1).F(x_2) dx_1 dx_2$$

$$= \iint_{\sqrt{X_1^2 + X_2^2} \leq x} \frac{1}{\sqrt{2\pi\sigma^2}} \cdot \frac{1}{\sqrt{2\pi\sigma^2}} \cdot e^{-\frac{x_1^2}{2\sigma^2}} \cdot e^{-\frac{x_2^2}{2\sigma^2}} dx_1 dx_2$$

$$= \iint_{\sqrt{X_1^2 + X_2^2} \leq x} \frac{1}{2\pi\sigma^2} \cdot e^{-\frac{(x_1^2 + x_2^2)}{2\sigma^2}} dx_1 dx_2 \quad (1)$$

Now, we perform a transformation of variables to simplify solution.

Let $x_1 = r \cos \theta$ and, $x_2 = r \sin \theta$.

$$\Rightarrow \sqrt{x_1^2 + x_2^2} = r$$

$$\text{Also, } \theta = \tan^{-1} \frac{x_2}{x_1}$$

Since we're using transformation of two variables, we use Jacobian matrix here

$$J_{x_1, x_2} = \begin{bmatrix} \frac{\partial x_1}{\partial r} & \frac{\partial x_1}{\partial \theta} \\ \frac{\partial x_2}{\partial r} & \frac{\partial x_2}{\partial \theta} \end{bmatrix} = \begin{bmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{bmatrix}$$

$$\det(J_{x_1, x_2}) = r \cos^2 \theta + r \sin^2 \theta = r$$

$$\text{Thus, } F(r, \theta) = r \cdot F(x_1, x_2)$$

where (x_1, x_2) are written in terms of (r, θ) .

Now, we can write (1) as

$$\iint_{r \leq x} \frac{1}{2\pi\sigma^2} \cdot e^{-\frac{r^2}{2\sigma^2}} \cdot r dr d\theta$$

or,

$$\int_{\theta=0}^{2\pi} \int_{r=0}^x \frac{1}{2\pi\sigma^2} \cdot e^{-\frac{r^2}{2\sigma^2}} \cdot r dr d\theta \quad (2)$$

$$\text{Let } u = -\frac{r^2}{2\sigma^2}$$

$$du = -\frac{2r}{2\sigma^2} dr$$

$$r dr = -\sigma^2 du$$

$$\text{where } 0 \leq r \leq x \text{ and } 0 \leq u \leq -\frac{x^2}{2\sigma^2}$$

Substituting u and r dr in (2), we get

$$\begin{aligned} & \int_{\theta=0}^{2\pi} \int_{u=0}^{-\frac{x^2}{2\sigma^2}} \frac{1}{2\pi\sigma^2} \cdot e^u \cdot (-\sigma^2) du d\theta \\ &= \int_{\theta=0}^{2\pi} -\frac{1}{2\pi} \int_{u=0}^{-\frac{x^2}{2\sigma^2}} e^u du d\theta = -\frac{1}{2\pi} \int_{\theta=0}^{2\pi} (e^{-\frac{x^2}{2\sigma^2}} - 1) d\theta \end{aligned}$$

Thus the CDF is derived as

$$F_A(x) = 1 - \exp(-x^2/2\sigma^2)$$




Fig 3: CDF Actual Vs Simulation

Problem 6.1.5

Find an expression for $p_A(x)$ using the definition.

Solution

The Pdf can be derived by differentiating the CDF expression from the previous problem 6.1.4

$$\begin{aligned}
 p_A(x) &= \frac{d}{dx} F_A(x) \\
 &= -e^{-\frac{x^2}{2\sigma^2}} \cdot \left(-\frac{2x}{2\sigma^2}\right) \\
 &= e^{-\frac{x^2}{2\sigma^2}} \cdot \frac{x}{\sigma^2}
 \end{aligned}$$




Fig 4: Pdf Actual Vs Simulation