

## Seminar 3

Octavian Custura

November 2020

8. Three contestants participate in a trivia game show. Their probabilities of answering a question correctly are 0.8, 0.9 and 0.75, respectively. If 10 questions are asked and every contestant answers every question, find the probability that one contestant (any one) answers exactly 7 questions correctly, while the other two give any other number of correct answers. (event A)?

*Proof.* Denote by  $A_1$ ,  $A_2$  and  $A_3$  the probability that the contestant no 1, 2, respectively 3 answers exactly 7 questions right. It can be seen that the 3 probabilities are computed using a **Binomial model**, since every one of them has equal probability for every iteration. We have

$$P(A_1) = C_n^k \cdot 0.8^7 \cdot 0.2^3 = 0.2$$

$$P(A_2) = C_n^k \cdot 0.9^7 \cdot 0.1^3 = 0.05$$

$$P(A_3) = C_n^k \cdot 0.75^7 \cdot 0.25^3 = 0.25$$

In order to compute  $P(A)$ , we need exactly one value from the 3 probabilities  $A_1$ ,  $A_2$  and  $A_3$ , thus using a **Poisson Model**.

$P(A)$  = the coefficient of  $x^1$  in the polynomial expansion of  $(P(A_1) \cdot x + 1 - P(A_1)) \cdot (P(A_2) \cdot x + 1 - P(A_2)) \cdot (P(A_3) \cdot x + 1 - P(A_3))$

$$(0.2 \cdot x + 0.8) \cdot (0.05 \cdot x + 0.95) \cdot (0.25 \cdot x + 0.75) = 0.0025 \cdot x^3 + 0.065 \cdot x^2 + 0.3625 \cdot x + 0.57 \quad (1)$$

We can see that the coefficient of  $x^1$  is  $0.3625 = P(A)$ , which is our final probability.  $\square$

9. In a department store at the mall, black and brown gloves are on sale. There are  $N$  (identical) pairs of black gloves and  $N$  (identical) pairs of brown gloves. If  $N$  customers come in, one at a time and randomly choose and buy 2 pairs each, find the probability of event A: each customer buys 2 pairs of different colors (one black and one brown).

*Proof.* There are  $N$  trials, where we need  $N$  successes, and every trial has a different probability. Thus, we need a **Poisson Model**. Denote by  $P(A_K)$  the

probability that the  $k$ th person chooses gloves of different colours, knowing that the first  $k - 1$  people took gloves of different colours.

$$P(A_1) = \frac{N}{2 \cdot N - 1}, P(A_2) = \frac{N - 1}{2 \cdot N - 3} \dots P(A_i) = \frac{N - i + 1}{2 \cdot N - 2 \cdot i + 1} \dots P(A_N) = \frac{1}{1}$$

Thus, we observe that the probability of  $N$  successes in  $N$  trials is the coefficient of  $x^N$  in the polynomial expansion of  $\prod_{i=1}^N \left( \frac{N - i + 1}{2 \cdot N - 2 \cdot i + 1} \cdot x + \frac{N - i}{2 \cdot N - 2 \cdot i + 1} \right)$ .

We can observe that the coefficient of  $x^N$  is the product of the coefficients of  $x$  in the above product, so

$$P(A) = \prod_{i=1}^N \frac{N - i + 1}{2 \cdot N - 2 \cdot i + 1} = \frac{N!}{(2 \cdot N - 1) \cdot (2 \cdot N - 3) \cdot \dots \cdot 3 \cdot 1}$$

Denote  $Q = 2 \cdot 4 \cdot 6 \dots 2n$ .  $Q = (1 \cdot 2)(2 \cdot 2)(3 \cdot 2) \dots (n \cdot 2) = n! \cdot 2^n$ . Also,  $(2n)! = (1 \cdot 3 \cdot 5 \dots (2n - 1)) \cdot (2 \cdot 4 \cdot 6 \dots 2n) = (1 \cdot 3 \cdot 5 \dots (2n - 1)) \cdot Q$ . So  $(1 \cdot 3 \cdot 5 \dots (2n - 1)) = \frac{(2n)!}{2^n \cdot n!}$ . In conclusion,

$$P(A) = \frac{N!^2 \cdot 2^N}{(2N)!}$$

□