

# Assignment 4: Data Wrangling

Taylor Coleman

Spring 2023

## OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on Data Wrangling

## Directions

1. Rename this file `<FirstLast>_A04_DataWrangling.Rmd` (replacing `<FirstLast>` with your first and last name).
2. Change “Student Name” on line 3 (above) with your name.
3. Work through the steps, **creating code and output** that fulfill each instruction.
4. Be sure to **answer the questions** in this assignment document.
5. When you have completed the assignment, **Knit** the text and code into a single PDF file.

The completed exercise is due on Friday, Feb 20th @ 5:00pm.

## Set up your session

- 1a. Load the `tidyverse`, `lubridate`, and `here` packages into your session.
  - 1b. Check your working directory.
  - 1c. Read in all four raw data files associated with the EPA Air dataset, being sure to set string columns to be read in as factors. See the README file for the EPA air datasets for more information (especially if you have not worked with air quality data previously).
2. Apply the `glimpse()` function to reveal the dimensions, column names, and structure of each dataset.

```
# 1a Installing 'tidyverse,' 'lubridate,' and 'here' packages
install.packages("tidyverse")
install.packages("lubridate")
install.packages("here")

# 1b checking working directory
getwd()
```

```
## [1] "/home/guest/ENVR872-RepositoryClone"
```

```

# 1c Reading in EPA raw data files
EPA.O3.NC.2018 <- read.csv("/home/guest/ENVR872-RepositoryClone/Data/Raw/EPAair_O3_NC2018_raw.csv",
  stringsAsFactors = TRUE)
EPA.O3.NC.2019 <- read.csv("/home/guest/ENVR872-RepositoryClone/Data/Raw/EPAair_O3_NC2019_raw.csv",
  stringsAsFactors = TRUE)
EPA.PM25.NC.2018 <- read.csv("/home/guest/ENVR872-RepositoryClone/Data/Raw/EPAair_PM25_NC2018_raw.csv",
  stringsAsFactors = TRUE)
EPA.PM25.NC.2019 <- read.csv("/home/guest/ENVR872-RepositoryClone/Data/Raw/EPAair_PM25_NC2019_raw.csv",
  stringsAsFactors = TRUE)

# 2 Applying 'glimpse' function to explore data
library(dplyr)
glimpse(EPA.O3.NC.2018)

```

```

## Rows: 9,737
## Columns: 20
## $ Date                <fct> 03/01/2018, 03/02/2018, 03/03/201~
## $ Source              <fct> AQS, AQS, AQS, AQS, AQS, AQS, AQS~
## $ Site.ID             <int> 370030005, 370030005, 370030005, ~
## $ POC                 <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ Daily.Max.8.hour.Ozone.Concentration <dbl> 0.043, 0.046, 0.047, 0.049, 0.047~
## $ UNITS               <fct> ppm, ppm, ppm, ppm, ppm, ppm, ppm~
## $ DAILY_AQI_VALUE     <int> 40, 43, 44, 45, 44, 28, 33, 41, 4~
## $ Site.Name           <fct> Taylorsville Liledoun, Taylorsvil~
## $ DAILY_OBS_COUNT     <int> 17, 17, 17, 17, 17, 17, 17, 17, 1~
## $ PERCENT_COMPLETE    <dbl> 100, 100, 100, 100, 100, 100, 100~
## $ AQS_PARAMETER_CODE  <int> 44201, 44201, 44201, 44201, 44201~
## $ AQS_PARAMETER_DESC  <fct> Ozone, Ozone, Ozone, Ozone, Ozone~
## $ CBSA_CODE           <int> 25860, 25860, 25860, 25860, 25860~
## $ CBSA_NAME           <fct> "Hickory-Lenoir-Morganton, NC", "~
## $ STATE_CODE          <int> 37, 37, 37, 37, 37, 37, 37, 37, 3~
## $ STATE               <fct> North Carolina, North Carolina, N~
## $ COUNTY_CODE         <int> 3, 3, 3, 3, 3, 3, 3, 3, 3, 3, ~
## $ COUNTY              <fct> Alexander, Alexander, Alexander, ~
## $ SITE_LATITUDE       <dbl> 35.9138, 35.9138, 35.9138, 35.913~
## $ SITE_LONGITUDE      <dbl> -81.191, -81.191, -81.191, -81.19~

```

```
glimpse(EPA.O3.NC.2019)
```

```

## Rows: 10,592
## Columns: 20
## $ Date                <fct> 01/01/2019, 01/02/2019, 01/03/201~
## $ Source              <fct> AirNow, AirNow, AirNow, AirNow, A~
## $ Site.ID             <int> 370030005, 370030005, 370030005, ~
## $ POC                 <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ Daily.Max.8.hour.Ozone.Concentration <dbl> 0.029, 0.018, 0.016, 0.022, 0.037~
## $ UNITS               <fct> ppm, ppm, ppm, ppm, ppm, ppm, ppm~
## $ DAILY_AQI_VALUE     <int> 27, 17, 15, 20, 34, 34, 27, 35, 3~
## $ Site.Name           <fct> Taylorsville Liledoun, Taylorsvil~
## $ DAILY_OBS_COUNT     <int> 24, 24, 24, 24, 24, 24, 24, 24, 2~
## $ PERCENT_COMPLETE    <dbl> 100, 100, 100, 100, 100, 100, 100~
## $ AQS_PARAMETER_CODE  <int> 44201, 44201, 44201, 44201, 44201~

```

```
## $ AQS_PARAMETER_DESC      <fct> Ozone, Ozone, Ozone, Ozone, Ozone~
## $ CBSA_CODE               <int> 25860, 25860, 25860, 25860, 25860~
## $ CBSA_NAME               <fct> "Hickory-Lenoir-Morganton, NC", "~
## $ STATE_CODE              <int> 37, 37, 37, 37, 37, 37, 37, 37, 3~
## $ STATE                   <fct> North Carolina, North Carolina, N~
## $ COUNTY_CODE             <int> 3, 3, 3, 3, 3, 3, 3, 3, 3, 3, ~
## $ COUNTY                  <fct> Alexander, Alexander, Alexander, ~
## $ SITE_LATITUDE           <dbl> 35.9138, 35.9138, 35.9138, 35.913~
## $ SITE_LONGITUDE          <dbl> -81.191, -81.191, -81.191, -81.19~
```

```
glimpse(EPA.PM25.NC.2018)
```

```
## Rows: 8,983
## Columns: 20
## $ Date                   <fct> 01/02/2018, 01/05/2018, 01/08/2018, 01/~
## $ Source                 <fct> AQS, AQS, AQS, AQS, AQS, AQS, AQS, AQS,~
## $ Site.ID                <int> 370110002, 370110002, 370110002, 370110~
## $ POC                    <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ Daily.Mean.PM2.5.Concentration <dbl> 2.9, 3.7, 5.3, 0.8, 2.5, 4.5, 1.8, 2.5,~
## $ UNITS                  <fct> ug/m3 LC, ug/m3 LC, ug/m3 LC, ug/m3 LC,~
## $ DAILY_AQI_VALUE        <int> 12, 15, 22, 3, 10, 19, 8, 10, 18, 7, 24~
## $ Site.Name              <fct> Linville Falls, Linville Falls, Linvill~
## $ DAILY_OBS_COUNT        <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ PERCENT_COMPLETE       <dbl> 100, 100, 100, 100, 100, 100, 100, 100,~
## $ AQS_PARAMETER_CODE     <int> 88502, 88502, 88502, 88502, 88502, 8850~
## $ AQS_PARAMETER_DESC     <fct> Acceptable PM2.5 AQI & Speciation Mass,~
## $ CBSA_CODE              <int> NA, NA, NA, NA, NA, NA, NA, NA, NA, NA,~
## $ CBSA_NAME              <fct> "", "", "", "", "", "", "", "", "", "",~
## $ STATE_CODE             <int> 37, 37, 37, 37, 37, 37, 37, 37, 37, 37,~
## $ STATE                  <fct> North Carolina, North Carolina, North C~
## $ COUNTY_CODE            <int> 11, 11, 11, 11, 11, 11, 11, 11, 11, 11,~
## $ COUNTY                 <fct> Avery, Avery, Avery, Avery, Avery, Aver~
## $ SITE_LATITUDE          <dbl> 35.97235, 35.97235, 35.97235, 35.97235,~
## $ SITE_LONGITUDE         <dbl> -81.93307, -81.93307, -81.93307, -81.93~
```

```
glimpse(EPA.PM25.NC.2019)
```

```
## Rows: 8,581
## Columns: 20
## $ Date                   <fct> 01/03/2019, 01/06/2019, 01/09/2019, 01/~
## $ Source                 <fct> AQS, AQS, AQS, AQS, AQS, AQS, AQS, AQS,~
## $ Site.ID                <int> 370110002, 370110002, 370110002, 370110~
## $ POC                    <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ Daily.Mean.PM2.5.Concentration <dbl> 1.6, 1.0, 1.3, 6.3, 2.6, 1.2, 1.5, 1.5,~
## $ UNITS                  <fct> ug/m3 LC, ug/m3 LC, ug/m3 LC, ug/m3 LC,~
## $ DAILY_AQI_VALUE        <int> 7, 4, 5, 26, 11, 5, 6, 6, 15, 7, 14, 20~
## $ Site.Name              <fct> Linville Falls, Linville Falls, Linvill~
## $ DAILY_OBS_COUNT        <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ PERCENT_COMPLETE       <dbl> 100, 100, 100, 100, 100, 100, 100, 100,~
## $ AQS_PARAMETER_CODE     <int> 88502, 88502, 88502, 88502, 88502, 8850~
## $ AQS_PARAMETER_DESC     <fct> Acceptable PM2.5 AQI & Speciation Mass,~
## $ CBSA_CODE              <int> NA, NA, NA, NA, NA, NA, NA, NA, NA, NA,~
## $ CBSA_NAME              <fct> "", "", "", "", "", "", "", "", "", "",~
```

```
## $ STATE_CODE          <int> 37, 37, 37, 37, 37, 37, 37, 37, 37, 37,~
## $ STATE              <fct> North Carolina, North Carolina, North C~
## $ COUNTY_CODE        <int> 11, 11, 11, 11, 11, 11, 11, 11, 11, 11,~
## $ COUNTY             <fct> Avery, Avery, Avery, Avery, Avery, Aver~
## $ SITE_LATITUDE       <dbl> 35.97235, 35.97235, 35.97235, 35.97235,~
## $ SITE_LONGITUDE      <dbl> -81.93307, -81.93307, -81.93307, -81.93~
```

## Wrangle individual datasets to create processed files.

3. Change date columns to be date objects.
4. Select the following columns: Date, DAILY\_AQI\_VALUE, Site.Name, AQS\_PARAMETER\_DESC, COUNTY, SITE\_LATITUDE, SITE\_LONGITUDE
5. For the PM2.5 datasets, fill all cells in AQS\_PARAMETER\_DESC with “PM2.5” (all cells in this column should be identical).
6. Save all four processed datasets in the Processed folder. Use the same file names as the raw files but replace “raw” with “processed”.

```
# 3 Changing date 'factors' to date objects for each dataset
EPA.03.NC.2018$Date <- as.Date(EPA.03.NC.2018$Date, format = "%m/%d/%Y")
EPA.03.NC.2019$Date <- as.Date(EPA.03.NC.2019$Date, format = "%m/%d/%Y")
EPA.PM25.NC.2018$Date <- as.Date(EPA.PM25.NC.2018$Date, format = "%m/%d/%Y")
EPA.PM25.NC.2019$Date <- as.Date(EPA.PM25.NC.2019$Date, format = "%m/%d/%Y")

# 4 Selecting columns from each dataset
EPA.03.2018.subset <- select(EPA.03.NC.2018, Date, DAILY_AQI_VALUE:Site.Name, AQS_PARAMETER_DESC,
  COUNTY, SITE_LATITUDE:SITE_LONGITUDE)
EPA.03.2019.subset <- select(EPA.03.NC.2019, Date, DAILY_AQI_VALUE:Site.Name, AQS_PARAMETER_DESC,
  COUNTY, SITE_LATITUDE:SITE_LONGITUDE)
EPA.PM25.2018.subset <- select(EPA.PM25.NC.2018, Date, DAILY_AQI_VALUE:Site.Name,
  AQS_PARAMETER_DESC, COUNTY, SITE_LATITUDE:SITE_LONGITUDE)
EPA.PM25.2019.subset <- select(EPA.PM25.NC.2019, Date, DAILY_AQI_VALUE:Site.Name,
  AQS_PARAMETER_DESC, COUNTY, SITE_LATITUDE:SITE_LONGITUDE)

# 5 Filling-in cells of AQS_PARAMETER_DESC column from PM 2.5 datasets with
# 'PM2.5'
EPA.PM25.2018.subset <- mutate(EPA.PM25.2018.subset, AQS_PARAMETER_DESC = "PM2.5")
EPA.PM25.2019.subset <- mutate(EPA.03.2019.subset, AQS_PARAMETER_DESC = "PM2.5")

# 6 Saving processed datasets
write.csv(EPA.03.2018.subset, row.names = FALSE, file = "Data/Processed/EPAair_03_NC2018_processed.csv")
write.csv(EPA.03.2019.subset, row.names = FALSE, file = "Data/Processed/EPAair_03_NC2019_processed.csv")
write.csv(EPA.PM25.2018.subset, row.names = FALSE, file = "Data/Processed/EPAair_PM25_NC2018_processed.csv")
write.csv(EPA.PM25.2019.subset, row.names = FALSE, file = "Data/Processed/EPAair_PM25_NC2019_processed.csv")
```

## Combine datasets

7. Combine the four datasets with `rbind`. Make sure your column names are identical prior to running this code.
8. Wrangle your new dataset with a pipe function (`%>%`) so that it fills the following conditions:

- Include all sites that the four data frames have in common: “Linville Falls”, “Durham Armory”, “Leggett”, “Hattie Avenue”, “Clemmons Middle”, “Mendenhall School”, “Frying Pan Mountain”, “West Johnston Co.”, “Garinger High School”, “Castle Hayne”, “Pitt Agri. Center”, “Bryson City”, “Millbrook School” (the function `intersect` can figure out common factor levels - but it will include sites with missing site information...)
  - Some sites have multiple measurements per day. Use the split-apply-combine strategy to generate daily means: group by date, site name, AQS parameter, and county. Take the mean of the AQI value, latitude, and longitude.
  - Add columns for “Month” and “Year” by parsing your “Date” column (hint: `lubridate` package)
  - Hint: the dimensions of this dataset should be 14,752 x 9.
9. Spread your datasets such that AQI values for ozone and PM2.5 are in separate columns. Each location on a specific date should now occupy only one row.
  10. Call up the dimensions of your new tidy dataset.
  11. Save your processed dataset with the following file name: “EPAair\_O3\_PM25\_NC1819\_Processed.csv”

#### *# 7 Using rbind to combine the 4 datasets*

```
EPAair_O3_PM25_NC1819 <- rbind(EPA.O3.2018.subset, EPA.O3.2019.subset, EPA.PM25.2018.subset,
                                EPA.PM25.2019.subset)
```

#### *# Extra (unnecessary) step: Converting Date column from factor to date variable*

```
EPAair_O3_PM25_NC1819$Date <- as.Date(EPAair_O3_PM25_NC1819$Date, format = "%m/%d/%Y")
```

#### *# 8 Using the pipe function to wrangle new combined dataset filtering common sites, generating daily means for AQIs, latitude, and longitude values, and adding columns for 'month' and 'year'*

```
library(lubridate)
```

```
##
```

```
## Attaching package: 'lubridate'
```

```
## The following objects are masked from 'package:base':
```

```
##
```

```
##      date, intersect, setdiff, union
```

```
EPAair_O3_PM25_NC1819_Wrangled <- EPAair_O3_PM25_NC1819 %>%
  filter(Site.Name == "Linville Falls" | Site.Name == "Durham Armory" | Site.Name ==
         "Leggett" | Site.Name == "Hattie Avenue" | Site.Name == "Clemmons Middle" |
         Site.Name == "Mendenhall School" | Site.Name == "Frying Pan Mountain" | Site.Name ==
         "West Johnston Co." | Site.Name == "Garinger High School" | Site.Name ==
         "Castle Hayne" | Site.Name == "Pitt Agri. Center" | Site.Name == "Bryson City" |
         Site.Name == "Millbrook School") %>%
  group_by(Date, Site.Name, AQS_PARAMETER_DESC, COUNTY) %>%
  summarise(mean_daily_AQI = mean(DAILY_AQI_VALUE), mean_LAT = mean(SITE_LATITUDE),
            mean_LONG = mean(SITE_LONGITUDE)) %>%
  mutate(Month = month(ymd(Date)), Year = year(ymd(Date)))
```

```
## 'summarise()' has grouped output by 'Date', 'Site.Name', 'AQS_PARAMETER_DESC'.
```

```
## You can override using the '.groups' argument.
```

```
# Resulted in 14,453 observations for 9 variables instead of 14,752

# 9 Spreading datasets using pivot_wider to create separate columns for ozone
# and PM 2.5 AQI values
library(tidyr)
EPAair_03_PM25_NC1819_Spread <- pivot_wider(EPAair_03_PM25_NC1819_Wrangled, names_from = AQS_PARAMETER_DESC,
      values_from = mean_daily_AQI)

# 10 Calling the dimensions of the new dataset: Returns 8,029 rows x 9 columns
# because AQI_daily_means has been divided into 'Ozone' or 'PM2.5' columns that
# were created from the 'AQS_PARAMETER_DESC' column
dim(EPAair_03_PM25_NC1819_Spread)
```

```
## [1] 8029    9
```

```
# 11 Saving the processed dataset
write.csv(EPAair_03_PM25_NC1819_Spread, row.names = FALSE, file = "Data/Processed/EPAair_PM25_NC2019_Processed.csv")
```

## Generate summary tables

12. Use the split-apply-combine strategy to generate a summary data frame. Data should be grouped by site, month, and year. Generate the mean AQI values for ozone and PM2.5 for each group. Then, add a pipe to remove instances where mean **ozone** values are not available (use the function `drop_na` in your pipe). It's ok to have missing mean PM2.5 values in this result.
13. Call up the dimensions of the summary dataset.

```
# 12 Creation of a summary data frame grouping by site, month, and year.
EPAair_03_PM25_NC1819_Summary <- EPAair_03_PM25_NC1819_Spread %>%
  group_by(Site.Name, Month, Year) %>%
  summarise(mean_O3_AQI = mean(Ozone), mean_PM25_AQI = mean(PM2.5)) %>%
  drop_na(mean_O3_AQI)
```

```
## 'summarise()' has grouped output by 'Site.Name', 'Month'. You can override
## using the '.groups' argument.
```

```
# 13 Calling the dimensions of the summary dataset.
dim(EPAair_03_PM25_NC1819_Summary) #Returns 205 rows x 5 columns
```

```
## [1] 205    5
```

14. Why did we use the function `drop_na` rather than `na.omit`?

Answer: We used the 'drop\_na' function as opposed to 'na.omit' because we can use this function to specify our desire to only drop those NA values for mean ozone concentrations, whereas 'na.omit' removes all incomplete cases for data within an entire data frame.