

- RAG Learning

2.4 Graphs

OCR GCSE Mathematics (J560) — Paper 1, 2 & 3 — Foundation and Higher

2.4.1 Coordinates and Midpoints

A point in 2D is given by its **coordinates** (x, y) , where x is the horizontal displacement from the origin and y is the vertical displacement.

The **midpoint** of the line segment joining (x_1, y_1) and (x_2, y_2) is:

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

The **distance** between two points (by Pythagoras):

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

Worked Example: Find the midpoint and length of the line segment joining $A(1, 3)$ and $B(7, 11)$.

Midpoint: $M = \left(\frac{1 + 7}{2}, \frac{3 + 11}{2} \right) = (4, 7)$

Length: $d = \sqrt{(7 - 1)^2 + (11 - 3)^2} = \sqrt{36 + 64} = \sqrt{100} = 10$


Exam Tip: The midpoint formula is an average of the two x -coordinates and an average of the two y -coordinates. You can think of it as "meet in the middle."

2.4.2 Straight Line Graphs

Any straight line can be written in the form $y = mx + c$, where:

- m = gradient (slope): how steep the line is = $\frac{\text{rise}}{\text{run}} = \frac{\Delta y}{\Delta x}$
- c = y -intercept: where the line crosses the y -axis.

Gradient and Equation of a Straight Line



Worked Example: Find the equation of the line through $(1, 3)$ and $(4, 9)$.

Gradient: $m = \frac{9 - 3}{4 - 1} = \frac{6}{3} = 2$

Using $y - y_1 = m(x - x_1)$ with point $(1, 3)$: $y - 3 = 2(x - 1)$

Expand: $y = 2x - 2 + 3 = 2x + 1$

Worked Example: A line has equation $3y - 6x = 12$. State the gradient and y -intercept.

Rearrange into $y = mx + c$ form: $3y = 6x + 12 \Rightarrow y = 2x + 4$

Gradient $m = 2$, y -intercept $= 4$ (i.e. the line crosses at $(0, 4)$).

Higher — Parallel and Perpendicular Lines

- **Parallel lines** have the same gradient: $y = 3x + 1$ and $y = 3x - 5$ are parallel (both have $m = 3$).
- **Perpendicular lines** have gradients that are negative reciprocals: if line 1 has gradient m , then any perpendicular line has gradient $-\frac{1}{m}$. Their product is -1 : $m_1 \times m_2 = -1$.




Worked Example: Find the equation of the line perpendicular to $y = 3x - 2$ that passes through $(6, 1)$.

Gradient of given line $= 3$. Perpendicular gradient $= -\frac{1}{3}$.

$$y - 1 = -\frac{1}{3}(x - 6)$$

$$y = -\frac{1}{3}x + 2 + 1 = -\frac{1}{3}x + 3$$

2.4.3 Quadratic, Cubic and Reciprocal Graphs



Key Features of Each Curve

Function	Shape	Key features
----------	-------	--------------

$y = x^2$	Parabola (U-shape)	Minimum at $(0, 0)$; symmetric about y -axis; always ≥ 0
$y = -x^2$	Parabola (n-shape)	Maximum at $(0, 0)$; always ≤ 0
$y = x^3$	S-shaped cubic	Passes through $(0, 0)$, $(-1, -1)$, $(1, 1)$; rotational symmetry about origin
$y = \frac{1}{x}$	Reciprocal (hyperbola)	Two branches; asymptotes along both axes ($x = 0$ and $y = 0$); undefined at $x = 0$

Plotting Quadratic Graphs: Table of Values

Worked Example: Plot $y = x^2 - 3x + 2$ for $-1 \leq x \leq 4$.

x	-1	0	1	2	3	4
x^2	1	0	1	4	9	16
$-3x$	3	0	-3	-6	-9	-12
$+2$	2	2	2	2	2	2
y	6	2	0	0	2	6

y -intercept: $(0, 2)$. x -intercepts (roots): $(1, 0)$ and $(2, 0)$.


Minimum between $x = 1$ and $x = 2$, at $x = 1.5$ (axis of symmetry).

Exam Tip: Always draw a smooth curve through plotted points — do not join with straight line segments. Plot at least 7 points for a quadratic.

2.4.4 Real-Life Graphs

Distance–Time Graphs

- **Gradient** of a distance–time graph = **speed** $\left(\text{speed} = \frac{\text{distance}}{\text{time}} \right)$.
- A horizontal line means the object is **stationary**.
- A steeper line means greater speed.
- A negative gradient means moving *back towards the start*.



Speed–Time Graphs

- **Gradient** of a speed–time graph = **acceleration** (in m/s^2).
- A horizontal line means **constant speed** (zero acceleration).
- **Area under the graph** = **distance travelled**.

Worked Example: A car accelerates from 0 to 20 m/s in 5 seconds, travels at 20 m/s for 10 seconds, then decelerates to 0 in 5 seconds. Find the total distance.

The graph is a trapezium with parallel sides 10 s and 20 s, height 20 m/s.

$$\text{Area} = \frac{1}{2}(10 + 20) \times 20 = \frac{1}{2} \times 30 \times 20 = 300 \text{ m}$$

Conversion Graphs

Straight-line graphs that convert between units (e.g. miles to km, £ to €). Read off values by locating one axis and reading across to the line, then down/up to the other axis.

Exam Tip: On distance–time graphs: steeper = faster. On speed–time graphs: steeper = greater acceleration. Area under speed–time graph = distance — and area under a velocity–time graph includes direction.

2.4.5 Transformations of Graphs [Higher]

Higher only

If $y = f(x)$ is the original graph, the transformed graphs are:

Transformation	Equation	Effect on graph
Translation up by a	$y = f(x) + a$	Shift the graph a units up
Translation down by a	$y = f(x) - a$	Shift the graph a units down
Translation left by a	$y = f(x + a)$	Shift the graph a units left (counter-intuitive!)
Translation right by a	$y = f(x - a)$	Shift the graph a units right
Reflection in x -axis	$y = -f(x)$	Reflect in the x -axis (negate all y values)
Reflection in y -axis	$y = f(-x)$	Reflect in the y -axis (negate all x values)
Stretch (vertical), scale factor a	$y = af(x)$	Multiply all y -coordinates by a
Stretch (horizontal), scale factor $\frac{1}{a}$	$y = f(ax)$	Multiply all x -coordinates by $\frac{1}{a}$ (compress if $a > 1$)

Memory aid for horizontal shifts: $y = f(x + a)$ shifts LEFT by a because the same y -value now occurs at a smaller x -value. Think "inside the bracket does the opposite."

Worked Example: The graph of $y = f(x)$ passes through $(2, 5)$. State the coordinates of the corresponding point on: (a) $y = f(x) + 3$; (b) $y = f(x - 4)$; (c) $y = -f(x)$.

(a) $y = f(x) + 3$: add 3 to y -coordinate: $(2, 8)$

(b) $y = f(x - 4)$: add 4 to x -coordinate (shift right): $(6, 5)$

(c) $y = -f(x)$: negate y -coordinate: $(2, -5)$

Exam Tip: Transformation questions often ask you to apply a transformation to a sketch or to state coordinates of key points after transformation. Always think about what happens to a single point before drawing.


2.4.6 Equation of a Circle [Higher]

Higher only

A circle with **centre at the origin** $(0, 0)$ and radius r has equation:

$$x^2 + y^2 = r^2$$

This follows directly from Pythagoras' theorem: for any point (x, y) on the circle, the distance to the centre is $\sqrt{x^2 + y^2} = r$.



Worked Example: (a) Write the equation of the circle with centre $(0, 0)$ and radius 7.
 (b) Does the point $(3, \sqrt{40})$ lie on, inside, or outside this circle?

(a) $x^2 + y^2 = 49$

(b) Compute $3^2 + (\sqrt{40})^2 = 9 + 40 = 49$. Since this equals $r^2 = 49$, the point lies **on** the circle.

Tangent to a Circle at a Point

A tangent to a circle at a point P is perpendicular to the radius OP at that point.

Worked Example: Find the equation of the tangent to the circle $x^2 + y^2 = 25$ at the point $(3, 4)$.

Gradient of radius OP : $m_{OP} = \frac{4 - 0}{3 - 0} = \frac{4}{3}$

Gradient of tangent (perpendicular): $m_t = -\frac{3}{4}$

Equation using point $(3, 4)$: $y - 4 = -\frac{3}{4}(x - 3)$

$y = -\frac{3}{4}x + \frac{9}{4} + 4 = -\frac{3}{4}x + \frac{25}{4}$

Or in integer form: $3x + 4y = 25$

Exam Tip: To verify a point is on a circle: substitute its coordinates into the equation and check. To find the tangent: find the gradient of the radius, then take the negative reciprocal.

Quick Reference Summary

Topic	Key formula / fact
Midpoint of (x_1, y_1) and (x_2, y_2)	$\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$
Gradient of line	$m = \frac{y_2 - y_1}{x_2 - x_1}$
Straight line equation	$y = mx + c$; or $y - y_1 = m(x - x_1)$
Parallel lines [H]	Same gradient m
Perpendicular lines [H]	Gradients satisfy $m_1 \times m_2 = -1$
Distance–time graph	Gradient = speed; horizontal = stationary
Speed–time graph	Gradient = acceleration; area under = distance
$y = f(x) + a$ [H]	Translate up by a
$y = f(x + a)$ [H]	Translate left by a
$y = -f(x)$ [H]	Reflect in x -axis
Circle (centre origin) [H]	$x^2 + y^2 = r^2$
Tangent to circle [H]	Perpendicular to radius at the point of contact

