

S5 — Redox, Electrochemistry and Transition Metals

5.1 Oxidation States and Redox Reactions

Oxidation States (Oxidation Numbers)

Oxidation state: A number assigned to an atom in a compound that represents the number of electrons it has lost, gained, or shared unequally compared to the uncombined element. It is a bookkeeping device for tracking electron transfer.

Rules for assigning oxidation states:

- Uncombined elements have an oxidation state of 0.
- In simple ions, the oxidation state equals the charge (e.g. $\text{Na}^+ = +1$, $\text{Cl}^- = -1$).
- Oxygen is -2 (except in peroxides where it is -1 , and in OF_2 where it is $+2$).
- Hydrogen is $+1$ (except in metal hydrides where it is -1).
- Fluorine is always -1 .
- The sum of oxidation states in a neutral compound is 0; in a polyatomic ion it equals the ionic charge.

Redox Definitions

Oxidation: Loss of electrons / increase in oxidation state.

Reduction: Gain of electrons / decrease in oxidation state.

Reducing agent: A substance that donates electrons (is itself oxidised).

Oxidising agent: A substance that accepts electrons (is itself reduced).

Remember: OIL RIG — Oxidation Is Loss, Reduction Is Gain (of electrons).

Half-Equations

A redox reaction can be split into two half-equations: one showing oxidation and one showing reduction. To write and balance half-equations:

1. Write the species being oxidised or reduced on each side.
2. Balance atoms other than O and H.
3. Balance oxygen by adding H_2O .
4. Balance hydrogen by adding H^+ .
5. Balance charge by adding electrons (e^-).

To combine half-equations, multiply each so the electrons cancel, then add the two equations together and simplify.

Exam Tip: When balancing redox equations, always check that atoms and charges balance on both sides. A common mistake is forgetting to balance hydrogen atoms after adding water molecules. Also check that the electrons cancel completely when combining half-equations.

5.2 Electrochemical Cells


Galvanic (Voltaic) Cells

Electrochemical cell: A device that converts chemical energy into electrical energy (galvanic cell) or electrical energy into chemical energy (electrolytic cell). A galvanic cell consists of two half-cells connected by a salt bridge and an external wire.

In a galvanic cell:

- The more reactive metal (more negative electrode potential) is the anode (negative terminal) where oxidation occurs.

- The less reactive metal (more positive electrode potential) is the cathode (positive terminal) where reduction occurs.
- Electrons flow through the external circuit from the anode to the cathode.
- The salt bridge completes the circuit by allowing ions to flow between the two half-cells, maintaining electrical neutrality.



Standard Electrode Potential

Standard electrode potential (E°): The EMF of a half-cell measured relative to the standard hydrogen electrode (SHE) under standard conditions (298 K, 100 kPa, 1.00 mol dm⁻³ solutions). The SHE is assigned a potential of 0.00 V by convention.

The standard hydrogen electrode consists of hydrogen gas at 100 kPa bubbled over a platinum electrode coated in platinum black, in contact with a 1.00 mol dm⁻³ solution of H⁺ ions at 298 K.

Standard Cell Potential

$$E^{\circ}_{\text{cell}} = E^{\circ}_{\text{cathode}} - E^{\circ}_{\text{anode}}$$

Alternatively: $E^{\circ}_{\text{cell}} = E^{\circ}_{\text{most positive half-cell}} - E^{\circ}_{\text{most negative half-cell}}$. The half-cell with the more positive E° value undergoes reduction (is the cathode); the other undergoes oxidation (is the anode).

Feasibility of Reactions

A reaction is thermodynamically feasible if $E^{\circ}_{\text{cell}} > 0$. This corresponds to a negative ΔG . The relationship is:

$$\Delta G^{\circ} = -nFE^{\circ}_{\text{cell}}$$

where n is the number of moles of electrons transferred and F is the Faraday constant ($96,485 \text{ C mol}^{-1}$).

Exam Tip: A positive E°_{cell} means the reaction is thermodynamically feasible, but it does not guarantee the reaction will occur at an observable rate. Kinetic factors (high activation energy) may prevent a feasible reaction from proceeding. Also, non-standard conditions can shift the equilibrium position and change the actual cell EMF.


Worked Example — Calculating E°_{cell} :

Given: $\text{Zn}^{2+}/\text{Zn } E^{\circ} = -0.76 \text{ V}$; $\text{Cu}^{2+}/\text{Cu } E^{\circ} = +0.34 \text{ V}$

$$E^{\circ}_{\text{cell}} = E^{\circ}(\text{cathode}) - E^{\circ}(\text{anode}) = E^{\circ}(\text{more positive}) - E^{\circ}(\text{more negative})$$

$$E^{\circ}_{\text{cell}} = +0.34 - (-0.76) = \mathbf{+1.10 \text{ V}}$$

Positive value means the reaction $\text{Zn} + \text{Cu}^{2+} \rightarrow \text{Zn}^{2+} + \text{Cu}$ is thermodynamically feasible. Zn is oxidised (anode), Cu^{2+} is reduced (cathode).



Electrolytic Cells

Electrolysis: The use of an external electrical energy source to drive a non-spontaneous chemical reaction. The electrolyte (an ionic compound, molten or in aqueous solution) is decomposed at the electrodes.

In electrolysis:

- The anode is positive — attracted by anions, where oxidation occurs.
- The cathode is negative — attracted by cations, where reduction occurs.
- In molten electrolytes, the ions discharged are straightforward (e.g. molten NaCl gives Na at the cathode and Cl_2 at the anode).
- In aqueous solutions, water may be preferentially discharged. At the cathode, H^+ is discharged if the metal cation has a very negative electrode potential. At the anode, OH^- is discharged in preference to most anions (except halide ions at moderate concentrations).

Fuel Cells

Fuel cell: An electrochemical cell that produces electricity continuously as long as fuel (typically hydrogen) and an oxidant (oxygen) are supplied. Unlike batteries, fuel cells do not need

recharging.

In a hydrogen fuel cell:

- Anode: $\text{H}_2(\text{g}) \rightarrow 2\text{H}^+(\text{aq}) + 2\text{e}^-$
- Cathode: $\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}(\text{l})$
- Overall: $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{O}(\text{l})$

Advantages: the only by-product is water (zero direct emissions), high efficiency compared to combustion engines. Disadvantages: hydrogen is difficult and expensive to store and transport, production of hydrogen often uses fossil fuels (unless from electrolysis using renewable energy), platinum catalysts are expensive.

5.3 Transition Metals

Definition and Electronic Configuration

Transition metal: A d-block element that forms at least one stable ion with a partially filled d sub-shell. Scandium (Sc^{3+} has an empty 3d sub-shell) and zinc (Zn^{2+} has a full 3d sub-shell) are d-block elements but are not transition metals by this definition.

The first row transition metals (Ti to Cu) have the general electron configuration $[\text{Ar}] 3\text{d}^n 4\text{s}^2$ (with exceptions for Cr and Cu). When forming ions, the 4s electrons are lost before the 3d electrons.

Variable Oxidation States


Transition metals can form ions with different oxidation states because the 3d and 4s energy levels are close together, so different numbers of electrons can be lost. Common examples:

Element	Common Oxidation States	Example Compounds/Ions
Titanium (Ti)	+3, +4	TiCl_3 , TiO_2
Vanadium (V)	+2, +3, +4, +5	V^{2+} (violet), V^{3+} (green), VO^{2+} (blue), VO_2^+ (yellow)

Manganese (Mn)	+2, +4, +7	Mn^{2+} , MnO_2 , MnO_4^-
Iron (Fe)	+2, +3	Fe^{2+} (green), Fe^{3+} (yellow/brown)
Copper (Cu)	+1, +2	Cu_2O , CuSO_4

Coloured Ions

Origin of colour: Transition metal ions in complexes are coloured because the d orbitals are split into two sets of different energies by the ligands. When visible light passes through the complex, photons of specific energy (wavelength) are absorbed to promote an electron from a lower d orbital to a higher one. The colour observed is the complementary colour of the light absorbed.



Factors that affect the colour of a transition metal complex:

- The identity of the metal ion and its oxidation state
- The type of ligand (different ligands cause different degrees of d-orbital splitting)
- The coordination number and geometry of the complex

Complex Ions

Complex ion: A central metal ion surrounded by ligands. The ligands are bonded to the metal ion by dative (coordinate) bonds, donating a lone pair of electrons into empty orbitals on the metal ion.

Ligand: A species with at least one lone pair of electrons that can form a dative bond to a central metal ion. Monodentate ligands donate one lone pair (e.g. H_2O , NH_3 , Cl^- , CN^-). Bidentate ligands donate two lone pairs from two different atoms (e.g. ethane-1,2-diamine, ethanedioate $\text{C}_2\text{O}_4^{2-}$). Polydentate ligands donate more (e.g. EDTA^{4-} is hexadentate).

Coordination number: The total number of dative bonds from ligands to the central metal ion.

Shapes of Complexes


Coordination Number	Shape	Bond Angle	Example
2	Linear	180°	$[\text{Ag}(\text{NH}_3)_2]^+$
4	Tetrahedral	109.5°	$[\text{CoCl}_4]^{2-}$
4	Square planar	90°	$[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ (cisplatin)
6	Octahedral	90°	$[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$, $[\text{Fe}(\text{CN})_6]^{3-}$

Six-coordinate complexes (coordination number 6) are typically octahedral. Four-coordinate complexes can be either tetrahedral or square planar depending on the metal and ligands. Tetrahedral is more common with larger ligands (e.g. Cl^-); square planar occurs especially with d^8 metals (e.g. Pt^{2+} , Ni^{2+} with strong-field ligands).

Shapes of Complex Ions

Octahedral


Coordination no. = 6



90° bond angles
[Cu(H₂O)₆]²⁺

Tetrahedral

Coordination no. = 4



109.5° bond angles
[CoCl₄]²⁻
Large ligands (Cl⁻)
favour tetrahedral

Square planar (CN=4, 90°): seen with d⁸ metals e.g. [Pt(NH₃)₂Cl₂] cisplatin

Ligand Substitution

Ligand substitution (ligand exchange) occurs when one ligand replaces another in a complex. This often causes a colour change because the new ligand splits the d orbitals differently.

Common examples with copper(II):

- [Cu(H₂O)₆]²⁺ (pale blue) + excess NH₃ → [Cu(NH₃)₄(H₂O)₂]²⁺ (deep blue) + 4H₂O
- [Cu(H₂O)₆]²⁺ (pale blue) + 4Cl⁻ → [CuCl₄]²⁻ (yellow/green) + 6H₂O

Note the change in coordination number from 6 (octahedral) to 4 (tetrahedral) when water is replaced by chloride ligands because Cl⁻ is a larger ligand.

Exam Tip: When writing equations for ligand substitution, always balance the equation carefully. The overall charge of the complex must be correct. Remember that replacing neutral ligands (H₂O, NH₃) with charged ligands (Cl⁻) changes the overall charge on the complex ion. When explaining colour, mention: (1) d-orbital splitting by ligands, (2) d-d transitions absorbing specific visible wavelengths, (3) complementary colour transmitted. If d⁰ or d¹⁰, no d-d transition is possible and the compound is white/colourless.

Catalytic Activity

Transition metals and their compounds are effective catalysts because:

- They have variable oxidation states, allowing them to accept and donate electrons during catalytic cycles.
- They can form complexes with reactant molecules, providing a surface (heterogeneous) or intermediate complex (homogeneous) that lowers the activation energy.

Examples: Fe in the Haber process ($\text{N}_2 + 3\text{H}_2 \rightarrow 2\text{NH}_3$), V_2O_5 in the Contact process ($2\text{SO}_2 + \text{O}_2 \rightarrow 2\text{SO}_3$), MnO_2 in the decomposition of H_2O_2 , Ni in hydrogenation of alkenes.

Precipitation Reactions

Adding sodium hydroxide or ammonia solution to aqueous transition metal ions produces coloured precipitates of metal hydroxides:

Ion	NaOH(aq) precipitate	Excess NaOH	NH ₃ (aq) precipitate	Excess NH ₃
Cu^{2+}	Blue, $\text{Cu}(\text{OH})_2$	No change	Blue, $\text{Cu}(\text{OH})_2$	Dissolves: deep blue $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$
Fe^{2+}	Green, $\text{Fe}(\text{OH})_2$	No change	Green, $\text{Fe}(\text{OH})_2$	No change
Fe^{3+}	Brown, $\text{Fe}(\text{OH})_3$	No change	Brown, $\text{Fe}(\text{OH})_3$	No change
Cr^{3+}	Green, $\text{Cr}(\text{OH})_3$	Dissolves: dark green $[\text{Cr}(\text{OH})_6]^{3-}$	Green, $\text{Cr}(\text{OH})_3$	Dissolves: purple $[\text{Cr}(\text{NH}_3)_6]^{3+}$

Colorimetry


Colorimetry: An analytical technique that measures the concentration of a coloured solution by measuring how much light of a specific wavelength it absorbs. The amount of light absorbed is proportional to the concentration of the coloured species (Beer-Lambert law).

$$A = \epsilon cl$$

where A is absorbance (no units), ϵ is the molar absorption coefficient ($\text{dm}^3 \text{mol}^{-1} \text{cm}^{-1}$), c is concentration (mol dm^{-3}), and l is the path length (cm).

A calibration curve is prepared by measuring the absorbance of solutions of known concentration. The absorbance of the unknown solution is then read off the calibration curve to determine its concentration. The wavelength of light used should be the one at which the solution absorbs most strongly (the complementary colour).

Redox Titration — KMnO_4 / Fe^{2+}



Burette

KMnO_4
(purple)

$\text{Fe}^{2+}(\text{aq})$
+ H_2SO_4

Conical flask

Half-equations:

$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$ (reduction)

$\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^-$ (oxidation, $\times 5$)

Overall: $\text{MnO}_4^- + 8\text{H}^+ + 5\text{Fe}^{2+} \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O} + 5\text{Fe}^{3+}$

End Point (self-indicating)

Endpoint: first permanent pink/purple colour

Before endpoint: MnO_4^- decolourised by excess Fe^{2+}

Before endpoint

$\text{MnO}_4^- \rightarrow \text{Mn}^{2+}$
purple $\rightarrow$ colourless

At endpoint

Excess MnO_4^-
stays pink/purple