

CP1.1: Complex Numbers

1. Introduction and Definition

Imaginary unit: The imaginary unit i is defined by $i^2 = -1$, so $i = \sqrt{-1}$.

Complex number: A complex number has the form $z = a + bi$ where $a, b \in \mathbb{R}$. The set of all complex numbers is denoted $\mathbb{C}$.

Real part: $\text{Re}(z) = a$

Imaginary part: $\text{Im}(z) = b$ (note: this is the real number b , not bi)

Every real number is also a complex number with imaginary part zero. Numbers of the form bi (where $b \neq 0$) are called *purely imaginary*.

Powers of i

The powers of i cycle with period 4:

Power	i^1	i^2	i^3	i^4	i^5
Value	i	-1	$-i$	1	i

To evaluate i^n : find the remainder when n is divided by 4 and use the table above.

Exam tip: When you see $\sqrt{-k}$ for $k > 0$, write it as $i\sqrt{k}$. Never leave a negative under a square root in your final answer. For example: $\sqrt{-9} = 3i$, not $3\sqrt{-1}$.

2. Arithmetic with Complex Numbers

Addition and Subtraction

Add or subtract real and imaginary parts separately:

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i$$

Multiplication

Expand using FOIL (like with surds), then substitute $i^2 = -1$:

$$(a + bi)(c + di) = ac + adi + bci + bdi^2 = (ac - bd) + (ad + bc)i$$

Example: Find $(3 + 2i)(1 - 4i)$.

$$\begin{aligned}(3 + 2i)(1 - 4i) &= 3(1) + 3(-4i) + 2i(1) + 2i(-4i) \\&= 3 - 12i + 2i - 8i^2 \\&= 3 - 10i - 8(-1) \\&= 3 - 10i + 8 = 11 - 10i\end{aligned}$$

The Complex Conjugate

Complex conjugate: If $z = a + bi$, then the conjugate is $z^* = a - bi$ (also written $\bar{z}$).

Key properties:

- $z + z^* = 2a$ (always real)
- $z \cdot z^* = a^2 + b^2$ (always real and non-negative)
- $(z^*)^* = z$
- $(z_1 z_2)^* = z_1^* z_2^*$
- $(z_1 + z_2)^* = z_1^* + z_2^*$

Division

To divide complex numbers, multiply both numerator and denominator by the conjugate of the denominator:

$$\frac{a + bi}{c + di} = \frac{(a + bi)(c - di)}{(c + di)(c - di)} = \frac{(ac + bd) + (bc - ad)i}{c^2 + d^2}$$

The denominator $c^2 + d^2$ is always real, giving a result in the form $x + yi$.

Example: Simplify $\frac{3 + 5i}{1 + 2i}$.

Multiply top and bottom by conjugate $(1 - 2i)$:

$$\frac{(3 + 5i)(1 - 2i)}{(1 + 2i)(1 - 2i)} = \frac{3 - 6i + 5i - 10i^2}{1 - 4i^2} = \frac{3 - i + 10}{1 + 4} = \frac{13 - i}{5} = \frac{13}{5} - \frac{1}{5}i$$

Exam tip: Always check your division: multiply your answer by the original denominator and verify you get the numerator back. Losing marks on arithmetic in division is very common.

3. Modulus and Argument


Modulus: $|z| = \sqrt{a^2 + b^2}$ — this is the distance of z from the origin in the Argand diagram.

Argument: $\arg(z) = \theta$ where $\tan \theta = \frac{b}{a}$ — this is the angle measured anticlockwise from the positive real axis.

The **principal argument** satisfies $-\pi < \arg(z) \leq \pi$.

Argand Diagram

A complex number $z = a + bi$ is represented as a point (a, b) in the complex plane, where the horizontal axis is the real axis and the vertical axis is the imaginary axis.



Calculating the Argument Carefully

Use $\tan \theta = b/a$ but you must place the angle in the correct quadrant. Always draw a sketch first:

Quadrant	Sign of a, b	Principal argument range	Method
1st	$a > 0, b > 0$	$0 < \theta \leq \frac{\pi}{2}$	$\theta = \arctan\left(\frac{b}{a}\right)$
2nd	$a < 0, b > 0$	$\frac{\pi}{2} < \theta \leq \pi$	$\theta = \pi - \arctan\left(\frac{b}{ a }\right)$
3rd	$a < 0, b < 0$	$-\pi < \theta \leq -\frac{\pi}{2}$	$\theta = -\pi + \arctan\left(\frac{ b }{ a }\right)$
4th	$a > 0, b < 0$	$-\frac{\pi}{2} < \theta < 0$	$\theta = -\arctan\left(\frac{ b }{a}\right)$

Example: Find $|z|$ and $\arg(z)$ for $z = -1 + \sqrt{3}i$.

$$|z| = \sqrt{(-1)^2 + (\sqrt{3})^2} = \sqrt{1 + 3} = 2$$

z is in the second quadrant ($a < 0, b > 0$).

$$\text{Reference angle: } \arctan\left(\frac{\sqrt{3}}{1}\right) = \frac{\pi}{3}$$

$$\text{Therefore } \arg(z) = \pi - \frac{\pi}{3} = \frac{2\pi}{3}$$

Exam tip: The most common error is computing $\arctan(b/a)$ and forgetting to adjust for quadrant. Always establish the quadrant FIRST, then compute the reference angle, then adjust.

Modulus-Argument Properties

$$|z_1 z_2| = |z_1| |z_2| \quad \arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$$

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|} \quad \arg\left(\frac{z_1}{z_2}\right) = \arg(z_1) - \arg(z_2)$$

(Arguments are taken modulo 2π to keep them in the principal range $(-\pi, \pi]$.)

4. Solving Quadratic Equations with Complex Roots

When a quadratic $az^2 + bz + c = 0$ has a negative discriminant ($b^2 - 4ac < 0$), the roots are complex conjugates of each other.

$$z = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Worked Example: Solve $z^2 + 4z + 13 = 0$.

Using the quadratic formula with $a = 1, b = 4, c = 13$:

$$z = \frac{-4 \pm \sqrt{16 - 52}}{2} = \frac{-4 \pm \sqrt{-36}}{2} = \frac{-4 \pm 6i}{2} = -2 \pm 3i$$

So the roots are $z = -2 + 3i$ and $z = -2 - 3i$.

Verification: Sum of roots $= (-2 + 3i) + (-2 - 3i) = -4 = -b/a \checkmark$

Product of roots $= (-2 + 3i)(-2 - 3i) = 4 + 9 = 13 = c/a \checkmark$

Conjugate Root Theorem: If the coefficients of a polynomial are all real, then complex roots occur in conjugate pairs. That is, if $z = p + qi$ is a root, then $z^* = p - qi$ is also a root.

5. Solving Cubic and Quartic Equations

Cubics with Complex Roots

A cubic with real coefficients has either:

- Three real roots, or
- One real root and one conjugate pair of complex roots

Strategy: If given one complex root α , immediately write down α^* as another root. The factor $(z - \alpha)(z - \alpha^*)$ is a real quadratic. Divide the cubic by this quadratic (polynomial long division or compare coefficients) to find the third root.

Example: Given that $2 + i$ is a root of $z^3 - 7z^2 + 17z - 15 = 0$, find all roots.

Since coefficients are real, $2 - i$ is also a root.

Form the quadratic factor:

$$(z - (2 + i))(z - (2 - i)) = (z - 2)^2 + 1 = z^2 - 4z + 5$$

Divide: $z^3 - 7z^2 + 17z - 15 \div (z^2 - 4z + 5)$

$$z^3 - 7z^2 + 17z - 15 = (z^2 - 4z + 5)(z - 3)$$

So the three roots are $z = 2 + i$, $z = 2 - i$, $z = 3$.

Quartics with Complex Roots

A quartic with real coefficients has either:

- Four real roots
- Two real roots and one conjugate pair
- Two conjugate pairs of complex roots

Strategy: If given one complex root α , use α^* to form one quadratic factor. Divide the quartic by it to get another quadratic, then solve that for the remaining roots.

Example: Given that $1 + 2i$ is a root of $z^4 - 2z^3 + 6z^2 - 2z + 5 = 0$, find all roots.

Conjugate pair: $1 + 2i$ and $1 - 2i$.

Quadratic factor: $(z - 1 - 2i)(z - 1 + 2i) = (z - 1)^2 + 4 = z^2 - 2z + 5$

Divide: $z^4 - 2z^3 + 6z^2 - 2z + 5 = (z^2 - 2z + 5)(z^2 + 1)$

Solve $z^2 + 1 = 0 \Rightarrow z^2 = -1 \Rightarrow z = \pm i$

All four roots: $1 + 2i$, $1 - 2i$, i , $-i$

Finding a Root by Substitution

If asked to verify or find a complex root directly, substitute into the polynomial and set real and imaginary parts equal to zero. This gives a system of equations.

Example: Find a and b if $z = 1 + 2i$ is a root of $z^2 + az + b = 0$ (where $a, b \in \mathbb{R}$).

Sum of roots: $(1 + 2i) + (1 - 2i) = 2 = -a \Rightarrow a = -2$

Product of roots: $(1 + 2i)(1 - 2i) = 5 = b \Rightarrow b = 5$

6. Equating Real and Imaginary Parts

Two complex numbers are equal if and only if their real parts are equal AND their imaginary parts are equal:

$$a + bi = c + di \iff a = c \text{ and } b = d$$

Example: Solve $z^2 = 5 + 12i$.

Let $z = x + yi$. Then $(x + yi)^2 = x^2 - y^2 + 2xyi = 5 + 12i$.

Equating real parts: $x^2 - y^2 = 5 \dots (1)$

Equating imaginary parts: $2xy = 12$, so $y = \frac{6}{x} \dots (2)$

Substituting (2) into (1): $x^2 - \frac{36}{x^2} = 5$

$$x^4 - 5x^2 - 36 = 0 \Rightarrow (x^2 - 9)(x^2 + 4) = 0$$

Since $x \in \mathbb{R}$: $x^2 = 9 \Rightarrow x = \pm 3$

When $x = 3$: $y = 2 \rightarrow z = 3 + 2i$

When $x = -3$: $y = -2 \rightarrow z = -3 - 2i$

Exam tip: When finding square roots of complex numbers by letting $z = x + yi$ and equating parts, remember that you always get two answers (one the negative of the other). Both must be stated.

7. Key Results Summary

Property	Formula
Definition of i	$i^2 = -1$
Conjugate	$z^* = a - bi$ if $z = a + bi$
Modulus	$ z = \sqrt{a^2 + b^2}$
Product $\times$ Conjugate	$zz^* = z ^2 = a^2 + b^2$

Argument (principal)	$-\pi < \arg(z) \leq \pi$
Modulus of product	$ z_1 z_2 = z_1 z_2 $
Argument of product	$\arg(z_1 z_2) = \arg z_1 + \arg z_2$
Conjugate root theorem	Real-coefficient polynomials: complex roots in conjugate pairs

8. Practice Checklist

- Can you add, subtract, multiply, and divide complex numbers in $a + bi$ form?
- Can you find the modulus and principal argument of any complex number?
- Can you solve quadratics with negative discriminant?
- Given one complex root of a cubic/quartic, can you find all other roots?
- Can you find square roots of complex numbers using the "let $z = x + yi$ " method?
- Can you use the conjugate root theorem to form factor pairs?