

1.5 Trigonometry

Mathematics — OCR A-Level (H240) — Paper 1: Pure Mathematics

Trigonometry began as the mathematics of triangles — ratios of sides used in surveying, navigation and astronomy — but at A-level it becomes a far richer subject about *periodic functions*. The same three ratios $\sin$, $\cos$ and $\tan$ that describe a right-angled triangle also describe circular motion, oscillations, alternating current, tides and sound waves. This chapter develops the machinery needed to work with them as functions: radian measure, graphs and transformations, the reciprocal functions $\sec$, $\csc$, $\cot$, a family of *identities* (Pythagorean, compound-angle, double-angle, the $R \sin(\theta + \alpha)$ form), methods for solving trig equations, and the inverse functions $\arcsin$, $\arccos$, $\arctan$. Throughout, the key idea is that every trig identity is a statement true for *all* valid angles, while every trig equation asks for the specific angles making a statement true.

1. Degrees and Radians

An angle can be measured in two principal units. Degrees split a full turn into 360 equal parts — convenient for everyday use but mathematically arbitrary. Radians, by contrast, measure an angle by the arc length it subtends on a unit circle, and are the natural unit for calculus: the derivative of $\sin x$ is $\cos x$ only when x is in radians.

Radian. One radian is the angle subtended at the centre of a circle by an arc equal in length to the radius. Consequently a full turn (circumference $2\pi r$) subtends 2π radians, so $[180^\circ = \pi \text{ rad}], \quad 1 \text{ rad} = \frac{180^\circ}{\pi} \approx 57.2958^\circ.$

Converting Between Units

To convert from degrees to radians multiply by $\pi/180$; to convert from radians to degrees multiply by $180/\pi$.

Degrees	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Radians	0	$\pi/6$	$\pi/4$	$\pi/3$	$\pi/2$	$2\pi/3$	$3\pi/4$	$5\pi/6$	π	$3\pi/2$	2π

Exam tip — set your calculator to the correct mode. If the question uses π , the angle is in radians; if it uses the $^\circ$ symbol, the angle is in degrees. A calculator in the wrong mode is the single most common cause of marks lost in trigonometry questions. Always check the small DEG or RAD indicator before evaluating $\sin 30$.

Arc Length and Sector Area

When θ is measured in *radians*, two elegant formulae follow directly from the definition:

Arc length. An arc of a circle of radius r , subtending an angle θ radians at the centre, has length $s = r\theta$.

Sector area. The area of a circular sector of radius r and angle θ radians is $A = \frac{1}{2}r^2\theta$. The area of the corresponding triangular *segment* (bounded by the chord and two radii) is $\frac{1}{2}r^2 \sin \theta$, so the crescent-shaped minor segment has area $\frac{1}{2}r^2(\theta - \sin \theta)$.

Both formulae fail in degrees: if θ is in degrees you must first convert or use $s = \pi r\theta/180$ and $A = \pi r^2\theta/360$.

Worked Example 1 — Arc, sector, segment

A sector of a circle has radius $r = 10$ cm and subtends $\theta = 1.2$ rad at the centre. Find the arc length, sector area, chord length and minor segment area.

1. Arc length: $s = r\theta = 10 \cdot 1.2 = 12$ cm.
2. Sector area: $A = \frac{1}{2}r^2\theta = \frac{1}{2} \cdot 100 \cdot 1.2 = 60$ cm².
3. Chord length (isosceles triangle, two sides 10, included angle 1.2 rad):
4. $c^2 = 10^2 + 10^2 - 2 \cdot 10 \cdot 10 \cdot \cos 1.2$ (cosine rule)
5. $= 200 - 200 \cdot 0.36236$
6. $= 200 - 72.47 = 127.53$

7. $c \approx 11.29 \text{ cm}.$
 8. Triangle area: $\frac{1}{2} \cdot 10 \cdot 10 \cdot \sin 1.2 = 50 \cdot 0.9320 \approx 46.60 \text{ cm}^2.$
 9. Segment area: $A - \text{triangle} = 60 - 46.60 = 13.40 \text{ cm}^2.$
- // The segment is the thin crescent between chord and arc.

Small-Angle Approximations

When an angle x is small and measured in *radians*, the trig functions simplify dramatically. These approximations are the starting point of calculus with trig functions.

Small-angle approximations. For x small and in radians, $\sin x \approx x$, $\cos x \approx 1 - \frac{x^2}{2}$, $\tan x \approx x$. The errors are of order x^3 , x^4 and x^3 respectively, so for $|x| < 0.05$ all three give four-decimal accuracy.

These come from the Taylor (power) series of the trig functions. Intuitively, for a small arc on the unit circle, the arc length x and the vertical drop $\sin x$ are nearly equal, and the horizontal coordinate $\cos x$ drops from 1 by approximately half x^2 .

Exam tip — the approximations only work in radians. $\sin x \approx x$ is false in degrees: $\sin 5^\circ = 0.0872$, not 5. If a question uses small-angle approximations, the angles must be converted to radians first. Writing $\sin 5 \approx 5$ in a script will lose every mark in the part.

Worked Example 2 — Small-angle limit

Use small-angle approximations to estimate $\lim_{x \rightarrow 0} \frac{\sin 3x}{x}$ and $\lim_{x \rightarrow 0} \frac{1 - \cos 2x}{x^2}$.

1. For $x \rightarrow 0$, use $\sin(3x) \approx 3x$:
2. $\sin(3x)/x \approx 3x/x = 3.$
3. For $\cos(2x)$ near 0 use $\cos(2x) \approx 1 - (2x)^2/2 = 1 - 2x^2$:
4. $1 - \cos(2x) \approx 1 - (1 - 2x^2) = 2x^2.$
5. $(1 - \cos 2x)/x^2 \approx 2x^2/x^2 = 2.$

```
// Both limits are rigorous – the approximation error vanishes as  $x \rightarrow 0$ .
```

2. Graphs of sin, cos and tan

Defining $\sin \theta$ and $\cos \theta$ as the y - and x -coordinates of the point on the unit circle at angle θ from the positive x -axis immediately extends them to *all* real angles, not just those in a right-angled triangle. This gives graphs that repeat: they are *periodic functions*.

$y = \sin x$

- **Domain:** all real x .
- **Range:** $-1 \leq \sin x \leq 1$.
- **Period:** 2π (or 360°).
- **Amplitude:** 1.
- **Roots:** $x = k\pi, k \in \mathbb{Z}$.
- **Maxima:** at $x = \pi/2 + 2k\pi$, value 1. **Minima:** at $x = -\pi/2 + 2k\pi$, value -1 .
- **Symmetry:** odd: $\sin(-x) = -\sin x$.

$y = \cos x$

- **Domain:** all real x . **Range:** $[-1, 1]$. **Period:** 2π .
- **Roots:** $x = \pi/2 + k\pi$.
- **Maxima:** $x = 2k\pi$. **Minima:** $x = \pi + 2k\pi$.
- **Symmetry:** even: $\cos(-x) = \cos x$. Also $\cos x = \sin(x + \pi/2)$ — the cosine curve is the sine curve shifted left by $\pi/2$.

$y = \tan x$

- **Domain:** all real x except $x = \pi/2 + k\pi$ (where $\cos x = 0$).
- **Range:** all real numbers.
- **Period:** π — shorter than sin and cos.
- **Vertical asymptotes:** at $x = \pi/2 + k\pi$.

- **Roots:** $x = k\pi$.
- **Symmetry:** odd: $\tan(-x) = -\tan x$.

Function	Period	Amplitude	Domain	Range
$\sin x$	2π	1	$\mathbb{R}$	$[-1, 1]$
$\cos x$	2π	1	$\mathbb{R}$	$[-1, 1]$
$\tan x$	π	undefined	$x \neq \pi/2 + k\pi$	$\mathbb{R}$

Transformations $y = a \sin(bx + c) + d$

Each of the four parameters reshapes the standard sine wave:

- a — **amplitude scale**. The range becomes $[d - |a|, d + |a|]$. If $a < 0$, the graph is reflected in the horizontal line $y = d$.
- b — **horizontal (frequency) scale**. Period becomes $2\pi/|b|$. Larger b squashes the graph horizontally.
- c — **phase shift**. The graph shifts left by c/b : $\sin(bx + c) = \sin(b(x + c/b))$.
- d — **vertical shift**. Moves the midline to $y = d$.

Exam tip — identifying a sinusoidal graph. Given a wave graph, read amplitude a from half the vertical distance between max and min; the vertical shift d is the midline $y = \frac{1}{2}(\max + \min)$; the period gives b via $b = 2\pi/T$; and the phase c is fixed by one known point such as a maximum. Work in that order for reliability.

Worked Example 3 — Reading a transformation

The function $y = 3 \sin(2x - \frac{\pi}{3}) + 1$ has what amplitude, period, phase shift, and range?

1. Compare with $a \sin(bx + c) + d$.
2. $a = 3$, $b = 2$, $c = -\pi/3$, $d = 1$.
3. Amplitude = $|a| = 3$.
4. Period = $2\pi/|b| = 2\pi/2 = \pi$.

5. Phase shift: $2x - \pi/3 = 2(x - \pi/6) \Rightarrow$ shift right by $\pi/6$.
6. Vertical shift: $d = 1$ (midline $y = 1$).
7. Range: $[1 - 3, 1 + 3] = [-2, 4]$.

3. Reciprocal Trig Functions

Three further trig functions are defined as reciprocals of the three main ones:

$$\sec x = \frac{1}{\cos x}, \quad \csc x = \frac{1}{\sin x}, \quad \cot x = \frac{1}{\tan x} = \frac{\cos x}{\sin x}.$$

(UK textbooks sometimes write $\operatorname{cosec} x$ for $\csc x$.) Each has vertical asymptotes wherever its defining function is zero, and a range that excludes the interval $(-1, 1)$ for $\sec$ and $\csc$.

Function	Definition	Period	Domain (excluded points)	Range
$\sec x$	$1/\cos x$	2π	$x \neq \pi/2 + k\pi$	$(-\infty, -1] \cup [1, \infty)$
$\csc x$	$1/\sin x$	2π	$x \neq k\pi$	$(-\infty, -1] \cup [1, \infty)$
$\cot x$	$\cos x/\sin x$	π	$x \neq k\pi$	$\mathbb{R}$

Between consecutive asymptotes, the graphs of $\sec$ and $\csc$ look like opposing cup and cap shapes — the portions of the cosine or sine curve between ± 1 and $\pm \infty$. The graph of $\cot$ decreases from $+\infty$ to $-\infty$ between each pair of zeros of $\sin$, mirror-image of $\tan$.

4. Exact Values

Certain special angles have trig values expressible as simple surds. These must be memorised: OCR expects them without a calculator. They follow from two simple geometric facts — the 45° - 45° - 90° isosceles right triangle (sides $1, 1, \sqrt{2}$) and the 30° - 60° - 90° half-equilateral triangle (sides $1, \sqrt{3}, 2$).

Angle (deg)	Angle (rad)	$\sin$	$\cos$	$\tan$
0°	0	0	1	0

30°	$\pi/6$	$1/2$	$\sqrt{3}/2$	$1/\sqrt{3} = \sqrt{3}/3$
45°	$\pi/4$	$\sqrt{2}/2$	$\sqrt{2}/2$	1
60°	$\pi/3$	$\sqrt{3}/2$	$1/2$	$\sqrt{3}$
90°	$\pi/2$	1	0	undefined

Exam tip — memory device. Write $\sin 0, \sin 30, \sin 45, \sin 60, \sin 90$ as $\frac{\sqrt{0}}{2}, \frac{\sqrt{1}}{2}, \frac{\sqrt{2}}{2}, \frac{\sqrt{3}}{2}, \frac{\sqrt{4}}{2}$ — a neat pattern. Cosine is the reverse order. $\tan$ is $\sin / \cos$. From these you can generate the values in all four quadrants using the CAST diagram.

5. Pythagorean Identity and its Corollaries

On the unit circle the point at angle θ has coordinates $(\cos \theta, \sin \theta)$. Since the radius is 1, Pythagoras's theorem gives the foundational identity

$$\sin^2 \theta + \cos^2 \theta = 1 \quad \text{for all } \theta.$$

Dividing through by $\cos^2 \theta$ (valid when $\cos \theta \neq 0$) gives the $\tan$ -sec form; dividing by $\sin^2 \theta$ gives the $\cot$ -csc form:

$$\tan^2 \theta + 1 = \sec^2 \theta, \quad 1 + \cot^2 \theta = \csc^2 \theta.$$

Identity	Rearrangement useful for
$\sin^2 \theta + \cos^2 \theta = 1$	Swapping between $\sin^2$ and $\cos^2$; simplifying under a surd.
$\sec^2 \theta = 1 + \tan^2 \theta$	Integrals like $\int \sec^2 x \, dx$; eliminating $\sec$ in equations.
$\csc^2 \theta = 1 + \cot^2 \theta$	Integrals like $\int \csc^2 x \, dx$; eliminating $\csc$.

Worked Example 4 — Using the Pythagorean identity to solve a quadratic in $\sin$

Solve $2 \cos^2 x + \sin x = 1$ for $0 \leq x \leq 2\pi$.

1. Replace $\cos^2 x$ using $\cos^2 x = 1 - \sin^2 x$:
2. $2(1 - \sin^2 x) + \sin x = 1$
3. $2 - 2\sin^2 x + \sin x = 1$
4. $2\sin^2 x - \sin x - 1 = 0$.
5. Let $s = \sin x$: $2s^2 - s - 1 = 0 \Rightarrow (2s + 1)(s - 1) = 0$.
6. $s = 1$ or $s = -1/2$.
7. Case $\sin x = 1$: $x = \pi/2$.
8. Case $\sin x = -1/2$: principal value $\arcsin(-1/2) = -\pi/6$,
9. in $[0, 2\pi]$: $x = \pi + \pi/6 = 7\pi/6$ or $x = 2\pi - \pi/6 = 11\pi/6$.
10. $\therefore x \in \{\pi/2, 7\pi/6, 11\pi/6\}$.

6. Compound-Angle Formulae

The compound-angle formulae express $\sin(A \pm B)$, $\cos(A \pm B)$ and $\tan(A \pm B)$ in terms of the trig values of A and B separately. They are the central toolkit of A-level trigonometry: every other identity on the syllabus (double-angle, $R \sin(\theta + \alpha)$) is a corollary.

Compound-angle formulae. $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$, $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$, $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$.

The top sign in each formula goes with $+$, the bottom with $-$. Note the sign *flip* in the cosine formula: $\cos(A + B) = \cos A \cos B - \sin A \sin B$.

Geometric Proof of $\sin(A + B)$

Consider a diagram with two right-angled triangles stacked. In a unit square, place a point P so that $\angle POQ = A$ along the horizontal, then rotate through a further angle B to reach R . Drop perpendiculars and compare lengths. The vertical projection of OR onto the y -axis decomposes as

$$\sin(A + B) = \underbrace{\sin A \cos B}_{\text{rise from rotating } A \text{ along horizontal}} + \underbrace{\cos A \sin B}_{\text{rise from rotating } B \text{ perpendicular}}.$$

A fully rigorous proof uses the rotation matrix $R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ and the identity $R(A + B) = R(A)R(B)$. Multiplying the two matrices and comparing entries gives both the sine and cosine compound-angle formulae simultaneously.

Exam tip — memory pattern for compound angles. $\sin(A + B)$ expands as "sincos + cossin" (the letters always appear in the order sin-cos, cos-sin). $\cos(A + B)$ expands as "coscos – sinsin". The signs in cos are opposite to the input sign; the signs in sin match. Writing a formula sheet from memory each week until exam is the most efficient way to lock these in.

Worked Example 5 — Exact value using compound angle

Find the exact value of $\sin 75^\circ$.

1. Write $75^\circ = 45^\circ + 30^\circ$.
2. $\sin(45^\circ + 30^\circ) = \sin 45^\circ \cos 30^\circ + \cos 45^\circ \sin 30^\circ$
3. $= (\sqrt{2}/2)(\sqrt{3}/2) + (\sqrt{2}/2)(1/2)$
4. $= \sqrt{6}/4 + \sqrt{2}/4$
5. $= (\sqrt{6} + \sqrt{2})/4$.
6. Numerically: $(2.449 + 1.414)/4 \approx 0.9659$ ✓ (matches calculator).

// Same idea: $\cos 75^\circ = (\sqrt{6} - \sqrt{2})/4$, $\tan 75^\circ = 2 + \sqrt{3}$.

Worked Example 6 — Proving an identity

Prove $\sin(x + \pi/2) = \cos x$ using the compound-angle formula.

1. $\sin(x + \pi/2) = \sin x \cos(\pi/2) + \cos x \sin(\pi/2)$
2. $= \sin x \cdot 0 + \cos x \cdot 1$
3. $= \cos x$. ✓

7. Double-Angle Formulae

Setting $B = A$ in the compound-angle formulae gives the *double-angle* formulae, which express trig functions of $2A$ in terms of trig functions of A .

Double-angle formulae. $\sin 2A = 2 \sin A \cos A$, $\cos 2A = \cos^2 A - \sin^2 A = 1 - 2\sin^2 A = 2\cos^2 A - 1$, $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$. The three equivalent forms of $\cos 2A$ follow from $\sin^2 A + \cos^2 A = 1$.

Derivation from Compound Angles

Take $B = A$ in $\sin(A + B) = \sin A \cos B + \cos A \sin B$: this gives $\sin 2A = \sin A \cos A + \cos A \sin A = 2 \sin A \cos A$. Similarly $\cos(A + A) = \cos^2 A - \sin^2 A$, and the alternative forms follow by substituting $\sin^2 A = 1 - \cos^2 A$ or $\cos^2 A = 1 - \sin^2 A$.

Half-Angle / Power-Reducing Identities

Rearranging the $\cos 2A$ identities gives the forms needed for integration:

$$\sin^2 A = \frac{1 - \cos 2A}{2}, \quad \cos^2 A = \frac{1 + \cos 2A}{2}.$$

These let you integrate $\sin^2$ and $\cos^2$ by linearising them — an essential calculus technique.

Exam tip — choosing the right form of $\cos 2A$. If the equation already contains only $\sin$, use $\cos 2A = 1 - 2\sin^2 A$. If it contains only $\cos$, use $\cos 2A = 2\cos^2 A - 1$. If it contains both, you may as well stick with $\cos^2 A - \sin^2 A$. Picking the right form saves a line of algebra and reduces slip risk.

Worked Example 7 — Solving with double angle

Solve $\sin 2x = \cos x$ for $0 \leq x \leq 2\pi$.

1. Use $\sin 2x = 2 \sin x \cos x$:
2. $2 \sin x \cos x = \cos x$
3. $2 \sin x \cos x - \cos x = 0$
4. $\cos x (2 \sin x - 1) = 0$.
5. $\cos x = 0 \Rightarrow x = \pi/2, 3\pi/2$.

$$6. \quad \sin x = 1/2 \Rightarrow x = \pi/6, 5\pi/6.$$

$$7. \quad \therefore x \in \{\pi/6, \pi/2, 5\pi/6, 3\pi/2\}.$$

// WARNING: dividing both sides by $\cos x$ loses the solutions $\cos x = 0$.

// Always factor; never divide by something that could be zero.

8. The $R \sin(\theta + \alpha)$ Form

Any linear combination $a \sin \theta + b \cos \theta$ can be written as a single sine wave $R \sin(\theta + \alpha)$ (or as a single cosine, depending on taste). This is essential for finding maxima and minima of expressions like $3 \sin x + 4 \cos x$ and for solving equations of the form $a \sin x + b \cos x = c$.

$R \sin(\theta + \alpha)$ form. For real constants a, b with at least one non-zero, $[a \sin \theta + b \cos \theta = R \sin(\theta + \alpha)]$ where $[R = \sqrt{a^2 + b^2} > 0, \text{ and } \tan \alpha = \frac{b}{a} \text{ (with } \alpha \text{ chosen so } \cos \alpha \text{ has the same sign as } a \text{)}]$. The maximum value of the expression is R (attained when $\theta + \alpha = \pi/2$) and the minimum is $-R$.

Derivation

Expand the target form using the compound-angle formula: $R \sin(\theta + \alpha) = R \sin \theta \cos \alpha + R \cos \theta \sin \alpha$. Matching coefficients of $\sin \theta$ and $\cos \theta$ in $a \sin \theta + b \cos \theta$ gives

$$a = R \cos \alpha, \quad b = R \sin \alpha.$$

Squaring and adding: $a^2 + b^2 = R^2(\cos^2 \alpha + \sin^2 \alpha) = R^2$, so $R = \sqrt{a^2 + b^2}$. Dividing the second equation by the first gives $\tan \alpha = b/a$.

Form	Expansion	Max	Min
$R \sin(\theta + \alpha)$	$R \cos \alpha \cdot \sin \theta + R \sin \alpha \cdot \cos \theta$	R at $\theta = \pi/2 - \alpha$	$-R$ at $\theta = -\pi/2 - \alpha$
$R \sin(\theta - \alpha)$	$R \cos \alpha \cdot \sin \theta - R \sin \alpha \cdot \cos \theta$	R at $\theta = \pi/2 + \alpha$	$-R$
$R \cos(\theta - \alpha)$	$R \cos \alpha \cdot \cos \theta + R \sin \alpha \cdot \sin \theta$	R at $\theta = \alpha$	$-R$

$R \cos(\theta + \alpha)$	$R \cos \alpha \cdot \cos \theta - R \sin \alpha \cdot \sin \theta$	$R \text{ at } \theta = -\alpha$	$-R$
---------------------------	---	----------------------------------	------

Exam tip — which form to pick. OCR questions usually tell you which target form to use ("express in the form $R \sin(\theta + \alpha)$ "). If the form is not specified, $R \sin(\theta + \alpha)$ matches $a \sin \theta + b \cos \theta$ cleanly and is the default. Make sure $R > 0$ and α is acute (in $[0, \pi/2]$) unless otherwise stated.

Worked Example 8 — Full $R \sin(\theta + \alpha)$ solution

Express $3 \sin \theta + 4 \cos \theta$ in the form $R \sin(\theta + \alpha)$, with $R > 0$ and $0 < \alpha < \pi/2$. Hence (a) state the maximum and minimum values, (b) solve $3 \sin \theta + 4 \cos \theta = 2$ for $0 \leq \theta \leq 2\pi$.

1. Match coefficients:
2. $a = 3, \quad b = 4.$
3. $R = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$
4. $\tan \alpha = 4/3 \Rightarrow \alpha = \arctan(4/3) \approx 0.9273 \text{ rad } (\approx 53.13^\circ).$
5. $\therefore 3 \sin \theta + 4 \cos \theta = 5 \sin(\theta + \alpha) \text{ with } \alpha = \arctan(4/3).$
6. (a) Maximum = $R = 5$, Minimum = $-R = -5$.
7. Max at $\theta + \alpha = \pi/2$, i.e. $\theta = \pi/2 - \arctan(4/3) \approx 0.6435 \text{ rad}.$
8. (b) Solve $5 \sin(\theta + \alpha) = 2 \Rightarrow \sin(\theta + \alpha) = 2/5 = 0.4.$
9. Principal value: $\theta + \alpha = \arcsin(0.4) \approx 0.4115 \text{ rad}.$
10. Second value (sin positive in Q2): $\theta + \alpha = \pi - 0.4115 \approx 2.7301 \text{ rad}.$
11. Add 2π to each for further solutions within shifted range.
12. Subtract $\alpha \approx 0.9273$:
13. $\theta_1 = 0.4115 - 0.9273 \approx -0.5158 \quad (\text{out of range; add } 2\pi)$
14. $\rightarrow 2\pi - 0.5158 \approx 5.7674 \text{ rad}$
15. $\theta_2 = 2.7301 - 0.9273 \approx 1.8028 \text{ rad} \quad (\text{in } [0, 2\pi]) \checkmark$
16. $\therefore \theta \approx 1.8028 \text{ rad} \quad \text{or} \quad 5.7674 \text{ rad} \quad (\text{to 4 d.p.}).$

// In degrees these are $\theta \approx 103.29^\circ$ and $\theta \approx 330.45^\circ$.

9. Solving Trigonometric Equations

Solving a trig equation means finding every angle that satisfies it — usually within a specified interval such as $[0, 2\pi]$ or $[0^\circ, 360^\circ]$, sometimes over all real numbers. The workflow is:

1. Manipulate the equation (using identities if necessary) until it has the form $\sin(\text{stuff}) = k$, $\cos(\text{stuff}) = k$ or $\tan(\text{stuff}) = k$.
2. Find the *principal value* using $\arcsin$, $\arccos$ or $\arctan$.
3. Use symmetry of the graph — or the CAST diagram — to list all other solutions in the range for the transformed argument.
4. Undo any substitution and check every candidate is in the stated interval.

The CAST Diagram

The CAST diagram summarises in which of the four quadrants each trig function is positive. Starting in the fourth quadrant and moving anticlockwise, the letters read C, A, S, T:

Quadrant	Angle range	Functions positive	Label
I	$0 \leq \theta < \pi/2$	$\sin, \cos, \tan$	A (All)
II	$\pi/2 \leq \theta < \pi$	$\sin$ only	S (Sine)
III	$\pi \leq \theta < 3\pi/2$	$\tan$ only	T (Tangent)
IV	$3\pi/2 \leq \theta < 2\pi$	$\cos$ only	C (Cosine)

Rules of thumb for moving from the principal value θ_0 to the others:

- If $\sin \theta = k$ with $k > 0$: solutions in Q1 and Q2, so $\theta = \theta_0$ and $\pi - \theta_0$.
- If $\cos \theta = k$ with $k > 0$: solutions in Q1 and Q4, so $\theta = \theta_0$ and $2\pi - \theta_0$.
- If $\tan \theta = k$: period π , so $\theta = \theta_0$ and $\theta_0 + \pi$.
- If k is negative, mirror the positive- k rules to the quadrants where the function is negative.

Exam tip — always state the range. The phrase "in the interval $0 \leq x \leq 2\pi$ " changes the answer. A solution outside the interval loses a mark, as does a missing solution inside. Sketch the graph or run through the CAST rule exhaustively, count solutions, and check your count matches

what the graph predicts (e.g. $\sin$ equation should have either 2 or 0 solutions per period, $\tan$ exactly 1).

Worked Example 9 — Linear argument

Solve $\sin 2x = \frac{\sqrt{3}}{2}$ for $0 \leq x \leq 2\pi$.

1. Let $u = 2x$, so u runs over $[0, 4\pi]$.
2. Principal value: $\arcsin(\sqrt{3}/2) = \pi/3$.
3. All solutions in one period $[0, 2\pi]$:
4. $u = \pi/3$ (Q1)
5. $u = \pi - \pi/3 = 2\pi/3$ (Q2)
6. Add 2π for second period (within $[0, 4\pi]$):
7. $u = 2\pi + \pi/3 = 7\pi/3$
8. $u = 2\pi + 2\pi/3 = 8\pi/3$.
9. Back-substitute $x = u/2$:
10. $x = \pi/6, \pi/3, 7\pi/6, 4\pi/3$.
11. $\therefore$ four solutions in $[0, 2\pi]$. ✓

Worked Example 10 — Reciprocal function

Solve $\sec x = 2$ for $0 \leq x \leq 2\pi$.

1. $\sec x = 2 \Leftrightarrow \cos x = 1/2$.
2. Principal value: $\arccos(1/2) = \pi/3$.
3. Solutions ($\cos > 0$ in Q1, Q4):
4. $x = \pi/3$ (Q1)
5. $x = 2\pi - \pi/3 = 5\pi/3$ (Q4).
6. $\therefore x \in \{\pi/3, 5\pi/3\}$.

Worked Example 11 — Quadratic in $\sin$

Solve $2\sin^2 x - 3\sin x + 1 = 0$ for $0 \leq x \leq 2\pi$.

1. Let $s = \sin x$: $2s^2 - 3s + 1 = 0 \Rightarrow (2s - 1)(s - 1) = 0$.
2. $s = 1/2$ or $s = 1$.
3. $\sin x = 1/2$: $x = \pi/6, 5\pi/6$.
4. $\sin x = 1$: $x = \pi/2$.
5. $\therefore x \in \{\pi/6, \pi/2, 5\pi/6\}$.

10. Inverse Trigonometric Functions

Trig functions are not one-to-one (they repeat), so strictly they have no inverse. To define an inverse we restrict the domain to a single monotonic piece. This gives the three *principal* inverse trig functions.

Inverse trig functions.

- $y = \arcsin x$: domain $[-1, 1]$, range $[-\pi/2, \pi/2]$. Equivalent to $\sin y = x$ with y in that range.
- $y = \arccos x$: domain $[-1, 1]$, range $[0, \pi]$. Equivalent to $\cos y = x$.
- $y = \arctan x$: domain $\mathbb{R}$, range $(-\pi/2, \pi/2)$. Equivalent to $\tan y = x$.

(Alternative notations: $\sin^{-1} x$, $\cos^{-1} x$, $\tan^{-1} x$. Note $\sin^{-1} x \neq 1/\sin x$ — the superscript -1 denotes function inverse, not reciprocal.)

Graphs

- $\arcsin$ rises from $(-1, -\pi/2)$ through the origin to $(1, \pi/2)$ — the reflection of the restricted $\sin$ in the line $y = x$.
- $\arccos$ decreases from $(-1, \pi)$ through $(0, \pi/2)$ to $(1, 0)$.
- $\arctan$ increases from the horizontal asymptote $y = -\pi/2$ through the origin to the asymptote $y = \pi/2$.

Useful Identities

$$\arcsin x + \arccos x = \frac{\pi}{2} \quad \text{for all } x \in [-1, 1].$$

$$\arctan x + \arctan(1/x) = \begin{cases} \pi/2, & x > 0 \\ -\pi/2, & x < 0 \end{cases}$$

Exam tip — restricted domains matter. $\arcsin(\sin \theta)$ equals θ only if $\theta \in [-\pi/2, \pi/2]$; otherwise you must reduce θ to the principal range first. For instance, $\arcsin(\sin(3\pi/4)) = \pi/4$, not $3\pi/4$, because $\sin(3\pi/4) = \sin(\pi/4)$ and $\pi/4$ is in range. OCR examiners frequently probe this subtlety.

Worked Example 12 — Evaluating inverse trig

Evaluate (a) $\arcsin(\sqrt{3}/2)$, (b) $\arccos(-1/2)$, (c) $\arctan(-1)$.

1. (a) $\arcsin(\sqrt{3}/2)$: need $y \in [-\pi/2, \pi/2]$ with $\sin y = \sqrt{3}/2 \Rightarrow y = \pi/3$.
2. (b) $\arccos(-1/2)$: need $y \in [0, \pi]$ with $\cos y = -1/2$.
3. $\cos$ is negative in Q2; reference angle is $\pi/3$ (since $\cos \pi/3 = 1/2$).
4. $y = \pi - \pi/3 = 2\pi/3$.
5. (c) $\arctan(-1)$: need $y \in (-\pi/2, \pi/2)$ with $\tan y = -1 \Rightarrow y = -\pi/4$.

11. Further Equations and Mixed Techniques

Worked Example 13 — Equation involving sec and tan

Solve $\sec^2 x - 3 \tan x - 1 = 0$ for $0 \leq x \leq 2\pi$.

1. Use $\sec^2 x = 1 + \tan^2 x$:
2. $(1 + \tan^2 x) - 3 \tan x - 1 = 0$
3. $\tan^2 x - 3 \tan x = 0$
4. $\tan x (\tan x - 3) = 0$.
5. $\tan x = 0$: $x = 0, \pi, 2\pi$.
6. $\tan x = 3$: principal value $\arctan 3 \approx 1.2490$ rad.
7. Period π , so $x \approx 1.2490$ and $x \approx 1.2490 + \pi \approx 4.3906$.

8. $\therefore x \in \{0, 1.2490, \pi, 4.3906, 2\pi\}$ (with endpoints 0 and 2π included if the interval is closed).

Worked Example 14 — $a \sin \theta + b \cos \theta = c$ using R-form

Solve $5 \sin \theta - 12 \cos \theta = 6$ for $0 \leq \theta \leq 2\pi$.

1. Choose the target form $R \sin(\theta - \alpha)$, with α acute, $R > 0$, so that $R \sin(\theta - \alpha) = R \cos \alpha \sin \theta - R \sin \alpha \cos \theta$.
 2. Matching: $R \cos \alpha = 5$, $R \sin \alpha = 12$.
 3. $R = \sqrt{5^2 + 12^2} = \sqrt{169} = 13$.
 4. $\tan \alpha = 12/5 \Rightarrow \alpha = \arctan(12/5) \approx 1.1760$ rad.
 5. Equation becomes: $13 \sin(\theta - \alpha) = 6 \Rightarrow \sin(\theta - \alpha) = 6/13$.
 6. Principal: $\theta - \alpha = \arcsin(6/13) \approx 0.4797$ rad.
 7. Second value: $\theta - \alpha = \pi - 0.4797 \approx 2.6619$ rad.
 8. Add back α :
 9. $\theta \approx 0.4797 + 1.1760 = 1.6557$ rad
 10. $\theta \approx 2.6619 + 1.1760 = 3.8379$ rad.
 11. Both lie in $[0, 2\pi]$, so there are exactly two solutions.
 12. $\therefore \theta \approx 1.6557$ rad or $\theta \approx 3.8379$ rad.
- // Check: $|6| \leq R = 13 \checkmark$ (so $|6/13| \leq 1$), hence solutions exist.
 // If the right-hand side exceeded R in absolute value, no solutions at all.

Worked Example 15 — Identity proof

Prove that $\frac{1 - \cos 2x}{\sin 2x} = \tan x$.

1. Numerator: $1 - \cos 2x = 1 - (1 - 2 \sin^2 x) = 2 \sin^2 x$.
2. Denominator: $\sin 2x = 2 \sin x \cos x$.
3. Ratio:
4. $(2 \sin^2 x)/(2 \sin x \cos x) = \sin x / \cos x = \tan x. \quad \checkmark$

// Valid wherever $\sin 2x \neq 0$ and $\cos x \neq 0$, i.e. $x \neq k\pi/2$.

12. Consolidated Summary and Exam Strategy

Chapter-wide checklist for Trigonometry.

- **Units first.** Are angles in degrees or radians? Set calculator accordingly. Small-angle and arc-length formulae assume radians.
- **Exact values at 0, 30°, 45°, 60°, 90°** must be automatic — no calculator needed.
- **Three Pythagorean identities:** $\sin^2 + \cos^2 = 1$, $\tan^2 + 1 = \sec^2$, $1 + \cot^2 = \csc^2$. Know which to deploy by looking at which functions already appear.
- **Compound angle** gives all other trig identities on the syllabus. If stuck, expand everything in $\sin$ and $\cos$ of single angles.
- **Double angle:** $\sin 2A = 2 \sin A \cos A$; $\cos 2A$ has three equivalent forms — pick the one matching the rest of the equation.
- **R sin($\theta + \alpha$):** $R = \sqrt{a^2 + b^2}$, $\tan \alpha = b/a$. Useful for optimisation (max = R , min = $-R$) and solving $a \sin \theta + b \cos \theta = c$ — solvable iff $|c| \leq R$.
- **Solving trig equations:** find principal value, use CAST / graph to list others in range, never divide by a quantity that could be zero (factor instead).
- **Inverse trig:** respect the restricted range; $\arcsin(\sin \theta) = \theta$ only for $\theta \in [-\pi/2, \pi/2]$.
- **State the interval** and list every solution within it; check by counting how many you expect from the period.

Typical OCR examiner complaints (from mark reports).

- Calculator in the wrong mode — degrees vs radians confusion.
- Dividing through by $\cos x$ (or similar) without checking $\cos x = 0$, losing half the solutions.
- Missing the second solution in the interval when $\sin$ or $\cos$ has two symmetric values per period.

- Using $\sin x \approx x$ for x in degrees — wrong by a factor of $\pi/180$.
- Forgetting to factor out R or using $\tan \alpha = a/b$ instead of b/a .
- Writing $\cos(A + B) = \cos A \cos B + \sin A \sin B$ — sign error: it's a minus.
- Giving $\arcsin(\sin(3\pi/4)) = 3\pi/4$ instead of $\pi/4$ — ignoring the principal range.
- Leaving an interval as $[0^\circ, 360^\circ]$ and giving answers in radians, or vice-versa.

Formula snapshot (memorise).

- Radian/degree: $180^\circ = \pi$ rad. Arc $s = r\theta$. Sector $A = \frac{1}{2}r^2\theta$. Segment area $\frac{1}{2}r^2(\theta - \sin \theta)$.
- Small-angle: $\sin x \approx x$, $\cos x \approx 1 - x^2/2$, $\tan x \approx x$ (x in radians).
- Pythagorean: $\sin^2 \theta + \cos^2 \theta = 1$; $\sec^2 \theta = 1 + \tan^2 \theta$; $\csc^2 \theta = 1 + \cot^2 \theta$.
- Compound: $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$; $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$; $\tan(A \pm B) = (\tan A \pm \tan B)/(1 \mp \tan A \tan B)$.
- Double: $\sin 2A = 2 \sin A \cos A$; $\cos 2A = \cos^2 A - \sin^2 A = 1 - 2 \sin^2 A = 2 \cos^2 A - 1$; $\tan 2A = 2 \tan A/(1 - \tan^2 A)$.
- Power-reducing: $\sin^2 A = (1 - \cos 2A)/2$; $\cos^2 A = (1 + \cos 2A)/2$.
- R-form: $a \sin \theta + b \cos \theta = R \sin(\theta + \alpha)$ with $R = \sqrt{a^2 + b^2}$, $\tan \alpha = b/a$; max R , min $-R$.
- Inverse: $\arcsin$ has range $[-\pi/2, \pi/2]$; $\arccos$ has range $[0, \pi]$; $\arctan$ has range $(-\pi/2, \pi/2)$; $\arcsin x + \arccos x = \pi/2$.