RAG Learning

AQA GCSE Geography (8035) — Paper 1: Living with the Physical Environment · 3.1.2 The Living World (option: hot deserts)

1.5 Hot Deserts

1. Location and Climate

Hot deserts are found predominantly at approximately **30° North and 30° South** of the equator, in belts corresponding to the **subtropical high pressure zones**. They also occur in **rain shadows** (sheltered side of mountain ranges) and deep **continental interiors** far from moisture sources.

> **Desert Definition:** An area receiving less than **250 mm of rainfall per year**. Hot deserts experience extreme heat by day and cold at night due to the absence of cloud cover and low humidity.

Climate Characteristics

Variable	Value	Detail
Annual rainfall	<250 mm/year (often <25 mm)	Rain is rare, unpredictable, and often arrives as intense short bursts causing flash floods
Daytime temperature	40–50°C in summer	No cloud cover means intense solar radiation reaches the ground directly
Night-time temperature	Can drop to 5–10°C	No cloud cover also means heat is rapidly lost at night via radiation — no "blanket" effect
Diurnal temperature range	Up to 40°C difference	Very large daily swing — extreme physical weathering (exfoliation/insolation weathering)

© RAG Learning · Not for redistribution

Humidity	Very low (<20%)	Low humidity accelerates evaporation; any moisture is rapidly lost to the atmosphere


Major Hot Desert Examples

Desert	Location	Notable Feature
Sahara	North Africa	World's largest hot desert (~9.2 million km²); only ~25% is sand — rest is rock (reg/hammada)
Arabian	Arabian Peninsula (Saudi Arabia, Oman, UAE)	World's largest sand desert (Rub' al Khali — Empty Quarter)
Atacama	West coast South America (Chile, Peru)	World's driest — some areas have had no recorded rainfall for centuries
Sonoran	SW USA and NW Mexico	Home of the saguaro cactus; most biodiverse desert
Australian Outback	Central Australia	Covers ~70% of Australia's land area

2. Causes of Desert Climate

The location of hot deserts at ~30°N/S is explained by the **Hadley Cell** — a large atmospheric circulation pattern driven by the sun heating the equator.

© RAG Learning · Not for redistribution



Other Causes of Desert Climate

Cause	Mechanism	Example
Subtropical high pressure	Hadley cell — dry air descends at 30°N/S, warms adiabatically, inhibits cloud formation and rainfall	Sahara, Arabian, Thar deserts
Rain shadow effect	Moist air rises over mountains, deposits rain on windward side; descends dry and warm on leeward side	Atacama Desert (behind the Andes); Patagonian Desert (behind Andes)
Distance from sea (continentality)	Air masses lose moisture as they travel inland; no maritime moisture source	Gobi Desert (Central Asia)
Cold ocean currents	Cold water chills overlying air, preventing it from rising and forming rain clouds; stable cold air = no convection	Atacama — cooled by the Humboldt Current; Namib — Benguela Current

© RAG Learning · Not for redistribution

3. Desert Weathering Processes

Weathering breaks rock down in situ (without moving it). In deserts, **mechanical (physical) weathering** dominates because of the extreme temperature ranges and the absence of vegetation roots. **Chemical weathering** also occurs, despite the apparent lack of water, whenever thin moisture films are present.

Mechanical (Physical) Weathering

> **Mechanical weathering:** The physical breakdown of rock into smaller fragments without any change to the rock's chemical composition. The rock splits or flakes but remains chemically identical to the parent rock.

Process	Mechanism (Step-by-Step)	Result
Insolation weathering (thermal expansion and contraction)	1. Intense solar radiation heats the outer rock surface to very high temperatures (up to 80°C at surface). 2. Rock expands as temperature rises. 3. At night, rapid heat loss causes the surface to contract. 4. The outer surface and inner rock expand/contract at different rates (rock is a poor conductor of heat, so inside stays cooler). 5. Over thousands of cycles, tensional stress builds up in the outer layers. 6. Eventually sheets or flakes crack off parallel to the surface.	Exfoliation (onion-skin weathering) — curved sheets of rock peel away; rounded boulders form; also causes granular disintegration where individual mineral grains loosen and fall away
Salt crystal growth (haloclasty)	1. Desert soils/rocks contain dissolved salts (sodium chloride, sodium sulphate, sodium nitrate). 2. When water (from dew, rare rain, or groundwater capillary action) evaporates, salts crystallise within rock pores and joints. 3. Crystal growth exerts pressures of up to 200 MPa within the pore — far exceeding the tensile strength of most rocks. 4. Repeated wetting and drying cycles cause progressive cracking.	Rock disintegrates from within; produces granular rubble and sand; particularly effective in coastal deserts where sea spray provides additional salt; a major weathering agent across the Sahara and Atacama

© RAG Learning · Not for redistribution

Frost shattering (freeze-thaw)	1. In hot deserts, nights can drop below 0°C. 2. Any water in rock joints freezes and expands by ~9% in volume. 3. Pressure (up to 2,000 kg/cm²) forces joints apart. 4. Repeated freeze-thaw cycles progressively widen cracks.	Angular rock fragments (scree); more significant at altitude or in cold desert margins; contributes to talus slopes at base of desert cliffs and escarpments
Pressure release (dilatation)	Deep rocks formed under enormous pressure; when overlying rock is eroded away, the pressure is released and the rock expands upward; curved joints develop parallel to the surface.	Exfoliation domes; inselbergs; contributes to the formation of tors and granite outcrops in desert landscapes

Chemical Weathering

Although deserts are dry, chemical weathering is not absent. Dew, fog, and rare rain provide enough moisture to drive chemical reactions, particularly at night or in coastal deserts where fog is frequent (e.g. the Atacama receives significant fog moisture from the Pacific — called "camanchaca").

Process	Mechanism	Evidence in Deserts
Hydration	Minerals absorb water molecules and expand; feldspar crystals in granite absorb water and swell, forcing grains apart. No actual chemical change to mineral composition — a physical-chemical boundary process.	Granular disintegration of granite rocks; produces sand-sized particles of feldspar and quartz
Oxidation	Iron-bearing minerals (e.g. iron pyrite, olivine) react with oxygen in the presence of thin moisture films; iron is oxidised to iron oxide (rust). $Fe^{2+} \rightarrow Fe^{3+}$ (iron oxide)	Red/orange colouring of desert rocks and soils (ferricrete); characteristic red sandstone landscapes of the Australian Outback and US Southwest

© RAG Learning · Not for redistribution

Hydrolysis	Weak carbonic acid (CO_2 dissolved in water) reacts with silicate minerals (especially feldspar), breaking them down into clay minerals and releasing silica in solution. This is the dominant chemical weathering process globally.	Rock varnish (dark patina of manganese and iron oxides); clay-lined wadis; desert caliche (calcium carbonate crust in soil)
Solution (carbonation)	Slightly acidic rainwater dissolves calcium carbonate in limestone rocks: $CaCO_3 + H_2O + CO_2 \rightarrow Ca(HCO_3)_2$	Limestone landscapes in desert fringes; calcrete formation; cave systems in desert limestone plateaus

Exam Tip — Weathering: Many students confuse weathering and erosion. **Weathering** = breakdown in situ (no movement). **Erosion** = wearing away AND transporting material. A question asking "explain how a desert landform is created" usually requires you to describe BOTH the weathering that weakens/breaks down rock AND the erosion/transportation that removes material to create the landform's shape.

4. Desert Landscapes and Landforms

Despite the popular image of endless sand dunes, only about **25% of the Sahara is erg (sand desert)**. Most desert landscapes are dominated by bare rock, gravel plains, and distinctive erosional landforms shaped by **wind (aeolian processes)** and rare but intense **water action**.

Wind Erosion Processes

Process	Mechanism	Result
Deflation	Wind picks up and removes loose, fine material (sand and dust) from the surface. Strong desert winds can carry particles at speeds above 30 km/h. Dust can be transported thousands of kilometres —	Deflation hollows (depressions in desert surface); desert pavements (reg/serir) where fine material removed leaving behind coarse lag gravel; deflation may expose fossil groundwater

	Saharan dust regularly reaches the UK and the Amazon rainforest.	
Abrasion (sandblasting)	Wind-carried sand particles act like sandpaper, grinding and polishing rock surfaces. Abrasion is most intense within 1 metre of the ground where sand transport is highest (saltation layer). This is why many desert landforms show undercutting near the base.	Smooth, polished rock surfaces; ventifacts (stones shaped by wind-blasting, often with facets); undercut cliffs; mushroom rocks (pedestal rocks)
Attrition	Sand particles collide with each other and with rock surfaces during transport; particles become progressively smaller and rounder.	Rounded, frosted sand grains (characteristic of aeolian sand under microscope — contrasts with angular water-transported grains)

Erosional Landforms


Landform	Formation	Appearance
Wadi	Dry river valley carved by rare but intense flash floods; water evaporates or soaks in quickly after rain	Flat-bottomed, steep-sided valley; dry most of the year but dangerous during rare rainstorms
Yardang	Wind erodes soft rock more rapidly than hard rock; wind-aligned ridges and troughs form parallel to prevailing wind	Long, streamlined ridges aligned with wind direction; can be hundreds of metres long
Zeugen (Tableland)	Horizontal layers of hard rock over soft rock; wind erosion undercuts the soft layer; hard cap protects the top	Flat-topped mesa or table rock with undercut base; eventually collapses when soft layer erodes completely
Desert pavement (reg/serir)	Wind removes fine sand, leaving behind a concentrated layer of coarse gravel and pebbles	Flat, stone-covered surface with a "mosaic" appearance;

© RAG Learning · Not for redistribution

		common across most of the Sahara
Mesa and Butte	Both formed from horizontal rock strata where a resistant cap rock (e.g. sandstone) protects softer underlying layers. A **mesa** is a large, wide flat-topped hill — erosion has removed surrounding material but the cap rock protects a large area. A **butte** is a smaller version of a mesa, typically narrower than it is tall. As erosion continues over millions of years: plateau → mesa → butte → isolated pillar → eventually removed.	Iconic flat-topped hills with near-vertical cliffs; mesas can be kilometres wide; buttes are narrow, tower-like; examples: Monument Valley, Utah; Ahaggar Mountains, Sahara
Inselberg	An isolated, steep-sided rocky hill rising abruptly from a flat desert plain (pediplain). Formed by deep chemical weathering along joint systems below the surface (when the landscape was wetter). The more jointed surrounding rock weathers faster; over millions of years, erosion strips away the weathered material, leaving the resistant core standing proud. Process called *pediplanation*. Can also form by pressure release doming in granite (dilatation).	Dramatic isolated hills of bare rock rising sharply from flat plains; may be smooth and rounded (granite bornhardts) or irregular; examples: Uluru (Ayers Rock), Australia; Brandberg Mountain, Namibia; inselbergs of the Sahara
Pediment	A gently sloping (2–7°) bedrock surface at the foot of a desert mountain or scarp. Formed by a combination of sheet flooding (thin sheets of water after rain) washing material across the surface, lateral erosion by streams cutting into the hillslope base, and weathering retreating the hillside backwards. The pediment is a rock-cut surface, often thinly covered with debris.	Smooth, gently inclined rock ramp connecting a steep rocky highland to a flat basin; often covered by a thin veneer of gravel; found at the base of most desert mountain ranges

Depositional Landforms

© RAG Learning · Not for redistribution



Landform	Type	Description
Barchan dune	Depositional (wind)	Crescent-shaped sand dune; horns point downwind. Forms where sand supply is limited. Can move ~15–20m/year — threatening settlements and roads.
Seif dune	Depositional (wind)	Long, linear dune parallel to prevailing wind. Can be hundreds of km long, tens of metres high. Form where wind direction varies seasonally.
Reg (stone desert)	Depositional/lag	Flat gravel/pebble surface left after wind removes all fine sand particles. Sometimes called "hammada" when the surface is bare bedrock.
Alluvial fan	Depositional (water)	A fan-shaped deposit of sediment where a stream or wadi exits a mountain canyon and enters a flat plain. The sudden loss of slope and confinement causes water to spread out and slow, dropping its sediment load. Coarser material (boulders, gravel) deposits at the top of the fan near the canyon mouth; fine silt deposits at the outer edge. Multiple fans merging together along a mountain front create a *bajada*. Examples: Death Valley alluvial fans (USA); Atacama piedmont fans.

Exam Tip — Landform Questions: For any landform, structure your answer around: (1) the type of rock and its resistance, (2) the process that erodes/deposits material, (3) the sequence of events that creates the shape. Landforms like mesas/buttes are excellent for showing understanding of

© RAG Learning · Not for redistribution

how *differential erosion* (hard rock protects soft rock) creates distinctive shapes. Inselbergs and pediments often appear in 6-mark questions asking you to describe and explain formation.

5. Water Action in Deserts

Water is paradoxically one of the most important agents shaping desert landscapes, despite deserts being defined by lack of rainfall. When rain does fall, it tends to arrive as **intense, short-duration convectional rainstorms**. Because desert soils are poorly developed, lack vegetation, and are often sealed by a hard surface crust, **infiltration is very low** — most water runs off immediately as surface flow.

Flash Floods and Wadis

Wadi: A dry river channel (or valley) found in desert environments. Most of the year they are completely dry, but during and after rare rainstorms they fill rapidly with fast-moving, sediment-laden floodwater. The word is Arabic and is used across the Middle East and North Africa. Equivalent terms: "arroyo" (SW USA/Mexico), "nullah" (South Asia).

Flash floods in wadis are among the most dangerous natural hazards in desert regions. Key characteristics:

- **Speed of onset:** Floodwaters can arrive with little or no warning — a storm 50 km away in mountains can send a wall of water down a normally dry wadi within minutes. A person in a wadi may hear a roaring sound before seeing the flood.
- **Sediment load:** Flash floods carry enormous amounts of sediment — the water appears brown or red and has a consistency almost like wet concrete at peak flow. Boulders are transported that would never move in normal river flow (high competence due to high velocity).
- **Duration:** Floods typically last only a few hours, then the water drains away rapidly or evaporates, leaving behind thick deposits of sand and gravel on the wadi floor.
- **Erosional power:** Repeated flash floods carve the characteristic flat-bottomed, steep-sided wadi profile. The flat floor results from lateral erosion and deposition during floods; the steep sides result from undercutting and collapse.

© RAG Learning · Not for redistribution



Alluvial Fans — Formation and Significance

When a wadi exits a confined mountain valley and reaches an open flat basin or plain, the flow loses velocity dramatically. The sudden change spreads water across a wide area. Sediment drops out of suspension in a characteristic fan shape:

- **Apex** (top, near canyon mouth): Coarse material — boulders, cobbles, coarse gravel. Deposited first as velocity drops rapidly.
- **Mid-fan:** Medium gravel and sand. Water has slowed further and spread out.
- **Toe** (outer edge of fan): Fine silt and clay; may pond into temporary desert lakes (playas).

Where multiple alluvial fans merge along a mountain front, they coalesce into a continuous sloping apron called a **bajada**. Alluvial fans are significant because they can contain freshwater aquifers (water soaks into the fan gravels during floods) and are often used as settlement sites in desert regions.

Desert Lakes — Playas

Playa: A flat desert basin that occasionally fills with shallow water after flash floods or heavy rain, then dries out to leave a flat, salt-encrusted surface. Also called a "sabkha" (Arabian Peninsula), "chott" (North Africa), or "salt flat." Formed in enclosed basins where water collects but cannot drain — evaporation concentrates minerals, leaving behind evaporite deposits (salt, gypsum, borax). Salar de Uyuni (Bolivia) is the world's largest salt flat at 10,582 km².

Exam Tip — Water in Deserts: A common misconception is that water does not shape desert landscapes. In fact, most geographers argue water action (fluvial processes) is MORE important than wind in shaping desert landforms overall — wind shapes dunes and creates yardangs, but

© RAG Learning · Not for redistribution

water shapes the wider landscape through wadis, alluvial fans, pediments, and playas. Make this point explicitly in high-mark answers.

6. Comparing Hot Deserts: Atacama vs Sahara

The Atacama and Sahara are both hot deserts but differ significantly in their causes, characteristics, and human use. Knowing the contrast is useful for evaluation questions.

Feature	Atacama (South America)	Sahara (North Africa)
Area	~105,000 km² (relatively small)	~9.2 million km² (world's largest hot desert)
Location	Narrow coastal strip, western South America; mainly Chile and Peru; 15°S–30°S	Stretches across North Africa from Atlantic coast to Red Sea; 15°N–35°N
Primary cause of aridity	Three reinforcing factors: (1) Rain shadow of the Andes (moist easterly air from the Amazon/Atlantic drops its moisture on the eastern Andes before reaching the Atacama). (2) Cold Humboldt Current (upwelling cold water cools overlying air, creating stable temperature inversion that suppresses rainfall). (3) Subtropical high pressure.	Primarily subtropical high pressure (Hadley Cell). Also continental interior away from Atlantic moisture. Trade winds blow across hot land before reaching desert interior.
Annual rainfall	<1 mm/year in driest areas (Arica has recorded 0 mm for over 14 consecutive years); some weather stations have never recorded measurable rain. World's driest non-polar desert.	Highly variable: 0–25 mm/year in central Sahara; up to 100 mm near margins. More variable and slightly higher than Atacama average.
Temperature	Cooler than typical hot desert due to cold Humboldt Current and coastal upwelling — coastal Atacama rarely exceeds 30°C; interior (at altitude) can be colder. Diurnal range large at altitude (4,000m+).	Extreme heat — summer maxima regularly 48–50°C in central Sahara; a reading of 58°C at Aziziya, Libya (1922)

© RAG Learning · Not for redistribution

		was invalidated by the WMO in 2012. Diurnal range up to 40°C.
Moisture source	Camanchaca (coastal fog) from Pacific provides limited moisture to coastal zones; supports unique fog-dependent ecosystems at 500–1,000m altitude.	Occasional incursions of Mediterranean moisture in north; rare tropical convection from south in wet years. Dust storms carry moisture long distances.
Vegetation	Sparse but unusual: lomas (fog-oasis plant communities); cacti at altitude; hypolithic algae under translucent rocks (survive using tiny amounts of light). Some areas are completely barren — the absolute Atacama.	More varied at margins: date palms in oases, acacia scrub in Sahel transition; wadis support ephemeral vegetation after rain; true Sahara core mostly barren.
Key resources	World's largest copper reserves (Escondida mine — 5% of world copper); world's largest lithium reserves in salt flats (Atacama Salt Flat); historic nitrate (saltpetre) deposits (guano). Critical for EV batteries — global strategic importance.	Oil and natural gas (Libya, Algeria, Egypt); iron ore; phosphates (Morocco — world's largest); gold. Solar potential enormous.
Human settlement	Sparse; mine workers in camps; ancient indigenous cultures (Atacameno people) used oases and llama herding. Pre-Inca cultures developed sophisticated irrigation (puquios — ancient spiral aqueducts).	Nomadic Bedouin and Tuareg peoples historically; oasis settlements (Siwa, Ghadames); modern cities only with oil wealth (Tripoli, Algiers near margins).
Key challenge	Mining uses enormous amounts of water in a near-waterless environment; lithium extraction processes deplete brine aquifers that sustain local wetlands (flamingo habitat). Environmental controversy.	Desertification threatening Sahel margins; sand dune encroachment on settlements and infrastructure; political instability across Sahel.

Exam Tip — Atacama vs Sahara: If asked to compare two desert environments, use the structure: similarity → difference → evaluation. For example: "Both receive <250mm rainfall (similarity),

© RAG Learning · Not for redistribution

however the Atacama is the world's driest desert receiving <1mm/year due to the triple effect of the Humboldt Current, Andean rain shadow, and subtropical high pressure, while the Sahara receives up to 25mm due to Hadley Cell alone (difference). This means the Atacama supports almost no life in its core while Sahara margins support sparse vegetation (evaluation/consequence)."

7. Plant Adaptations (Xerophytes)

Xerophyte: A plant that has evolved structural or physiological adaptations to survive in environments with very little water. The name comes from Greek: xero (dry) + phyte (plant).

Plant / Adaptation	Mechanism	Benefit
Cactus — thick, fleshy stem	Stem is enlarged to store water (up to 750 litres in a large saguaro). Accordion-shaped ribs expand when water is absorbed.	Stores water during rare rainfall events to survive months or years of drought
Cactus — spines instead of leaves	Leaves have been replaced entirely by spines (modified leaves)	Spines have very low surface area — dramatically reduces water loss by transpiration. Also deters animals from eating the plant for its water.
Waxy cuticle	Thick waterproof coating on leaf/stem surfaces	Prevents evaporation from plant surfaces between rainfall events
Deep tap roots	Some desert plants grow roots 10–50m deep	Reaches permanent water table deep underground; accesses moisture between rare rain events
Wide, shallow roots	Roots spread out to cover a large area just below surface	Captures maximum rainfall from even small showers before it evaporates; saguaro roots cover an area equal to the above-ground canopy

© RAG Learning · Not for redistribution

Drought-avoiding seeds	Seeds germinate only when specific rainfall threshold is met; can remain viable in soil for years	Ensures germination and rapid flowering only when water is available; desert blooms after rare rain
CAM photosynthesis	Stomata open only at night to take in CO_2; photosynthesis occurs during the day with stored CO_2	Prevents water loss from open stomata during the hot day; can reduce water use by up to 90% compared to normal photosynthesis
Pale/silvery colouring	Light-coloured leaves reflect solar radiation	Reduces heating of leaf surface; lowers temperature difference between plant and air

8. Animal Adaptations

Animal	Structural Adaptations	Behavioural Adaptations
Camel	Fat stored in hump (not water — released for energy when food scarce); oval red blood cells function even when dehydrated; wide feet prevent sinking in sand; long eyelashes and closable nostrils for sandstorms; can tolerate body temperature varying from 34–41°C	Rests in shade during peak heat; faces sun to minimise exposed surface area; can drink 115 litres in 15 minutes when water is found
Fennec fox	Very large ears (up to 15cm) with thin skin and many blood vessels near surface	Nocturnal — active only at night; burrows deep in sand during day to escape heat
Sidewinder rattlesnake	Flattened body; heat-sensing pit organs to detect prey at night	Sidewinding movement minimises contact with hot sand; nocturnal; burrows during day
Desert tortoise	Thick, insulating shell; stores water in bladder (can store up to 40% of body weight as dilute urine)	Active only at dawn/dusk; hibernates during coldest months; digs shallow burrows

© RAG Learning · Not for redistribution

Kangaroo rat	Concentrated urine (minimises water loss); efficient kidneys; obtains most water from metabolising dry seeds	Strictly nocturnal; never needs to drink liquid water — metabolic water from seeds is sufficient

Exam Tip: When asked about adaptations, always explain the *mechanism* — don't just say "camels store fat." Say: "Camels store fat in their hump which is broken down to release energy (and water as a metabolic by-product) when food and water are scarce." Explaining HOW the adaptation works gains full marks.

9. Interdependence in the Hot Desert Ecosystem

The parts of a hot desert ecosystem — climate, soils, water, plants, animals and people — depend closely on one another, so a change to one affects the others.

- **Climate → plants:** very low, unreliable rainfall means plants are sparse and adapted to drought (xerophytes), with long roots and small or waxy leaves.
- **Plants → soils:** sparse vegetation adds little leaf litter, so soils are thin, sandy or stony, and low in organic matter — but rich in salts drawn up by evaporation.
- **Plants → animals:** plants provide food, moisture and shade for animals; animals in turn spread seeds (e.g. birds and rodents) and fertilise the soil with dung.
- **Water → everything:** oases and wadis support dense clusters of plants, animals and people. When rare rains fall, ephemeral plants flower rapidly, triggering a short burst of insect, bird and mammal activity.
- **People → ecosystem:** overgrazing by goats and camels removes vegetation, exposing soil to wind erosion; over-abstraction of groundwater dries out oases.

Exam Tip: For "explain the interdependence" questions, write chains of cause and effect: "Low rainfall limits plant growth, which means little organic matter is added to the soil, so soils are infertile, which in turn limits the number of animals the area can support."

10. Development Opportunities

© RAG Learning · Not for redistribution

Despite harsh conditions, hot deserts offer significant economic opportunities that are increasingly being exploited as technology advances.

Opportunity	Detail	Example
Mineral and oil extraction	Desert rocks contain vast mineral wealth formed over millions of years. Lack of vegetation makes extraction easier. Oil and gas formed from ancient organic deposits.	Saudi Arabia, Libya, Algeria (oil/gas); Chile Atacama (lithium, copper, nitrates); Western Australia (iron ore, gold)
Solar energy	Deserts receive ~3,000+ hours of sunshine per year — far more than temperate regions. Large, flat, unpopulated land available for solar farms.	Sahara solar potential: enough to power Europe 10x over. Noor Ouarzazate Solar Complex (Morocco) — one of the world's largest concentrated solar power plants. DESERTEC project proposal.
Tourism	Unique landscapes, cultural heritage, extreme environments attract adventure tourism and cultural visitors.	Wadi Rum (Jordan) — UNESCO site, 4WD safaris, Bedouin camps; Namib Desert; Dubai desert tours; Sahara camel treks
Agriculture with irrigation	Hot deserts receive intense sunlight ideal for crop growth IF water is supplied. Centre-pivot irrigation uses deep aquifer water.	Nile Valley, Egypt (year-round crops); Saudi Arabia used to grow wheat in desert using fossil water aquifers; California's Imperial Valley (Colorado River water)

11. Tourism: Opportunity and Challenge

Tourism in hot deserts has grown rapidly since the 1990s, driven by rising global incomes, improved transport links to remote areas, and growing demand for "adventure" and "extreme" experiences. Desert tourism is now a significant source of income for countries including the UAE, Jordan, Morocco, Namibia, and Peru (Atacama).

Opportunities from Desert Tourism

- **Economic income:** Tourism generates foreign exchange, taxes, and employment for local communities. In Jordan, Wadi Rum tourism generates millions of dollars annually; local Bedouin

families run camps and guided tours.

- **Infrastructure development:** Tourism investment often funds road construction, airport expansion, and utilities (electricity, communications) that also benefit local populations.
- **Cultural preservation:** Tourist interest in traditional cultures (Bedouin, Tuareg, Berber) can incentivise communities to maintain traditional practices, crafts, language, and music that might otherwise be lost to modernisation.
- **Conservation funding:** Ecotourism revenues can fund national park management, anti-poaching, and habitat restoration. Namibia's community conservancies, partly funded by safari tourism, have seen significant wildlife recovery (desert-adapted lions, elephants, rhinos).
- **UAE/Dubai example:** Dubai's desert tourism (dune bashing, camel safaris, falconry, desert dining camps) forms a key part of its diversification away from oil dependency. The Dubai Desert Conservation Reserve (225 km²) protects desert biodiversity while enabling tourism.

Challenges from Desert Tourism

- **Environmental damage:** Off-road vehicles (4x4s, quad bikes) compact desert soils, destroy fragile cryptobiotic soil crusts (which take decades to re-form), and crush vegetation. Tyre tracks in Peruvian deserts have damaged ancient Nazca Lines.
- **Water consumption:** Tourist hotels and resorts in desert cities (Dubai, Las Vegas, Marrakech) consume enormous quantities of water for swimming pools, golf courses, and gardens — deepening water scarcity problems. Dubai uses ~550 litres/person/day (far above global average of 170 litres).
- **Litter and waste:** Remote desert areas have no waste management infrastructure; tourist sites accumulate plastic waste, particularly along major desert trekking routes.
- **Cultural conflict:** Mass tourism can undermine traditional cultures — traditional nomadic lifestyles commercialised into inauthentic "shows" for tourists; sacred sites can be damaged by visitor footfall (e.g. climbing Uluru was banned in 2019 due to Aboriginal spiritual significance and environmental damage).
- **Carbon footprint:** Desert tourism typically requires long-haul flights, air-conditioned vehicles, and energy-intensive accommodation — a high carbon cost per tourist visit.
- **Carrying capacity:** Fragile desert ecosystems have a low carrying capacity — even small numbers of visitors can cause irreversible damage to rare species, rock formations, and archaeological sites. The Atacama's famous "flowering desert" (desierto florido) events have attracted huge tourist numbers that trample the very blooms visitors come to see.

© RAG Learning · Not for redistribution

Exam Tip — Tourism as Opportunity and Challenge: Questions may ask you to "assess the extent to which tourism is an opportunity for desert areas." Structure your answer: explain the opportunities (with place-specific examples and data), then explain the challenges/costs, then make a justified conclusion. A strong conclusion might argue that *ecotourism* (small-scale, community-managed, environmentally sensitive) offers the best balance, while mass tourism does more harm than good.

12. Development Challenges

Challenge	Impact	Example
Extreme heat	Dangerous working conditions (heatstroke, dehydration); requires air conditioning (high energy cost); equipment overheats	Construction workers in Dubai/Qatar work limited hours in summer; a 2021 investigation reported ~6,500 South Asian migrant worker deaths in Qatar in the decade after it won the World Cup bid (not all on World Cup sites; heat was thought to be a factor in many)
Water scarcity	All water must be imported, piped from distant sources, or produced by desalination (very expensive — $1–2/m³ vs $0.10 for freshwater)	Saudi Arabia relies on desalination for ~70% of freshwater; fossil water aquifers are being depleted irreversibly
Remote location / poor infrastructure	High transport costs for workers, equipment, and food supplies; isolation reduces labour supply	Mining operations in the Atacama or Australian outback require fly-in-fly-out (FIFO) workers with high wages
Sandstorms	Damage precision equipment, block air intakes, reduce visibility to near zero; can last days	Haboob sandstorms in Sudan/Egypt can be 1km high; cause millions of dollars of damage to oil infrastructure annually

© RAG Learning · Not for redistribution

Fragile ecosystem / environmental risk	Desert ecosystems are extremely fragile and slow to recover from disturbance; oil spills in desert soils can persist for decades	Oil pollution in Kuwait after Gulf War fires (1991) still visible in soils; camel grazing destruction in Australia's arid interior

Water Management in Deserts — Saudi Arabia in Depth

Saudi Arabia represents one of the most extreme examples globally of the tension between water demand and desert environment. It is almost entirely desert (the Rub' al Khali alone covers 650,000 km²) yet supports a population of ~35 million and aspirations for major economic development.

> **Fossil water (non-renewable aquifer):** Groundwater stored in underground rock formations (aquifers) over thousands to millions of years during past wetter climatic periods. It is called "fossil" because it is not currently being recharged by rainfall — extracting it is effectively mining a finite resource that cannot be replaced on any human timescale.

- **The Saudi fossil water situation:** Saudi Arabia relies on the **Saq Aquifer** — a vast sandstone aquifer system storing fossil water that accumulated 10,000–30,000 years ago during the last Ice Age when the Arabian Peninsula was much wetter. Extraction rates are approximately **17 times the natural recharge rate**.
- **Scale of depletion:** Between 1980 and 2012, Saudi Arabia extracted an estimated **300–400 km³ of fossil water** — close to the entire volume of Lake Erie. Water table levels have dropped by up to 100 metres in some areas.
- **Wheat production collapse:** Saudi Arabia used fossil water to grow wheat in the desert from the 1970s to 2000s, briefly making it a net wheat exporter. By 2008, recognising aquifer depletion was threatening domestic water security, the government announced it would phase out wheat growing entirely by 2016. Production dropped from 4 million tonnes (peak) to near zero — a complete policy reversal.
- **Desalination dependency:** Saudi Arabia now has 27 desalination plants producing approximately 70% of the country's drinking water — it is the world's largest producer of desalinated water. The Ras Al-Khair plant (2014) is one of the world's largest desalination facilities, producing 1.025 million m³/day. However, desalination consumes enormous amounts of energy (powered mainly by fossil

© RAG Learning · Not for redistribution

fuels), is very expensive (~$1/m³ vs $0.01 for pumped groundwater), and produces hypersaline brine that is discharged into the Gulf, damaging marine ecosystems.

- **NEOM and future challenges:** Saudi Arabia's NEOM mega-city project (including 'The Line', originally planned for 9 million people in the Tabuk desert but heavily scaled back since 2024) will require extraordinary water supplies — planned to come entirely from solar-powered desalination. Critics question whether this is environmentally or economically sustainable.

Exam Tip — Water Management Evaluation: When evaluating water management strategies, consider: (1) Is the source sustainable or finite? Fossil water and desalination solve short-term problems but create long-term dependence. (2) What are the economic costs? Desalination is 10–100x more expensive than surface water. (3) What are the environmental costs? Brine discharge, energy consumption. (4) What happens when the source runs out or becomes unaffordable? Saudi Arabia's wheat abandonment shows that even wealthy countries must eventually confront environmental limits.

13. Desertification

Desertification: The process by which fertile land in **dryland areas (arid, semi-arid, and dry sub-humid)** gradually degrades and becomes desert-like. It involves loss of topsoil, vegetation, and land productivity. Defined by the **UNCCD** (United Nations Convention to Combat Desertification). Affects approximately 1.5 billion people globally and threatens ~40% of Earth's land surface.

Location: The Sahel Region

The **Sahel** is a semi-arid belt running across Africa from **Mauritania and Senegal in the west to Eritrea and Ethiopia in the east** — approximately 5,400 km long. It sits between the Sahara Desert to the north and the tropical savanna to the south. It is the world's most at-risk region for desertification.

Causes of Desertification

Cause	Mechanism	Human or Physical

Overgrazing	Too many livestock (cattle, goats, camels) consume vegetation faster than it can regenerate; roots die; bare soil exposed to wind erosion; soil compacted by hooves	Human (population growth, cattle as status symbol)
Deforestation (fuelwood collection)	Trees and shrubs cut for firewood (80% of sub-Saharan Africa relies on wood for cooking fuel); roots no longer hold soil; wind erosion increases	Human (poverty, lack of alternatives)
Over-cultivation	Land farmed without rotation or fallow periods; soil nutrients exhausted; soil structure degrades; wind and rain erode bare soil between crops	Human (food demand, poverty)
Population growth	More people = more demand for food and fuel from same fixed land area; fragile marginal land brought into cultivation; pressure increases	Human
Climate change / drought	Reduced and increasingly erratic rainfall across the Sahel (linked to warming Indian Ocean and shifts in ITCZ); longer drought periods mean vegetation cannot recover between dry seasons	Physical (exacerbated by human climate change)

Effects of Desertification

- **Soil erosion:** Topsoil blown away by wind or washed by flash floods, exposing infertile subsoil
- **Reduced crop yields:** Degraded soil cannot support adequate harvests; food insecurity worsens
- **Famine and malnutrition:** Crop failures lead to starvation — particularly affecting children under 5
- **Migration:** People abandon degraded land and move to urban areas or other countries (climate/environmental refugees)
- **Conflict:** Competition for remaining productive land and water resources between farmers and herders has intensified conflicts (e.g. Sahel violence in Burkina Faso, Mali, Niger)
- **Economic loss:** UNCCD estimates desertification costs ~$42 billion per year in lost economic output

14. Case Study: Desertification in the Sahel

© RAG Learning · Not for redistribution

Case Study: Desertification in the Sahel

Location: A belt of semi-arid land approximately 350–500 km wide stretching ~5,400 km across Africa: Mauritania, Senegal, Mali, Burkina Faso, Niger, Chad, Sudan, South Sudan, Ethiopia, Eritrea.

Population: Well over 100 million people depend on this region for subsistence agriculture and pastoralism. Population is growing at ~3% per year — among the world's fastest rates — increasing pressure on land dramatically.

Historical famines:

- 1968–1973: Prolonged drought; ~100,000 deaths; millions displaced
- 1984–1985: Worst drought of the 20th century in Ethiopia/Sudan; hundreds of thousands of deaths (estimates up to 1 million); Live Aid response
- 2011–2012: Horn of Africa drought declared a famine; 260,000 deaths in Somalia alone
- 2021–2023: Ongoing food crisis with tens of millions of people facing acute food insecurity in the Sahel

The Great Green Wall project: A pan-African initiative launched in 2007 by the African Union — a planned 8,000 km × 15 km belt of native trees and vegetation stretching from Senegal to Djibouti, designed to combat desertification, improve food security, and sequester carbon (250 million tonnes by 2030).

- Progress by 2023: Only a small fraction of the 2030 restoration target (estimates range from about 4% to 20%) had been achieved
- Challenges: Funding gaps, political instability in the Sahel (multiple coups in Burkina Faso, Mali, Niger 2020–2023), drought, maintenance after planting, conflict making access impossible in some areas
- Successes: In Senegal and Ethiopia, restored areas show improved soil health, food production, and biodiversity

Water harvesting — Zai pits (Burkina Faso): Small pits (20–30 cm diameter, 20cm deep) dug in rows across degraded farmland. Fill with compost/manure. Rainwater collects in pits, allowing crops to be grown where soil is too hard for rain to penetrate. Revived and improved by farmer Yacouba Sawadogo from the 1980s, whose methods spread widely across Burkina Faso and Niger. A proven low-tech, low-cost solution.

© RAG Learning · Not for redistribution

Drought-resistant crops: NGOs and the CGIAR research network have developed drought-tolerant varieties of sorghum, millet, and cowpea. WEMA (Water Efficient Maize for Africa) project distributes drought-tolerant maize seeds to small farmers.

15. Case Study: Atacama Desert Mining (Chile)

Case Study: Resource Extraction in the Atacama Desert, Chile

Location: The Atacama Desert in northern Chile, primarily the Antofagasta Region. Elevation ranges from sea level to over 5,000m (altiplano). The driest non-polar desert on Earth.

Why it matters globally: The Atacama underpins the global green energy transition. Chile holds approximately **around a third of the world's known lithium reserves**, mostly in the Atacama Salt Flat (Salar de Atacama). Lithium is a critical component of lithium-ion batteries used in electric vehicles (EVs), smartphones, and grid-scale energy storage. Demand for lithium is projected to increase 40-fold by 2040 as the world transitions away from fossil fuels.

Copper: Chile also produces approximately **a quarter of the world's copper** — much of it from Atacama mines. The Escondida mine (Antofagasta Region, partly owned by BHP) is the world's largest copper mine, producing ~5% of global supply. Copper is essential for electrical wiring, electric motors, and renewable energy infrastructure.

Economic benefits:

- Mining accounts for around half of Chile's total export earnings and roughly 10–15% of GDP
- Creates hundreds of thousands of direct and indirect jobs (though many are FIFO — fly-in-fly-out — workers from distant cities rather than local indigenous communities)
- Antofagasta city has grown significantly as a mining service hub; high wages support local businesses
- Government royalties from mining fund public services nationally (education, healthcare)
- Chile's lithium wealth has led to intense international investment (US, China, EU competing for access)

Environmental challenges — water: This is the critical problem. Lithium is extracted from brine (salty water) pumped up from beneath salt flats. The process:

© RAG Learning · Not for redistribution

- Companies pump brine from underground aquifers into large evaporation ponds; sun evaporates water, leaving lithium-rich residue over 12–18 months
- This extraction significantly lowers the water table in the Salar de Atacama — a UNESCO Biosphere Reserve
- Flamingo populations (Andean, Chilean, James's flamingos) that feed on algae in Atacama wetlands have declined — a 2022 study linked falls of around 10% in some flamingo populations over a decade partly to lithium extraction
- The Atacameno/Lickanantay indigenous people (who have lived around the salar for thousands of years) argue their water rights are being violated; traditional farming and llama herding made impossible in some areas
- An often-quoted estimate is that around 2 million litres of water are evaporated for every tonne of lithium produced — in a desert that receives <1mm of rain per year

Environmental challenges — other:

- Tailings (mining waste) contain toxic heavy metals (arsenic, lead, mercury) that can leach into groundwater
- Dust from mine operations and tailings dams creates air pollution affecting nearby settlements
- Road and infrastructure construction through pristine desert destroys unique biological soil crusts and archaeological sites (the Atacama contains some of the world's oldest human mummies — Chinchorro mummies — and pre-Inca ruins)

Evaluation — the lithium paradox: Lithium mining in the Atacama presents a genuine paradox: extracting this resource damages a unique desert ecosystem and indigenous communities, yet the lithium itself is needed for the electric vehicles and batteries that could reduce global carbon emissions and slow climate change. Chile's government is attempting to balance this through a National Lithium Strategy (2023) giving the state majority control of new lithium projects via public–private partnerships (e.g. Codelco with SQM) to ensure greater state control, environmental standards, and indigenous community benefit-sharing — but tensions remain unresolved.

Key statistics to remember: Chile: around a third of world lithium reserves; Escondida: 5% of world copper production; ~2 million litres of water per tonne of lithium.

16. Managing Desertification

Strategy Overview

Strategy	Detail	Evaluation
Afforestation	Planting trees to stabilise soil, reduce wind speed, and restore water cycle through evapotranspiration	Effective long-term; takes years to establish; needs maintenance and water; wrong species choices can fail
Stone lines / bunds	Rows of stones laid along contour lines to slow water runoff and encourage infiltration; retains soil during rare rainstorms	Simple, cheap, can be built by local communities; well-evidenced in Burkina Faso to increase crop yields by 50%; see detail below
Zai pits	Traditional technique (see case study above) — pits with organic matter concentrate water and nutrients for crops in degraded soil	Proven highly effective; low-cost; community-led; Yacouba Sawadogo was awarded the "Right Livelihood Award" in 2018
Reduced stocking density	Limiting the number of livestock per hectare to sustainable levels; allows vegetation to recover between grazing periods	Difficult to enforce in nomadic cultures where livestock = wealth; requires community agreement and often land tenure reform
Drought-resistant crops	Breeding or GM varieties that require less water and mature faster (before drought intensifies)	Effective yield improvement; some opposition to GM crops; needs seed distribution networks to reach smallholders
International aid and agreements	WFP provides emergency food; UNCCD coordinates global anti-desertification policy; Great Green Wall funding	Emergency aid addresses symptoms, not causes; long-term development investment more valuable; UNCCD has limited enforcement power
FMNR (see below)	Farmer Managed Natural Regeneration — encouraging existing tree stumps and roots to regenerate rather than planting new trees	Faster, cheaper, and more ecologically appropriate than afforestation; transformational results in Niger — see below

© RAG Learning · Not for redistribution

Stone Bunds — Evidence and Detail

Stone bunds are simple but highly effective contour barriers. The technique involves:

1. Identifying the contour lines of a degraded slope (using a simple A-frame level made from two sticks and a string)
2. Collecting rocks from the surrounding area and arranging them in lines following the contour — typically stones 10–20cm in diameter, placed in rows 20–30cm high
3. Rainwater flowing down the slope is slowed and pooled behind the stone lines, allowing it to infiltrate the soil rather than running off
4. Sediment carried by runoff is also trapped behind the bunds, gradually rebuilding topsoil

Documented outcomes in Burkina Faso:

- Crop yields increased by **40–50%** in the first season after bund construction
- In Yatenga Province, stone bunds combined with zai pits have reclaimed an estimated **300,000 hectares** of degraded land since the 1980s
- Water table levels have risen by up to **3 metres** in some areas where stone bunds have been used widely — improving well water supply for domestic use
- Cost: approximately **$50–100 per hectare** using local labour and stone — extremely cost-effective compared to other interventions
- Self-maintaining: communities maintain their own bunds as they directly benefit from improved harvests

FMNR — Farmer Managed Natural Regeneration

> **FMNR (Farmer Managed Natural Regeneration):** A land restoration technique developed in Niger by Australian agronomist Tony Rinaudo (known as "The Man Who Stopped the Desert") in the 1980s. Rather than planting new trees (which are expensive and often fail to survive in harsh conditions), FMNR involves protecting and managing the natural regrowth of trees and shrubs from existing root systems that survived below-ground even when above-ground vegetation was destroyed by overgrazing, drought, or fuelwood cutting.

The FMNR method:

© RAG Learning · Not for redistribution

- Farmers select the strongest shoots emerging from dormant underground tree stumps and root systems (which can remain alive for decades)
- They prune competing weak shoots and protect the selected shoot from grazing animals
- Trees can grow to 2–3 metres within 3–4 years — far faster than planting new seedlings
- Farmers actively "manage" the trees — selectively harvesting branches for firewood and fodder while keeping the main trunk growing

Results in Niger (the "Re-greening of the Sahel"):

- An estimated **5 million hectares** of Niger's agricultural land has been restored using FMNR since the 1980s — one of the largest environmental transformations in Africa
- Approximately **200 million trees** have been regenerated on farmer fields across Niger
- Crop yields increased by **10–85%** depending on location — trees provide shade (reducing crop water stress), fix nitrogen (improving soil fertility), and reduce wind erosion
- Food security improved: surveys found that households in FMNR areas were more food secure, with extra food, fodder and income from trees
- Fuelwood supply improved dramatically — women (who collect fuelwood) reduced collection time from 2 hours to 30 minutes per day
- Cost: near-zero, as no seeds or planting required; farmers do the work themselves
- Technique has now been adopted across 24 countries in sub-Saharan Africa and beyond

Exam Tip — Management Strategy Comparison: FMNR vs Great Green Wall is an excellent comparison for evaluation questions. FMNR is: *bottom-up* (farmer-led), *low-cost* (near-zero), *fast* (results in 3–4 years), *locally appropriate* (uses species already adapted to local conditions), and *proven* (5 million hectares in Niger). The Great Green Wall is *top-down* (government/international), *expensive* (billions of dollars pledged), *slow* (only a small fraction complete after more than 15 years), and faces *political barriers* (coups, instability). This does NOT mean large-scale projects are useless — they can fund FMNR training and provide political will — but the lesson is that community-owned solutions tend to deliver better outcomes per dollar spent.

Exam Tip: For desertification questions, make sure you can distinguish between **physical** (climate) and **human** causes — many questions ask you to explain how BOTH contribute. The

strongest answers also link causes together (e.g. population growth increases overgrazing, which combined with reduced rainfall creates a vicious cycle of degradation).

Key Comparison: In the exam you may be asked to compare management strategies. Low-tech, community-led approaches (zai pits, stone bunds, FMNR) are often *more effective in practice* than large-scale international projects (Great Green Wall) because they are owned by the community, use local knowledge, are cheap, and do not depend on political stability or international funding.

© RAG Learning · Not for redistribution