

1.4 Weather and Climate

Eduqas GCSE Geography (C/4) — Component 1

1. Weather vs Climate

Weather: The short-term atmospheric conditions at a specific place and time — including temperature, rainfall, wind speed, humidity, cloud cover, and air pressure. Weather can change from hour to hour.

Climate: The *average* weather conditions of a place over a **long period of time** (typically 30 years). Climate describes the expected pattern of weather for a region rather than day-to-day variation.

UK Climate Characteristics — Temperate Maritime

The UK has a **temperate maritime climate**, shaped primarily by its proximity to the Atlantic Ocean and the warming North Atlantic Drift current.

Season	Temperature	Rainfall	Key influence
Winter	Mild (rarely below −5°C in lowlands)	Wet; frequent frontal rain and depressions	Atlantic Ocean stores summer heat and releases it slowly; North Atlantic Drift
Summer	Warm (rarely above 28°C in most areas)	Drier, though convectional rain possible	Longer days increase solar radiation; Atlantic limits extremes
Autumn/Spring	Transitional — no sharp boundaries	Variable; changeable	Interaction of air masses

Key characteristics of the UK climate:

- No extreme temperatures — cool summers and mild winters (compared to same latitude inland)
- Rainfall distributed throughout the year (no dry season)
- Western areas receive more rainfall than eastern areas (prevailing south-westerly winds)
- Highly variable, changeable day-to-day weather
- Influenced by several different **air masses** (Polar Maritime, Tropical Maritime, Polar Continental, Tropical Continental)

Exam tip: Don't confuse weather and climate. Weather = short-term, specific. Climate = long-term, average. A common exam question asks you to “distinguish between weather and climate” — always include both the time-scale difference and the idea of averages vs day-to-day conditions.


2. Factors Affecting UK Climate

Factor	Effect on UK Climate	Example
Latitude	The UK (50–61°N) receives less concentrated solar radiation than tropical regions because the Sun is lower in the sky. The same amount of solar energy is spread over a larger surface area. Higher latitudes are cooler.	Scotland is colder than southern England because it is at higher latitude (more northerly)
Altitude	Temperature decreases with height at an average rate of 6.5°C per 1,000 m (the environmental lapse rate). High ground is colder and receives more precipitation. Snow more likely on summits.	Cairngorm summit (1,245 m) is on average ~8°C colder than nearby Aviemore in the valley; over 100 days of snow cover/year
Distance from the sea (continentality)	The sea heats and cools more slowly than the land (<i>maritime effect</i>). Areas near the sea have smaller temperature ranges (mild winters, cool summers). Inland areas have a <i>continental climate</i> — hotter summers, colder winters, less rainfall.	London (inland) is hotter in summer and colder in winter than Plymouth (coastal). Eastern Europe has extremes not seen in W England.

Prevailing winds	UK's prevailing winds come from the south-west , bringing warm, moist air from the Atlantic Ocean. This gives the west coast more rainfall. Easterly winds from continental Europe bring cold, dry conditions in winter and hot, dry air in summer.	Cardiff receives c. 1,150 mm rain/year; London receives c. 590 mm/year (sheltered from south-westerly winds by Wales)
Ocean currents (North Atlantic Drift)	The North Atlantic Drift (NAD) — an extension of the Gulf Stream — flows north-eastward past the British Isles, bringing warm tropical water. This keeps the UK significantly warmer than other places at the same latitude (e.g. Newfoundland, Canada).	Stornoway, Scotland (58°N) has January average of 5°C; Labrador, Canada (same latitude) averages –15°C in January

3. Relief Rainfall

Relief (orographic) rainfall: Rainfall caused when moist air is forced to rise over a mountain barrier (orographic uplift). As the air rises, it cools (at the dry adiabatic lapse rate — 9.8°C/1,000 m until saturation; then at the saturated adiabatic lapse rate — ~6°C/1,000 m). Once cooled to the dew point, water vapour condenses into clouds and falls as rain on the *windward* side. On the *leeward* side (the rain shadow), descending air warms and dries out, producing lower rainfall.




Named example — the Pennines: The Pennine hills run north–south through northern England. Prevailing south–westerly winds bring moist Atlantic air that rises over the Pennines. Manchester (west side) receives c. 850 mm of rain/year; York (east side, in the rain shadow) receives c. 650 mm/year. The difference is even

more pronounced in the higher Lake District fells — Seathwaite in the Lake District averages over 3,000 mm/year (one of England's wettest places), while the Vale of York to the east averages c. 600 mm/year.

4. Global Atmospheric Circulation

Global atmospheric circulation: The large-scale movement of air around the Earth driven by unequal heating. The equator receives more solar radiation than the poles, creating temperature and pressure differences. This drives a system of convection cells (Hadley, Ferrel, and Polar cells) that redistribute heat energy around the globe and create the major wind belts and climate zones.



Pressure Belt / Zone	Latitude	Conditions	Climate result
ITCZ (Inter-Tropical Convergence Zone)	0° (equator)	Rising air, intense heating, low pressure	Equatorial climate: hot, very wet all year; tropical rainforest
Sub-tropical high pressure	~30°N/S	Descending air, high pressure, dry	Hot desert climates (Sahara, Arabian, Australian deserts)


Sub-polar low pressure	~60°N/S	Cold polar air meets warm tropical air; fronts; rising air	Temperate climate with frontal rainfall (UK, NW Europe)
Polar high pressure	90°N/S	Descending cold air, high pressure, very dry	Polar/tundra climate: very cold, low precipitation (mainly snow)

5. Depressions

Depression (mid-latitude cyclone): An area of low atmospheric pressure formed when cold polar air meets warm tropical air. The boundary between these air masses is called the **polar front**. Warm air rises over cold air, creating low pressure at the surface. Depressions bring unsettled, cloudy, wet, and windy weather to the UK. They move broadly west to east across the UK, driven by the jet stream.

Structure of a Depression

Feature	Description
Warm front	Leading edge of the warm air mass, shown on maps as a line with semicircles. Warm air rises gently over cold air ahead of it — produces wide cloud cover and steady, prolonged rain (or drizzle). Passes before the warm sector arrives.
Warm sector	Area of warm air between the warm and cold fronts. Relatively mild, drier, and clearer spells (though some low cloud possible).
Cold front	Trailing edge where cold air undercuts and lifts the warm air rapidly — shown on maps as a line with triangles. Produces brief but intense rainfall, heavy showers, thunderstorms, then cold, showery conditions as the front passes.
Occluded front	As the faster cold front catches up with the warm front, the warm sector is lifted off the ground, forming an occluded front. The depression is weakening at this stage.



Weather sequence as a depression passes over the UK (west to east):

1. **Before warm front:** Cirrus clouds high in sky, then thickening cloud (cirrostratus, altostratus, nimbostratus). Pressure falling. Steady, prolonged rain begins.
2. **In warm sector:** Warmer temperatures. Rain eases. Possible drizzle; low cloud. Wind veers (e.g. from S to SW).
3. **Cold front passes:** Heavy, brief rain or thunderstorms. Winds strengthen and veer to NW. Temperature drops sharply.
4. **After cold front:** Cold, bright, showery conditions. Cumulus/cumulonimbus clouds. Good visibility between showers. Pressure rising.

6. Anticyclones

Anticyclone: An area of **high atmospheric pressure** where air descends from upper levels toward the surface, then diverges outward. Descending air is warmed by compression, which suppresses cloud formation. Anticyclones bring settled, dry, calm weather. They rotate clockwise in the Northern Hemisphere. They can persist for days or weeks, causing blocking weather patterns.

Feature	Summer Anticyclone	Winter Anticyclone
Temperature	Hot, sunny (can reach 30°C+ in SE England)	Very cold (frost common; temperatures below 0°C at night)

Precipitation	Dry; drought risk with prolonged anticyclones	Dry; any moisture forms fog or frost rather than rain
Cloud cover	Clear skies; intense sunshine	Clear at night (radiation fog); possible low stratus by day
Wind	Calm, light winds	Calm; cold temperatures persist because winds don't bring Atlantic warmth
Hazards	Heatwave, drought, UV risk, hosepipe bans	Black ice, freezing fog, hypothermia risk for elderly
Air quality	Poor in urban areas — pollutants trapped (temperature inversions possible)	Smog possible; poor visibility in fog

Depression vs Anticyclone Comparison

Feature	Depression	Anticyclone
Pressure	Low (<1000 mb)	High (>1013 mb)
Air movement	Air converges and rises	Air descends and diverges
Wind speed	Strong; isobars close together	Light or calm; isobars widely spaced
Wind direction (N. Hemisphere)	Anticlockwise (cyclonic)	Clockwise (anticyclonic)
Cloud / precipitation	Heavy cloud; prolonged rain or showers	Clear skies; dry
Duration	Days (pass through quickly)	Days to weeks (blocking high)
Temperature	Variable; warm sector, then colder	Hot in summer; very cold in winter

Exam tip: A common 4-mark question asks you to describe the weather associated with either a depression or anticyclone. Use the table above and refer to temperature, precipitation, cloud cover, and wind. Always specify summer or winter for anticyclones as they produce very different conditions.

7. UK Extreme Weather Events

Case Study: Summer Heatwaves

2003 European Heatwave: France and southern UK experienced temperatures 6–7°C above average for over two weeks. UK recorded 38.5°C (Brogdale, Kent). Caused by a persistent anticyclone blocking Atlantic systems. Over 70,000 excess deaths across Europe; 2,000+ in UK. Crops failed; wildfires; power stations shut as river water temperatures rose too high for cooling.

2018 UK Heatwave: Prolonged high-pressure system from June to August. Temperatures reached 35.3°C (Faversham, Kent). UK's joint-hottest summer on record at the time. Drought conditions: rivers at record lows; hosepipe bans; moorland fires in Greater Manchester (Saddleworth Moor).

2022 UK Heatwave: On 19 July 2022, the UK recorded its highest ever temperature: **40.3°C** at Coningsby, Lincolnshire. Caused by a plume of hot air from North Africa channelled north by an anticyclone. Wildfires in London suburbs; rail lines buckled; Luton Airport tarmac melted; schools and businesses closed. Linked to climate change — Met Office stated such temperatures would have been virtually impossible without human-caused climate change.

Case Study: 2013–14 Somerset Levels Flooding

Winter 2013–14 was the UK's wettest in 248 years of records. A series of Atlantic depressions brought relentless storms. The Somerset Levels — a low-lying floodplain in SW England — were inundated from late December 2013 to March 2014.

- 65 sq km of farmland flooded; water depths up to 1 m; 17 million tonnes of water on the Levels
- Villages including Moorland and Muchelney were cut off for weeks by road
- Over 600 homes flooded and 16 farms evacuated; about 1,000 livestock moved to safety
- Somerset County Council initially estimated over £10 million of damage; a later economic impact assessment put the total cost to the county at up to £147.5 million

- Controversy over the Environment Agency's decision to reduce dredging of rivers Tone and Parrett in prior years
- Long-term response: £100 million Somerset Levels Flood Action Plan; 8 km of rivers dredged; upgraded pumping stations; raised flood banks

Exam tip: When answering extreme weather questions, always include: specific dates, specific statistics (temperatures, cost, areas affected), social impacts AND economic impacts, and — for higher marks — link to the cause (e.g. blocking anticyclone, exceptional run of Atlantic depressions).

8. Global Climate Zones

Climate Zone	Location	Temperature	Precipitation	Example
Equatorial	0–5° N/S (equator)	Hot all year: 25–28°C; no seasons	Very heavy: 2,000– 3,000 mm/yr; rain every afternoon	Amazon Basin; Congo Basin; Borneo
Tropical (savanna)	5–20° N/S	Hot all year: 20–32°C; wet season cooler	Strong wet and dry seasons: 750–1,500 mm (seasonal)	Nigeria; Kenya; northern Australia
Arid / Desert	20–30° N/S (sub- tropical)	Extreme range: 40– 50°C day; near 0°C night	Very low: <250 mm/yr; unreliable	Sahara; Arabian Desert; Atacama
Mediterranean	30–40° N/S (W coasts)	Hot dry summers (25– 35°C); mild wet winters (8–15°C)	500–800 mm; mostly in winter	Spain; Italy; California; SW Australia
Temperate (maritime)	40–65° N/S (W coasts)	Mild: 0–22°C; no extremes	Regular rainfall all year: 600–1,500 mm	UK; France; NW USA; New Zealand

Continental	40–65°N (inland)	Extreme range: –20 to +30°C; cold winters	Low–moderate: 300–600 mm; concentrated in summer	Russia; Canada; central USA
Polar / Tundra	>65° N/S	Very cold: –30 to +5°C; short cool summer	Very low: <300 mm/yr (mostly snow)	Arctic; Antarctica; northern Siberia

9. Evidence of Climate Change

Climate change: A long-term shift in global or regional climate patterns. Since the mid-20th century, the term predominantly refers to human-caused (anthropogenic) changes, particularly global warming driven by increased greenhouse gas emissions. Global average surface temperature has risen by approximately **+1.1°C since pre-industrial levels** by 2011–2020 (IPCC Sixth Assessment Report, 2021), and by the mid-2020s the figure was around 1.3°C.

Evidence	Data/Detail	Significance
Rising global temperatures	+1.1°C above pre-industrial average (1850–1900 baseline). The 2010s were the warmest decade on record at the time. 2024 was the hottest year on record globally (about 1.55°C above pre-industrial), beating 2023.	Direct measure of global warming; affects ecosystems, food production, and sea levels
Glacier retreat	Rhône Glacier (Switzerland): retreated well over 1 km since the mid-1800s. September Arctic sea ice extent has decreased by ~13% per decade since 1979. Himalayan glaciers losing mass rapidly.	Contributes to rising sea levels; threatens freshwater supply for millions
Rising sea levels	Global average sea level has risen ~20 cm since 1900; the rate averaged ~3.4 mm/year since 1993 and is now over 4 mm/year (roughly double the early-1990s rate).	Increased flood risk in low-lying coastal areas (Bangladesh, Maldives, Pacific islands, coastal UK cities)

More frequent extreme weather	Heatwaves more frequent and intense (UK 40°C, 2022); storms more powerful; droughts longer; flooding events more common in UK winters.	Economic damage, loss of life, food insecurity, displacement of people
Earlier spring	UK blossom dates (cherry, hawthorn) advancing by ~10–15 days since 1950s. Bird nesting dates earlier. Swallows arriving in UK earlier each decade.	Disrupts seasonal timing (phenological mismatch) between plants, insects, and birds
Shifting species ranges	UK butterfly species moving northward by c. 35–50 km per decade. Mediterranean species (e.g. little egret, bee-eater) colonising southern England. Bleaching of Great Barrier Reef (record ocean temperatures 2016, 2022, 2024).	Biodiversity loss; ecosystem disruption; extinction risk for cold-adapted species

10. Causes of Climate Change

Natural Causes

Milankovitch cycles: Long-term, predictable changes in Earth's orbit and orientation relative to the Sun, which alter the amount and distribution of solar energy received. Three components:

- **Eccentricity** — the shape of Earth's orbit changes from circular to more elliptical over ~100,000 years. A more elliptical orbit increases the seasonal variation in solar radiation.
- **Axial tilt (obliquity)** — the tilt of Earth's axis varies between 22.1° and 24.5° over ~41,000 years. Greater tilt = stronger seasons (hotter summers, colder winters).
- **Precession** — Earth wobbles on its axis like a spinning top, completing a cycle over ~26,000 years. This alters which hemisphere points toward the Sun during which season.

Milankovitch cycles explain the timing of ice ages but are too slow to explain rapid 20th/21st century warming.

Volcanic eruptions: Large volcanic eruptions inject sulphur dioxide (SO₂) into the stratosphere. SO₂ reacts with water vapour to form sulphate aerosols, which reflect incoming solar radiation back to


space, causing *short-term cooling*. Example: Mt Pinatubo eruption (Philippines, 1991) lowered global temperatures by $\sim 0.5^{\circ}\text{C}$ for 1–2 years. Volcanoes also emit CO_2 , but this is a tiny fraction of human emissions.

Solar output variations: Changes in the Sun's energy output (solar irradiance) affect Earth's temperature. The 11-year sunspot cycle causes minor variations. The *Maunder Minimum* (1645–1715) was a period of reduced solar activity coinciding with the Little Ice Age. However, solar output has been flat or declining slightly since 1980 while temperatures have risen — ruling out solar activity as the main cause of recent warming.

Human Causes — The Enhanced Greenhouse Effect

Greenhouse effect (natural): Solar radiation passes through the atmosphere to warm Earth's surface. Earth re-emits this energy as long-wave infrared radiation. Greenhouse gases (water vapour, CO_2 , methane, nitrous oxide) in the atmosphere absorb some of this outgoing radiation and re-emit it back toward Earth, keeping the planet $\sim 33^{\circ}\text{C}$ warmer than it would otherwise be. This is a natural and essential process.

Enhanced (anthropogenic) greenhouse effect: Human activities have increased concentrations of greenhouse gases far beyond natural levels, trapping more heat and causing additional warming. CO_2 has risen from ~ 280 ppm (pre-industrial) to over **420 ppm** in 2024.



Greenhouse Gas	Main Human Sources	Relative Impact
Carbon dioxide (CO₂)	Burning fossil fuels (coal, oil, gas) for energy and transport; cement production; deforestation (trees absorb CO ₂ when cut down this store is released and future absorption is lost)	Currently 420 ppm; responsible for ~64% of enhanced greenhouse effect
Methane (CH₄)	Livestock farming (cattle belching/manure); rice paddies; landfill waste decomposition; natural gas leaks	~80x more powerful than CO ₂ over 20 years; responsible for ~17% of warming
Nitrous oxide (N₂O)	Fertiliser use in agriculture; burning fossil fuels	~300x more powerful than CO ₂ responsible for ~6% of warming
Deforestation (indirect)	Removing forests (for farming, logging, urbanisation) releases stored carbon AND reduces future CO ₂ absorption. Deforestation and other land-use change cause roughly 10–15% of global emissions.	Brazilian Amazon deforestation rose by around 70% between 2018 and 2021, before falling sharply in 2023–24; Congo Basin and SE Asian forests also under threat

Exam tip: For higher marks on causes of climate change, clearly distinguish between **natural** causes (Milankovitch cycles, volcanic activity, solar variation) and **human** causes (fossil fuel burning, deforestation, agriculture). Examiners expect you to explain the *mechanism* — don't just list sources; explain how each cause leads to warming (or cooling). For the greenhouse effect, describe the process: solar radiation in → absorbed by surface → re-emitted as infrared → absorbed by GHGs → re-emitted back to Earth → warming.

Exam tip — extended (6–8 mark) questions: On Eduqas extended questions about climate change, structure your answer as: evidence → natural causes → human causes → relative significance (human causes dominate since 1950 — IPCC consensus). Use named statistics throughout.