


Visualizing Quadratics


Computer Vision

Carnegie Mellon University (Kris Kitani)



Equation of a circle

$$1 = x^2 + y^2$$




Equation of a 'bowl' (paraboloid)

$$f(x, y) = x^2 + y^2$$

If you slice the bowl at


$$f(x, y) = 1$$

what do you get?



Equation of a circle

$$1 = x^2 + y^2$$




Equation of a 'bowl' (paraboloid)

$$f(x, y) = x^2 + y^2$$

If you slice the bowl at

$$f(x, y) = 1$$

what do you get?



$$f(x, y) = x^2 + y^2$$


can be written in matrix form like this...

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Just two different ways to express a quadratic.

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

‘sliced at 1’



What happens if you **increase**
coefficient on ***x***?

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$


and slice at 1

What happens if you **increase**
coefficient on **x**?

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

and slice at 1

decrease width in **x**!




What happens if you **increase**
coefficient on **x**?

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

and slice at 1

decrease width in **x**!



What happens to the gradient in **x**?

increases gradient in **x**
'thins the bowl in x'

What happens if you **increase**
coefficient on **y**?


$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

and slice at 1

What happens if you **increase**
coefficient on **y**?

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

and slice at 1




What happens if you **increase**
coefficient on **y**?

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

and slice at 1

What happens to the gradient in **y**?




What happens if you **increase**
coefficient on **y**?

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

and slice at 1

What happens to the gradient in **y**?

increases gradient in **y**
'thins the bowl in y'




$$f(x, y) = x^2 + y^2$$

can be written in matrix form like this...

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

What's the shape?
What are the eigenvectors?
What are the eigenvalues?




Recall

Singular Value Decomposition

$$\underset{n \times m}{\mathbf{A}} = \underset{n \times n}{\mathbf{U}} \overset{\text{ortho-normal}}{\underset{n \times m}{\mathbf{\Sigma}}} \overset{\text{diagonal}}{\underset{m \times m}{\mathbf{V}}} \overset{\text{ortho-normal}}{\mathbf{T}}$$

unit norm constraint



orthogonal: inner (dot) product between columns/rows is zero

norm (unit vector): magnitude of each column/row is equal to 1

$$f(x, y) = x^2 + y^2$$

can be written in matrix form like this...

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Result of Singular Value Decomposition (SVD)

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}^T$$

$$f(x, y) = x^2 + y^2$$

can be written in matrix form like this...

$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Result of Singular Value Decomposition (SVD)

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \boxed{1} & \boxed{0} \\ \boxed{0} & \boxed{1} \end{bmatrix} \begin{bmatrix} \boxed{1} & 0 \\ 0 & \boxed{1} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}^T$$

eigenvectors
eigenvalues
along diagonal

axis of the
gradient of the
‘ellipse slice’
quadratic along



the axis

But what do these values correspond to geometrically?


$$\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \boxed{1} & 0 \\ \boxed{0} & 1 \end{bmatrix} \begin{bmatrix} \boxed{1} & 0 \\ 0 & 1 \end{bmatrix}^T \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Eigenvectors Eigenvalues


Eigenvectors




Recall:


$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

you can smash this bowl in the **y** direction




$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

you can smash this bowl in the **x** direction


$$f(x, y) = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} 4 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \boxed{1} & 0 \\ 0 & \boxed{4} \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}^T$$



Eigenvalues Eigenvectors Eigenvectors



$$\mathbf{A} = \begin{bmatrix} 3.25 & 1.30 \\ 1.30 & 1.75 \end{bmatrix} = \begin{bmatrix} 0.50 & -0.87 \\ -0.87 & -0.50 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} 0.50 & -0.87 \\ -0.87 & -0.50 \end{bmatrix}^T$$

Eigenvalues



Eigenvectors



$$\mathbf{A} = \begin{bmatrix} 7.75 & 3.90 \\ 3.90 & 3.25 \end{bmatrix} = \begin{bmatrix} 0.50 & -0.87 \\ -0.87 & -0.50 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 10 \end{bmatrix} \begin{bmatrix} 0.50 & -0.87 \\ -0.87 & -0.50 \end{bmatrix}^T$$

Eigenvalues

Eigenvectors



We will need this to understand...


Error function

(for Harris corners, which we'll cover next)

The surface $E(u,v)$ is locally approximated by a quadratic form

$$E(u,v) \approx [u \ v] M \begin{bmatrix} u \\ v \end{bmatrix}$$

$$M = \sum \begin{bmatrix} I_x^2 & I_x I_y \\ I_x I_y & I_y^2 \end{bmatrix}$$



Conic section of Error function

Since M is symmetric, we have $M = R^{-1} \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} R$

We can visualize M as an ellipse with axis lengths determined by the eigenvalues and orientation determined by R

Ellipse equation:

$$\begin{bmatrix} u & v \end{bmatrix} M \begin{bmatrix} u \\ v \end{bmatrix} = 1$$

'isocontour'

