

The trigonometric identity is  $\sin^2(\theta) + \cos^2(\theta) = 1$ .

# Formulas

The trigonometric identity is  $\sin^2(\theta) + \cos^2(\theta) = 1$ .

The trigonometric identity  
is `$ \sin^2(\theta) + \cos^2(\theta) = 1 $`.

# Formulas

The trigonometric identity is  $\sin^2(\theta) + \cos^2(\theta) = 1$ .

The trigonometric identity  
is `$ \sin^2(\theta) + \cos^2(\theta) = 1 $`.

```
\usepackage{amsmath,amssymb}  
\usepackage{commath,mathtools}
```

# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	$\$$ $\$$	$\sqrt[3]{8}$	$\$$ $\$$
$\frac{2}{3}$	$\$$ $\$$	$x_1$	$\$$ $\$$
$6 \geq 3$	$\$$ $\$$	$x_1^2$	$\$$ $\$$
$a^2 + b^2$	$\$$ $\$$	$a^{2+b^2}$	$\$$ $\$$

# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	$\$ \text{\textcolor{blue}{\code{\sqrt{2}}} \$$	$\sqrt[3]{8}$	$\$ \quad \$$
$\frac{2}{3}$	$\$ \quad \$$	$x_1$	$\$ \quad \$$
$6 \geq 3$	$\$ \quad \$$	$x_1^2$	$\$ \quad \$$
$a^2 + b^2$	$\$ \quad \$$	$a^{2+b^2}$	$\$ \quad \$$

# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	$\$ \backslash\text{sqrt}\{2\} \$$	$\sqrt[3]{8}$	$\$ \quad \quad \quad \$$
$\frac{2}{3}$	$\$ \backslash\text{frac}\{2\}\{3\} \$$	$x_1$	$\$ \quad \$$
$6 \geq 3$	$\$ \quad \quad \$$	$x_1^2$	$\$ \quad \$$
$a^2 + b^2$	$\$ \quad \quad \$$	$a^{2+b^2}$	$\$ \quad \quad \$$

# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	<code>\$ \sqrt{2} \$</code>	$\sqrt[3]{8}$	<code>\$ \sqrt[3]{8} \$</code>
$\frac{2}{3}$	<code>\$ \frac{2}{3} \$</code>	$x_1$	<code>\$ x_1 \$</code>
$6 \geq 3$	<code>\$ 6 \geq 3 \$</code>	$x_1^2$	<code>\$ x_1^2 \$</code>
$a^2 + b^2$	<code>\$ a^2 + b^2 \$</code>	$a^{2+b^2}$	<code>\$ a^{2+b^2} \$</code>

# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	<code>\$ \sqrt{2} \$</code>	$\sqrt[3]{8}$	<code>\$ \sqrt[3]{8} \$</code>
$\frac{2}{3}$	<code>\$ \frac{2}{3} \$</code>	$x_1$	<code>\$ x_1 \$</code>
$6 \geq 3$	<code>\$ 6 \geq 3 \$</code>	$x_1^2$	<code>\$ x_1^2 \$</code>
$a^2 + b^2$	<code>\$ a^2 + b^2 \$</code>	$a^{2+b^2}$	<code>\$ a^{2+b^2} \$</code>



# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	<code>\$ \sqrt{2} \$</code>	$\sqrt[3]{8}$	<code>\$ \sqrt[3]{8} \$</code>
$\frac{2}{3}$	<code>\$ \frac{2}{3} \$</code>	$x_1$	<code>\$ x_1 \$</code>
$6 \geq 3$	<code>\$ 6 \geq 3 \$</code>	$x_1^2$	<code>\$ x_1^2 \$</code>
$a^2 + b^2$	<code>\$ a^2 + b^2 \$</code>	$a^{2+b^2}$	<code>\$ a^{2+b^2} \$</code>

# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	<code>\$ \sqrt{2} \$</code>	$\sqrt[3]{8}$	<code>\$ \sqrt[3]{8} \$</code>
$\frac{2}{3}$	<code>\$ \frac{2}{3} \$</code>	$x_1$	<code>\$ x_1 \$</code>
$6 \geq 3$	<code>\$ 6 \geq 3 \$</code>	$x_1^2$	<code>\$ x_1^2 \$</code>
$a^2 + b^2$	<code>\$ a^2 + b^2 \$</code>	$a^{2+b^2}$	<code>\$ a^{2+b^2} \$</code>

# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	<code>\$ \sqrt{2} \$</code>	$\sqrt[3]{8}$	<code>\$ \sqrt[3]{8} \$</code>
$\frac{2}{3}$	<code>\$ \frac{2}{3} \$</code>	$x_1$	<code>\$ x_1 \$</code>
$6 \geq 3$	<code>\$ 6 \geq 3 \$</code>	$x_1^2$	<code>\$ x_1^2 \$</code>
$a^2 + b^2$	<code>\$ a^2 + b^2 \$</code>	$a^{2+b^2}$	<code>\$ a^{2+b^2} \$</code>

# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	<code>\$ \sqrt{2} \$</code>	$\sqrt[3]{8}$	<code>\$ \sqrt[3]{8} \$</code>
$\frac{2}{3}$	<code>\$ \frac{2}{3} \$</code>	$x_1$	<code>\$ x_1 \$</code>
$6 \geq 3$	<code>\$ 6 \geq 3 \$</code>	$x_1^2$	<code>\$ x_1^2 \$</code>
$a^2 + b^2$	<code>\$ a^2 + b^2 \$</code>	$a^{2+b^2}$	<code>\$ a^{2 + b^2} \$</code>

# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	<code>\$ \sqrt{2} \$</code>	$\sqrt[3]{8}$	<code>\$ \sqrt[3]{8} \$</code>
$\frac{2}{3}$	<code>\$ \frac{2}{3} \$</code>	$x_1$	<code>\$ x_1 \$</code>
$6 \geq 3$	<code>\$ 6 \geq 3 \$</code>	$x_1^2$	<code>\$ x_1^2 \$</code>
$a^2 + b^2$	<code>\$ a^2 + b^2 \$</code>	$a^{2+b^2}$	<code>\$ a^{2 + b^2} \$</code>

`$ x^{22} $`:  $x^{22}$

# Formulas: The basics

Formula	Code	Formula	Code
$\sqrt{2}$	<code>\$ \sqrt{2} \$</code>	$\sqrt[3]{8}$	<code>\$ \sqrt[3]{8} \$</code>
$\frac{2}{3}$	<code>\$ \frac{2}{3} \$</code>	$x_1$	<code>\$ x_1 \$</code>
$6 \geq 3$	<code>\$ 6 \geq 3 \$</code>	$x_1^2$	<code>\$ x_1^2 \$</code>
$a^2 + b^2$	<code>\$ a^2 + b^2 \$</code>	$a^{2+b^2}$	<code>\$ a^{2 + b^2} \$</code>

`$ x^22 $` :  $x^22$  | `$ x^{\{22\}} $` :  $x^{22}$

## Formulas: Symbols

Formula	Code	Formula	Code
$x_1, \dots, x_n$	<code>\$</code>	$5 \cdot 6$	<code>\$</code>
$\alpha, \beta, \gamma$	<code>\$</code>	$A, B, \Gamma$	<code>\$</code>
$\epsilon, \varepsilon$	<code>\$</code>	$\mathcal{P}$	<code>\$</code>
$\phi, \varphi$	<code>\$</code>	$\mathbb{P}$	<code>\$</code>

## Formulas: Symbols

Formula	Code	Formula	Code
$x_1, \dots, x_n$	<code><math>\text{\textcolor{blue}{x\_1}, \text{\textcolor{blue}{\dots}}, x\_n}</math></code>	$5 \cdot 6$	
$\alpha, \beta, \gamma$		$A, B, \Gamma$	
$\epsilon, \varepsilon$		$\mathcal{P}$	
$\phi, \varphi$		$\mathbb{P}$	



## Formulas: Symbols

Formula	Code	Formula	Code
$x_1, \dots, x_n$	<code>\$ x_1, \dots, x_n \$</code>	$5 \cdot 6$	<code>\$ \$</code>
$\alpha, \beta, \gamma$	<code>\$ \alpha, \beta, \gamma \$</code>	$A, B, \Gamma$	<code>\$ \$</code>
$\epsilon, \varepsilon$	<code>\$ \$</code>	$\mathcal{P}$	<code>\$ \$</code>
$\phi, \varphi$	<code>\$ \$</code>	$\mathbb{P}$	<code>\$ \$</code>

## Formulas: Symbols

Formula	Code	Formula	Code
$x_1, \dots, x_n$	<code>\$ x_1, \dots, x_n \$</code>	$5 \cdot 6$	<code>\$ \$</code>
$\alpha, \beta, \gamma$	<code>\$ \alpha, \beta, \gamma \$</code>	$A, B, \Gamma$	<code>\$ \$</code>
$\epsilon, \varepsilon$	<code>\$ \epsilon, \varepsilon \$</code>	$\mathcal{P}$	<code>\$ \$</code>
$\phi, \varphi$	<code>\$ \$</code>	$\mathbb{P}$	<code>\$ \$</code>

## Formulas: Symbols

Formula	Code	Formula	Code
$x_1, \dots, x_n$	<code>\$ x_1, \dots, x_n \$</code>	$5 \cdot 6$	<code>\$ \$</code>
$\alpha, \beta, \gamma$	<code>\$ \alpha, \beta, \gamma \$</code>	$A, B, \Gamma$	<code>\$ \$</code>
$\epsilon, \varepsilon$	<code>\$ \epsilon, \varepsilon \$</code>	$\mathcal{P}$	<code>\$ \$</code>
$\phi, \varphi$	<code>\$ \phi, \varphi \$</code>	$\mathbb{P}$	<code>\$ \$</code>

## Formulas: Symbols

Formula	Code	Formula	Code
$x_1, \dots, x_n$	<code>\$ x_1, \dots, x_n \$</code>	$5 \cdot 6$	<code>\$ 5\cdot 6 \$</code>
$\alpha, \beta, \gamma$	<code>\$ \alpha, \beta, \gamma \$</code>	$A, B, \Gamma$	<code>\$ A, B, \Gamma \$</code>
$\epsilon, \varepsilon$	<code>\$ \epsilon, \varepsilon \$</code>	$\mathcal{P}$	<code>\$ \mathcal{P} \$</code>
$\phi, \varphi$	<code>\$ \phi, \varphi \$</code>	$\mathbb{P}$	<code>\$ \mathbb{P} \$</code>

Formula	Code	Formula	Code
$x_1, \dots, x_n$	<code>\$ x_1, \dots, x_n \$</code>	$5 \cdot 6$	<code>\$ 5\cdot 6 \$</code>
$\alpha, \beta, \gamma$	<code>\$ \alpha, \beta, \gamma \$</code>	$A, B, \Gamma$	<code>\$ A, B, \Gamma \$</code>
$\epsilon, \varepsilon$	<code>\$ \epsilon, \varepsilon \$</code>	$\mathcal{P}$	<code>\$ \mathcal{P} \$</code>
$\phi, \varphi$	<code>\$ \phi, \varphi \$</code>	$\mathbb{P}$	<code>\$ \mathbb{P} \$</code>

# Formulas: Symbols

Formula	Code	Formula	Code
$x_1, \dots, x_n$	<code>\$ x_1, \dots, x_n \$</code>	$5 \cdot 6$	<code>\$ 5\cdot 6 \$</code>
$\alpha, \beta, \gamma$	<code>\$ \alpha, \beta, \gamma \$</code>	$A, B, \Gamma$	<code>\$ A, B, \Gamma \$</code>
$\epsilon, \varepsilon$	<code>\$ \epsilon, \varepsilon \$</code>	$\mathcal{P}$	<code>\$ \mathcal{P} \$</code>
$\phi, \varphi$	<code>\$ \phi, \varphi \$</code>	$\mathbb{P}$	<code>\$ \mathbb{P} \$</code>

## Formulas: Symbols

Formula	Code	Formula	Code
$x_1, \dots, x_n$	<code>\$ x_1, \dots, x_n \$</code>	$5 \cdot 6$	<code>\$ 5 \cdot 6 \$</code>
$\alpha, \beta, \gamma$	<code>\$ \alpha, \beta, \gamma \$</code>	$A, B, \Gamma$	<code>\$ A, B, \Gamma \$</code>
$\epsilon, \varepsilon$	<code>\$ \epsilon, \varepsilon \$</code>	$\mathcal{P}$	<code>\$ \mathcal{P} \$</code>
$\phi, \varphi$	<code>\$ \phi, \varphi \$</code>	$\mathbb{P}$	<code>\$ \mathbb{P} \$</code>

# Formulas: Vectors

Formule	Code	Formule	Code
$\vec{x}$	$\$ \backslash\mathrm{vec}\{\mathrm{x}\} \$$	$\vec{F}_{\mathrm{tot}}$	$\$ \backslash\mathrm{vec}\{\mathrm{F}\}_{\backslash\mathrm{text}\{\mathrm{tot}\}} \$$
$\mathbf{x}$	$\$ \backslash\mathrm{mathbf}\{\mathrm{x}\} \$$	$\hat{i} + 6 \hat{k}$	$\$ \backslash\mathrm{hat}\{\backslash\mathrm{imath}\} + 6\backslash,\backslash\mathrm{hat}\{\mathrm{k}\} \$$
$\ \vec{x}\ $	$\$ \backslash\mathrm{norm}\{\backslash\mathrm{vec}\{\mathrm{x}\}\} \$$	$\nabla \times \mathbf{A}$	$\$ \backslash\mathrm{nabla}\backslash\mathrm{times}\backslash\mathrm{mathbf}\{\mathrm{A}\} \$$

$$\vec{F}_{tot}, \vec{F}_{\mathrm{tot}}$$



# Formulas: Calculus

```
\usepackage{commath}
```

```
\dod{\sin(x)}{x}, \dod{f(x,y)}{x}, \partial_x f
```

```
\int_{0}^{\infty} e^{-x} \dif x = 1
```

$$\frac{d \sin(x)}{dx}, \frac{\partial f(x,y)}{\partial x}, \partial_x f$$

$$\int_0^\infty e^{-x} dx = 1$$

Formula	Code	Formula	Code
$a \leq b$	<code>\$ a \leq b \$</code>	$a \geq b$	<code>\$ a \geq b \$</code>
$a < b$	<code>\$ a &lt; b \$</code>	$a > b$	<code>\$ a &gt; b \$</code>
$a \ll b$	<code>\$ a \ll b \$</code>	$a \gg b$	<code>\$ a \gg b \$</code>
$a = b$	<code>\$ a = b \$</code>	$a \simeq b$	<code>\$ a \simeq b \$</code>
$a \neq b$	<code>\$ a \neq b \$</code>	$a \approx b$	<code>\$ a \approx b \$</code>
$a \sim b$	<code>\$ a \sim b \$</code>	$a \stackrel{*}{=} b$	<code>\$ a \stackrel{*}{=} b \$</code>

# Formulas: Arrows and operators

```
\DeclareMathOperator{\Image}{Image}
```

```
a \iff b, a\implies b, a\mapsto b  
\lim_{x\to 0}\frac{\sin(x)}{x} = 1  
\Image(f) = \mathbb{R}_{\geq 0}
```

$$a \iff b, a \implies b, a \mapsto b$$

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$

$$\text{Image}(f) = \mathbb{R}_{\geq 0}$$

So many! And there are lots more :-)

CTAN symbol list:

<http://mirrors.ctan.org/info/symbols/comprehensive/symbols-a4.pdf>

Detexify:

<http://detexify.kirelabs.org/classify.html>

# Equation

The trigonometric identity is

`$ \sin^2(\theta) + \cos^2(\theta) = 1 $`.

The trigonometric identity is

`\begin{equation}`

`\sin^2(\theta) + \cos^2(\theta) = 1.`

`\end{equation}`

The trigonometric identity is  $\sin^2(\theta) + \cos^2(\theta) = 1$ .

The trigonometric identity is

$$\sin^2(\theta) + \cos^2(\theta) = 1. \quad (1)$$

# Align

The double-angle formula can now be rewritten as

```
\begin{align}
\cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta) \\
&= 2\cos^2(\theta) - 1.
\end{align}
```

The double-angle formula can now be rewritten as

$$\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta) \tag{1}$$

$$= 2\cos^2(\theta) - 1. \tag{2}$$

# Align

The double-angle formula can now be rewritten as

```
\begin{align}
\cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta) \\
&= 2\cos^2(\theta) - 1.
\end{align}
```

The double-angle formula can now be rewritten as

$$\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta) \quad (1)$$

$$= 2\cos^2(\theta) - 1. \quad (2)$$

# Align

The double-angle formula can now be rewritten as

```
\begin{align}
\cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta)
\nonumber\\
&= 2\cos^2(\theta) - 1.
\end{align}
```

The double-angle formula can now be rewritten as

$$\begin{aligned}\cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta) \\ &= 2\cos^2(\theta) - 1.\end{aligned}\tag{1}$$



# Align

The double-angle formula can now be rewritten as

```
\begin{align*}
\cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta) \\
&= 2\cos^2(\theta) - 1.
\end{align*}
```

The double-angle formula can now be rewritten as

$$\begin{aligned}\cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta) \\ &= 2\cos^2(\theta) - 1.\end{aligned}$$

# Align

The double-angle formula can now be rewritten as

```
\begin{align*}
\cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta) \\
&= 2\cos^2(\theta) - 1. \tag{$ * $}
\end{align*}
```

The double-angle formula can now be rewritten as

$$\begin{aligned}\cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta) \\ &= 2\cos^2(\theta) - 1.\end{aligned}\tag{*}$$

# Align

We do this with the double-angle formula

```
\begin{align*}
\cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta),
\end{align*}
```

which we can rewrite as

```
\begin{align*}
&= \cos^2(\theta) - (1 - \cos^2(\theta))\\
&= 2\cos^2(\theta) - 1.
\end{align*}
```

We do this with the double-angle formula

$$\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta),$$

which we can rewrite as

$$\begin{aligned} &= \cos^2(\theta) - (1 - \cos^2(\theta)) \\ &= 2\cos^2(\theta) - 1. \end{aligned}$$

# Align

We do this with the double-angle formula

```
\begin{align*}
\cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta), \\
\intertext{which we can rewrite as}
&= \cos^2(\theta) - (1 - \cos^2(\theta)) \\
&= 2\cos^2(\theta) - 1.
\end{align*}
```

We do this with the double-angle formula

$$\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta),$$

which we can rewrite as

$$\begin{aligned} &= \cos^2(\theta) - (1 - \cos^2(\theta)) \\ &= 2\cos^2(\theta) - 1. \end{aligned}$$

## Also in use

AA	<code>\(\sqrt{2}\)</code>
BB	<code>\[\sqrt{3}\]</code>
CC	<code>\$\$\sqrt{4}\$\$</code>

AA	$\sqrt{2}$	BB
	$\sqrt{3}$	
CC	$\sqrt{4}$	

# Left-right

```
\begin{align*}
&f(\sum_{i=1}^n x_i) \\
&f\left(\sum_{i=1}^n x_i\right)
\end{align*}
```

$$f\left(\sum_{i=1}^n x_i\right)$$

$$f\left(\sum_{i=1}^n x_i\right)$$

# Delimiter point

```
\begin{align*}
  \left.\left[x^2\right]\right|_{x=0}^{\phantom{x^2}} = 4
\end{align*}
```

$$\left[x^2\right]\bigg|_{x=0}^{x=2} = 4,$$

```

\begin{align*}
R(\theta) &= \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix} \\
\abs{x} &= \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}
\end{align*}

```

$$R(\theta) = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}, \quad |x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$