

BROAD SPECTRUM DISINFECTANT, EMERGING VIRAL PATHOGEN, TIER 3 ANTIMICROBIAL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOSPITAL/HEALTH CARE

FOR USE AS A HOSPITAL DISINFECTANT VIRUCIDE*, DEODORIZER CLEANER:

1. Pre-clean surface. 2. Apply solution to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, auto scrubber, mechanical spray device, hand pump trigger spray device. For spray applications, spray 6 – 8 inches from surface. Do not breathe spray. 3. Treated surfaces must remain visibly wet for 10 minutes. 4. Wipe dry with a clean cloth or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished.

CLEANING AND DISINFECTING HARD, NON-POROUS SURFACES ON PERSONAL PROTECTIVE EQUIPMENT :

Pre-clean equipment, Apply solution to surfaces of the respirator with a sponge, brush, cloth, by mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Rub with brush, cloth, or sponge. Treated surfaces must remain visibly wet for 10 minutes. Remove excess solution from equipment prior to storage. The user must comply with all OSHA regulations for cleaning respiratory protection equipment (29 CFR § 1910.134).

BOTANICAL HVAC COIL DISINFECTANT & FUNGICIDE.

FOR USE AS A CLEANER, FUNGICIDE AND DISINFECTANT ON COILS:

For use on HVAC and/or Cooling Tower Coils to prevent fouling, mold and odor causing bacteria growth 1. Turn off system associated with coil. 2. The system must be inspected and cleaned with BotaniMax and duct work repaired to ensure that the system is in sound mechanical condition prior to cleaning the coil. 3. Apply Ready-to-Use BotaniMax to clean the coil, using brush or towel. 4. Apply Ready-to-Use BotaniMax to disinfect coil surfaces with a sponge, brush, cloth, by immersion, mechanical spray/mist device, hand pump trigger spray device. For spray applications, spray 6 – 8 inches from surface. Do not breathe spray. 5. Treated surfaces must remain visibly wet for 10 minutes. 6. Let drain and dry. 7. Use as needed to clean and disinfect coils.

ANIMAL PREMISES

DIRECTIONS FOR USE IN LIVESTOCK -PRODUCING AND/OR HOUSING FACILITIES:

For Initial Disinfection: 1. Empty all troughs, racks, and other feeding and watering appliances. 2. Remove all gross soil from walls to ceilings of livestock housing facilities and other areas occupied or traversed by livestock. 3. Thoroughly pre-clean all surfaces with BotaniMax. 4. For animal pathogen control close off room to contain BotaniMax, then thoroughly saturate with BotaniMax, using spraying or misting methods. 5. Allow all treated areas to remain visibly wet for at least 10 minutes. 6. After cleaning: Immerse equipment used for maintaining facilities in the BotaniMax and let dry. Allow to remain in contact for 10 minutes.

For Maintenance to Control Animal Pathogens: 1. For animal pathogen control close off room to contain BotaniMax, then thoroughly saturate area with BotaniMax, using spraying or misting methods. 2. Allow all treated areas to remain in contact with the solution for at least 2 minutes. 3. Recommend applying BotaniMax at least twice per week to control animal pathogens. Livestock includes, but is not limited to cattle, horses, sheep, swine, poultry, reptiles and rodents.

CONTINUE...

EPA Reg. No.
92089-2-103661
EPA Est. No.
92089-GA-1/72268-AL-1



**KILLS THE
SARS-CoV-2
THE VIRUS THAT CAUSES
COVID-19**

**CLEANER • DISINFECTANT • DEODORIZER
BACTERICIDE • VIRUCIDE* • FUNGICIDE** • FUNGISTAT**

Active Ingredient:

Citric Acid.....2.06%

Other Ingredients:.....97.94%

Total:.....100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for First Aid Statements and
Additional Precautionary Statements.

DIRECTIONS FOR USE IN HATCHERY ROOMS, INCUBATORS AND HATCHER FACILITIES:

For Initial Disinfection: 1. Empty all troughs, racks, and other feeding and watering appliances. 2. Remove all gross soil from walls to ceilings of facilities. 3. Thoroughly pre-clean all surfaces with BotaniMax. 4. For animal pathogen control close off room to contain BotaniMax, then thoroughly saturate with BotaniMax, using spraying or misting methods. 5. Allow all treated areas to remain in contact with this solution for at least 10 minutes. 6. After cleaning: Immerse equipment used for maintaining facilities in the BotaniMax and let dry. Allow to remain in contact for 10 minutes.

For Maintenance to Control Animal Pathogens: 1. For animal pathogen control close off room to contain product, then thoroughly saturate area with BotaniMax, using spraying or misting methods. 2. Allow all treated areas to remain in contact with the solution for at least 2 minutes. 3. Recommend applying BotaniMax at least twice per week to control animal pathogens.

DEODORIZING/CLEANING/DEGREASER

FOR USE AS A CLEANER AND/OR DEODORIZER AND/OR DEGREASER: Apply a use solution of 2 fl. oz. of this product per gal. of water or equivalent use dilution to hard, non-porous surfaces. Wipe up excess liquid with a paper towel and Allow to air dry. For heavy-duty use, add mix 4 fl. oz. of this product per gal. of water to clean hard, non-porous surfaces.

TO CLEAN/REMOVE SOAP SCUM: Apply a use solution of 2 fl. oz. of this product per gal. of water onto soils and wipe clean with a dry paper towel of lint-free cloth or microfiber cloth or sponge. No rinsing necessary. Repeat for visibly soiled areas. For stubborn stains or visibly soiled areas or tougher jobs, allow product to penetrate dirt and or soap scum before wiping. For best results, use regularly to prevent dirt and soap scum build-up.

GENERAL DEODORIZATION: To deodorize, apply 2 fl. oz. of this product per gal. of water or equivalent use dilution to hard, non-porous surfaces. Wipe up excess liquid with a paper towel and or Allow to air dry.

GLASS CLEANING/DEODORIZING DIRECTIONS: Use a 2 fl. oz. of this product per gal. of water use solution to clean and deodorize windows, mirrors, and glass surfaces. Use a coarse spray device. For spray applications, spray 6 – 8 inches from surface. Do not breathe spray. Rub with sponge or cloth. Change cloth, sponge or towels frequently to avoid re-deposition of soil.

GENERAL DEGREASER: To degrease, apply 2 fl. oz. of this product per gal. of water or equivalent use dilution to hard, non-porous surfaces. Wipe up excess liquid with a paper towel and or Allow to air dry.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store only in original container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Non-Refillable Container. Do not reuse or refill this container. Offer for recycling if available or place in trash.

FIRST AID

In case of emergency, call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS. CAUTION.** May cause eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Do not mix with bleach or mildew stain removers as irritating fumes may result.

DISTRIBUTED BY:
SECURE LOGIC, LLC
1221 BAYOU BLACK DR.
HOUMA, LA 70360



NET CONTENTS:
128 FL. Oz.