KAYA SKIN CLINIC & DERMATOLOGY INSTITUTE

Department of Clinical Dermatology & Polarized Dermoscopy
Unit 4, Ground Floor, Phoenix Marketcity, Viman Nagar, Pune, Maharashtra 411014 • Tel: +91 20 6689 3200

VALIDATED CLINICAL REPORT
DERM-ID: #KY-PN-88214
UHID: DERM-2026-44109

Patient Name:	**Mrs. Priya Venkatesh Iyer**	**Consultation Date:**	28/10/2026 • 04:30 PM
Age / Sex:	29 Years • Female	**Consulting Specialist:**	Dr. Meera Sanzgiri, MD, DNB (Dermatology)
Anatomical Site:	**Left Forearm & Antecubital Fossa**	**Presenting Complaint:**	Intense pruritus, erythematous dry patches × 3 weeks

CLINICAL PRESENTATION & EXAMINATION

Well-to-poorly demarcated erythematous papules and plaques localized over the flexor surface of the left distal arm and forearm. Notable xerosis, fine superficial collarette desquamation, and linear excoriation marks secondary to persistent scratching. Fitzpatrick Skin Phototype IV.



Site Photo #01
Left Forearm (Flexural) Polarized Dermoscopy Ref.

DERMOSCOPIC FINDINGS (POLARIZED 10×)

- **Vascular Architecture:** Clustered and patchy dotted vessels on a faint pinkish erythematous background, consistent with subacute epidermal inflammation.
- **Scaling & Surface:** Diffuse yellowish-white superficial micro-scales without follicular plugs or silvery hyperkeratotic micaceous scales.
- **Pigmentary Network:** Irregular post-inflammatory reticular hyperpigmentation; no atypical pigment network or atypical vascular structures observed.

CLINICAL IMPRESSION

1. **Subacute Eczematous Dermatitis / Atopic Flare (Flexural):** Characteristic clinical and dermoscopic features of active eczematous dermatitis.
2. **Excoriative Micro-trauma:** No secondary impetiginization or overt bacterial superinfection detected.

RECOMMENDED MANAGEMENT & PRESCRIPTION

- **Emollient Therapy:** Ceramide-based moisturizing lotion applied within 3 minutes of bathing twice daily.
- **Topical Anti-inflammatory:** Mometasone furoate cream 0.1% apply sparingly once daily at night for 7 days, then taper.
- **Antihistaminic:** Tab. Bilastine 20 mg once daily at bedtime for 10 days for pruritus control.
- **Review:** Follow-up after 14 days or earlier if signs of weeping/infection appear.

Dr. Meera Sanzgiri, MBBS, MD, DNB (Dermatology, Venereology & Leprosy)
Consultant Dermatologist & Dermatosurgeon
Reg. No: MMC / 2012 / 09 / 3381
Kaya Skin Clinic, Viman Nagar, Pune

VERIFIED CLINICAL RECORD
ID: DERM-PUN-7741
Pune, Maharashtra