

INVITED REVIEW

Integrated strategies for the diagnosis of venous thromboembolism

P. S. WELLS

Department of Medicine, Ottawa Hospital, Ottawa Health Research Institute, and the University of Ottawa, Ottawa, ON, Canada

To cite this article: Wells PS. Integrated strategies for the diagnosis of venous thromboembolism. *J Thromb Haemost* 2007; 5 (Suppl. 1): 41–50

Summary. Diagnosis of deep vein thrombosis (DVT) and pulmonary embolism (PE) is an important medical problem because of the high fatality rate from PE and the large number of cases not diagnosed before causing death. Over the last decade, there has been considerable research into the diagnostic process. It is widely accepted that venous ultrasound imaging is an accurate test for the diagnosis of DVT and is the imaging test of choice. For PE, computer tomographic pulmonary angiography (CTPA) is replacing ventilation perfusion lung scanning. Technology for CTPA is rapidly evolving and multi-row detector scans have quite reasonable sensitivity and specificity. Despite the accuracy of imaging tests, the post-test probability of disease is highly dependent on pretest probability. Clinical evaluation tools have developed that enable us to accurately categorize patients' risk prior to diagnostic imaging. One advantage of this characterization is an ability to exclude the diagnosis of DVT or PE if clinical probability is sufficiently low and when the D-dimer is negative. There are now a number of D-dimer assays that have well-defined specificities and sensitivities, which enable use in conjunction with clinical probability. A careful combination of clinical assessment, D-dimer and imaging enables safe PE rule out protocols without imaging, an ability to suspect false positive imaging results, and more accurate determination of true positive imaging. These integration strategies result in safer, more convenient and cost-effective care for patients.

Keywords: clinical probability, computerized tomographic pulmonary angiography, D-dimer, deep vein thrombosis, diagnosis, pulmonary embolism, venous ultrasound imaging.

Introduction

Venous thromboembolism (VTE), manifesting as deep vein thrombosis (DVT) or pulmonary embolism (PE), is one of the

most common cardiovascular disorders in industrialized countries, affecting about 5% of people in their lifetime [1]. PE is highly fatal and in 22% of cases is not diagnosed before causing death [2,3]. The signs and symptoms of both DVT and PE are largely non-specific and as a consequence many patients presenting with leg pain or swelling, or chest pain or dyspnea, are investigated but do not have DVT or PE. Mismanagement of PE has been a frequent problem [4], at least in part because of limitations of diagnostic tests. Imaging tests still have limitations but these can be better managed as the diagnostic workup for suspected PE has now evolved to an integrated approach that includes clinical pretest probability assessment and D-dimer testing in combination with imaging. In fact, we are currently observing an encouraging decrease in mortality from PE, which may reflect both more accurate diagnosis and the use of diagnostic algorithms [5–7]. In this review, I summarize the literature on clinical probability assessment, D-dimer, and the imaging tests used for VTE, with a focus on integration of these diagnostic modalities. Integrated strategies enable application of Bayes Theorem, that is, pretest odds times the likelihood ratio (especially pertinent is the negative likelihood ratio of 1 minus sensitivity divided by specificity) = post-test odds [8,9].

Diagnosis of DVT

Imaging tests for DVT

The test of choice for clinically suspected DVT is the highly specific venous ultrasound; a positive result is sufficiently predictive in most patients that treatment can be initiated. [10–13]. The exceptions are patients with a previous history of DVT and low pretest probability in which the positive predictive value is less [14].

In many centres ultrasound testing is limited to the proximal veins (distally to the region of the calf veins where they join the popliteal vein) as the sensitivity for proximal DVT has been reported as 97% but for calf DVT it appears to be considerably less (73%) [15]. In these centers, it has been suggested that a negative ultrasound should be repeated 1 week later (serial testing) to detect extending calf DVT [11]. However, in symptomatic patients, only 10% to 20% of thrombi detected are isolated to the calf, and only 20–30% of these extend, so routine serial testing is inefficient and inconvenient. Indeed,

Correspondence: Philip Wells, Suite F6-49, 1053 Carling Avenue, Ottawa Hospital Civic Campus, Ottawa, Ontario, Canada K1Y 4E9. Tel.: + 1 613 761 4127; fax: + 1 613 761 5351; e-mail: pwells@ohri.ca

Dr Wells is the recipient of a Canada Research Chair in Thromboembolic Diseases.

Received 2 February 2007, accepted 26 February 2007

studies suggest very few patients (1–2% in two recent studies) who have a negative initial ultrasound test will be confirmed to have proximal DVT upon serial testing [15,16]. As a result, serial testing is not cost-effective [17,18]. Three recent studies have suggested that imaging of the entire deep vein system, excluding DVT with a single negative result, is a safe strategy [19–21]. However, no prospective randomized trials have compared the safety of this with strategies that only evaluate the proximal venous system. It is unknown if the single whole-leg vein assessment can be widely applied with the accuracy described in these studies, it is more laborious, and it would still result in many needless tests. As I will discuss, a strategy that employs a combination of D-dimer, clinical probability and ultrasound imaging is ideal.

The clinical diagnosis of deep vein thrombosis

Although the clinical diagnosis of DVT is non-unspecific, it has now been well established that a clinical prediction rule incorporating signs, symptoms and risk factors, can be accurately applied to categorize patients as low, moderate or high probability for DVT (Table 1). Alternatively, the same rule can be used to categorize patients as 'DVT likely' or 'DVT unlikely' [22]. A recent systematic review (all studies included used the same clinical prediction rule) demonstrated that the incidence of DVT in the low, moderate and high clinical probability groups was 5.0% (95% CI, 4.0–8.0%), 17% (95% CI, 13–23%) and 53% (95% CI, 44–61%), respectively [23]. Inter-observer reliability has not been widely evaluated, but the reported studies involved many different physicians with a wide range of clinical experience, including junior residents. One study specifically demonstrated the model's reproducibility by resident physicians [24]. Determination of pretest probability allows for several potential diagnostic strategies. Used in

combination with ultrasound imaging it has been demonstrated that patients at low pretest probability can have DVT safely excluded on the basis of a single negative ultrasound test without serial testing [14]. However, it is my opinion that pretest probability determination is best used in algorithms that also incorporate D-dimer.

D-dimer for the diagnosis of DVT or PE

D-dimer, a degradation product of cross-linked fibrin, is typically elevated with acute VTE. D-dimer levels may also be increased by a variety of non-thrombotic disorders, including recent major surgery, hemorrhage, trauma, malignancy or sepsis, and D-dimer levels increase with age and through pregnancy [25–28]. D-dimer assays are sensitive but non-specific markers for VTE so positive D-dimer results are *not* useful to 'rule in' the diagnosis; rather the potential value is for a negative test result to 'rule out' the diagnosis. Although the negative predictive value of the D-dimer increases proportionately with increase in sensitivity, as with all tests the negative predictive value of the D-dimer is inversely related to the incidence of VTE in the population under study. Specificity is important because use of a very non-specific assay or the testing of very ill hospitalized patients would be of limited value because of the expected high number of positive results, many in patients with no VTE (i.e. false positives). Data suggest that most D-dimer assays lie on the same receiver operating characteristic curve [29–31], although one meta-analysis suggests three tests had significantly worse diagnostic odds ratios than the reference D-dimer used in the analysis (the VIDAS assay) [32]. I believe it is best to consider D-dimer assays as those with moderate sensitivity and moderate specificity or those with high sensitivity and poor specificity. The former consist of latex agglutination assays (qualitative or quantitative) and the whole blood assays, which are predominantly qualitative. The sensitivities and specificities for the agglutination assays have been in the range of 85–90%, and 50–80%, respectively [23,32,33]. Qualitative D-dimer assays have the advantages that they are simple, have a rapid turn round time and are inexpensive, but as they are interpreted by visual inspection, it is advisable that only trained observers perform and interpret them. Recent studies have proven that the sensitivity is highest for many of the quantitative assays but the specificity is less with these methods. Rapid, automated ELISA assays such as the Vidas DD test® (bioMérieux, Marcy l'Etoile, France), have demonstrated the highest sensitivity at an average of 95–98% but with specificity of 36–58%.

Approach to patients for the diagnosis of a first episode of deep vein thrombosis

Patients with leg symptoms compatible with DVT should initially have a determination of pretest probability of DVT using an established prediction model/rule such as the one we have validated (Table 1). It is important that a history and physical examination be performed first and only if DVT

Table 1 Simplified clinical model for assessment of DVT*

Clinical variable	Score
Active cancer (treatment ongoing or within previous 6 months or palliative)	1
Paralysis, paresis, or recent plaster immobilization of the lower extremities	1
Recently bedridden for 3 days or more, or major surgery within the previous 12 weeks requiring general or regional anesthesia	1
Localized tenderness along the distribution of the deep venous system	1
Entire leg swelling	1
Calf swelling at least 3 cm larger than that on the asymptomatic leg (measured 10 cm below the tibial tuberosity)	1
Pitting edema confined to the symptomatic leg	1
Collateral superficial veins (non-varicose)	1
Previously documented DVT	1
Alternative diagnosis at least as likely as DVT	–2

DVT, deep vein thrombosis. * ≥ 2 , probability of DVT is 'likely'. ≤ 1 , probability for DVT is 'unlikely'. Alternatively, < 1 is low probability, moderate is 1 or 2, and high is > 2 .

remains a diagnostic possibility should the model be applied. If it is obvious that the symptoms are due to an alternative problem DVT does not need to be a diagnostic possibility. If doubt remains or if DVT risk factors exist the physician may be wise to apply the model but individual judgment must be applied. After the pretest probability is determined a D-dimer test should be performed. If moderate sensitivity D-dimers are used then the pretest clinical probability should be $< 10\%$ to enable a negative D-dimer to exclude the diagnosis without the need for ultrasound. In our centre a score of ≤ 1 (unlikely DVT) in our model is sufficient to use with a qualitative D-dimer such as the IL-test or the SimpliRED™. However, most studies have employed our earlier scoring system and as such use a score of zero or less to enable exclusion of DVT with a negative moderate sensitivity D-dimer [23,34]. This strategy should be applicable to up to 40% of patients referred with suspected DVT. It should be kept in mind that ultrasound may provide information helpful to establish an alternative diagnosis but ultrasound imaging for DVT is not required to rule out DVT with low/unlikely probability and a negative D-dimer. Because of its better negative likelihood ratio of 0.08 (vs. 0.20 for moderate sensitivity assays) a negative Vidas DD test® (and other high sensitivity assays) may be used to exclude the diagnosis of venous thromboembolism when the pretest probability is less than 20%, which is a score ≤ 2 [35]. In my opinion, no D-dimer assay should be used to exclude DVT in patients who have high pretest probability.

Clinical assessment and D-dimer testing have the further advantage of enabling management at times when imaging is not available. Patients with moderate/high clinical probability may receive an injection of low-molecular-weight heparin (treatment dose) and imaging arranged the following day (Figs 1 and 2) [36,37]. Patients at low risk (by clinical models or negative sensitive D-dimer) may have diagnostic imaging delayed for a 12- to 24-h period without the need for anticoagulants, although typically we also anticoagulate these patients. When imaging is indicated, ultrasound is performed, limited to the proximal veins from the groin proceeding distally




Fig. 1. Diagnostic algorithm using clinical probability, D-dimer and ultrasound in patients with suspected DVT. DVT, deep vein thrombosis; US, ultrasound; LMWH, low-molecular-weight heparin.




Fig. 2. Diagnostic algorithm using clinical probability of DVT likely, D-dimer and ultrasound in patients with suspected DVT. DVT, deep vein thrombosis; US, ultrasound. *Use low-molecular-weight heparin if imaging is to be delayed.

to the trifurcation region. If the test is positive (lack of complete vein compressibility) the diagnosis is confirmed. If the veins are completely compressible a repeat ultrasound no more than 1 week later has generally been recommended. Five studies reported sufficient data to determine that a negative D-dimer result with a clinical probability estimate of moderate/high and normal initial ultrasound enables exclusion of DVT without a serial ultrasound test [22,37–40]. This strategy without use of pretest probability was validated in at least one study [41]. Serial testing after an initially normal ultrasound result can be confined to high probability patients with positive D-dimer results (Fig. 2). Note that false negative D-dimer results may occur in patients with prolonged symptoms of DVT or after prolonged heparin therapy (more than 24 h) [42]. Finally, it is important that the D-dimer should only be employed if the physician is convinced that DVT is a diagnostic possibility and not as a screening test. Indiscriminant use of D-dimer as a screening test will result in many unnecessary ultrasound tests.

Recurrent DVT

A randomized trial demonstrated the safety of combining clinical probability, D-dimer and ultrasound for diagnosis of recurrent DVT. [22]. The biggest concern is false positive ultrasound results. It is helpful to recognize that if the ultrasound reveals thrombosis that is echogenic, non-occlusive or discontinuous then chronic DVT should be considered. Serial testing or venography can help clarify the issue. If previous ultrasound results are available, an increase in clot diameter by 4 mm is suggestive of recurrence [43]. A recent study has suggested a negative sensitive D-dimer result can exclude recurrence without ultrasound, but Bayes theorem demonstrates the danger of this strategy as high pretest

probability patients may have over 20% probability of DVT after a negative highly sensitive D-dimer test [44].

Diagnosis of pulmonary embolism

Imaging procedures for pulmonary embolism

Pulmonary angiography is regarded as the gold standard test for the diagnosis of PE, and although the procedure is usually well tolerated, it is invasive, expensive and requires a skilled radiologist and a cooperative patient [45,46]. In addition, a negative result does not entirely exclude VTE, because in the PIOPED study 1.6% of patients with normal results developed PE over the 1-year follow-up, most in the first month [47,48].

Ventilation-perfusion (V/Q) lung scanning has been the imaging procedure of choice in patients with suspected PE. A normal scan essentially excludes the diagnosis of PE (1% VTE rate in follow-up), and a high-probability lung scan has an 85–90% predictive value for PE [47–49]. Unfortunately, most lung scans fit into a non-diagnostic category, in which the incidence of PE varies from 10% to 30% and further investigation is necessary. As a consequence the first imaging test in many centers is now computerized tomographic pulmonary angiography (CTPA). However, CTPA also has limitations, many of which are not appreciated by clinicians. CTPA is an evolving technology, with early single-slice detectors unable to visualize subsegmental arteries sufficiently [50]. Indeed, the first studies comparing single-slice detector CT to pulmonary angiography determined that CTPA had a sensitivity between 53% and 100% and a specificity between 81% and 100% [51]. However, in a more recent meta-analysis the pooled sensitivity and the specificity of CTPA were 86% and 93.7%, respectively [52]. A recent study suggests higher sensitivity with multi-detector row CTPA [53]. However, the less than perfect sensitivity and specificity mandates a need for management studies with CTPA. As discussed below, management studies have now been carried out, and the initial fears that CT would miss many PE seem to be unfounded and indeed, with current multi-detector row scanners, diagnostic sensitivity is such that false detection of PE, or detection of clinically irrelevant PE is a now a pressing issue. Magnetic resonance pulmonary angiography also holds promise as an imaging modality but will also inevitably benefit from integrated approaches [54–56].

Clinical diagnosis of pulmonary embolism

Pulmonary embolism is suspected in many patients with respiratory or chest complaints because of the non-specific nature of the presenting signs and symptoms. Despite the limitations of the individual clinical predictors [57–67], the PIOPED investigators and others have demonstrated that the clinicians' overall diagnostic impression could be useful in management [48,68–70]. Physician gestalt (empiric assessment) has worked but most of this data came from centres that also use predefined clinical decision tools, so, it is not clear if empiric assessment can be generalized. In addition, with empiric

assessment, the exact methods used by each clinician to estimate pretest probability are difficult to measure or reproduce [71], clinicians often disagree (even for broad categories) on the pretest probability of pulmonary embolism [72], the clinician's experience level appears to influence assessment [73], and probability estimates tend to follow a middle road, categorizing fewer into the more useful low- or high-probability groups. Thus, the empirical method has drawbacks, but it is the easiest method to use.

Several explicit clinical models have been described to determine pretest probability for PE using clinical findings, ECG and chest X-ray [62,74–77]. The most widely applied model has been published by our Canadian group (Table 2) [74,78]. At least 12 studies and over 10 000 patients have been evaluated, including five studies and over 5800 patients using the dichotomous scoring system of PE unlikely (score ≤ 4) or PE likely (score of > 4) [26,74,79–87]. The main limitation of this model is the need for the physician to consider an alternative diagnosis, which may be dependent on the physician's experience. However, the Kappa for inter-observer variability was reasonable, despite performing the repeat assessment up to 18 h after the first assessment [88]. At least three studies have demonstrated moderate to substantial interrater agreement and reproducibility of the Wells *et al.* model [83,88,89] but one study noted only moderate agreement [70]. The latter study reported a higher interobserver agreement for the Charlotte rule but that rule only has safe and unsafe categories. To my knowledge no other prediction rule has evaluated interrater agreement and reproducibility.

Other clinical assessment/prediction rules have also been reported. Miniati reported that a combination of clinical predictors (symptoms, ECG findings and chest X-ray findings) had a negative predictive value of 94%, and PE could be excluded in 42% of patients in their validation set [90]. Wicki *et al.* [69,75,91] devised a model in emergency room patients that in comparison to our model seems to be equally effective.

Table 2 Variables used to determine patient pretest probability for pulmonary embolism*

Clinical variable	Score
Clinical signs and symptoms of DVT (minimum of leg swelling and pain with palpation of the deep veins)	3
PE as or more likely than an alternative diagnosis	3
Heart rate greater than 100	1.5
Immobilization or surgery in the previous 4 weeks	1.5
Previous DVT/PE	1.5
Hemoptysis	1
Malignancy (on treatment, treated in the last 6 months or palliative)	1

DVT, deep vein thrombosis; PE, pulmonary embolism. * > 4 , probability of PE is 'likely'. ≤ 4 , probability for PE is 'unlikely'. Alternatively, < 2 is low probability, moderate is 2–6, and high is > 6 .

This model has subsequently been revised (called the Geneva Rule) and no longer requires a blood gas [76].

Kline *et al.*, in a series of studies, have evaluated PE diagnosis in the emergency department, where the need to rule out PE generally results in screening many patients with a low incidence of PE [92–94]. Their strategy doubled the rate of screening for PE, had a false negative rate of less than 1%, and did not increase imaging [95]. Their decision rule was: ‘a patient aged 50 years or older or any (patient with) a pulse rate greater than the systolic blood pressure, and either (i) unexplained hypoxemia (SaO₂ on pulse oximetry < 95% while breathing room air) or (ii) unilateral leg swelling or recent surgery or hemoptysis, was unsafe’ and required a V/Q scan or CTPA. If the physician felt that his/her unstructured estimate of PE was high regardless of the above, patients underwent imaging. In a further study, they suggested that if all the above factors are negative and the patient has no prior VTE and is not on hormone therapy, the patient is at low risk for PE [77]. Finally, a rule published by Hyers *et al.* [96] has demonstrated utility in two management studies [97,98].

In summary, there are several prediction rules to choose from, and not much evidence exists to prefer one above the other, but the use of these rules appears to help categorize patient pretest probability and should improve the diagnostic process as I outline below. All these rules have limitations, predominantly the fact that they have several variables and complex scoring systems. Efforts to simplify the rule are ongoing. We have published one potential simplification and it appears to work in our patient data bases but it has not yet been validated [99]. The Wells *et al.* [100] model used in the Christopher Study was re-evaluated with the intent of developing an easier model. The simplified model assigns one point to all the variables in the model and if any point is present imaging is indicated. This new model appears to work well in this data set but further validation will be necessary.

Approach to patients with suspected pulmonary embolism

As with the diagnosis of DVT the safety of a protocol for the diagnosis of PE is primarily defined by the rate of PE eventually detected in patients in whom the protocol excluded the diagnosis (i.e. the false-negative rate). As protocols are very unlikely to result in a zero post-test probability, a low threshold of about 1–2% is targeted. This threshold is comparable with the rate of PE or DVT at follow-up after a normal pulmonary angiogram, a negative CTPA [101] or a normal result on a V/Q scan [102–104]. Trying to achieve an even lower PE rate with a negative result seems unrealistic, as the rate of PE discovered in a composite population of hospitalized patients and outpatients without recognized signs or symptoms of PE, but who underwent contrast-enhanced CT of the chest, ranged from 1.5% to 3.4% [105,106]. Finally, a cut-off lower than 1% would lead to an unacceptable trade-off in increased imaging, and increased false-positive diagnosis of PE. Many studies have now been published using integrated strategies and the low

follow-up VTE event rates validate these approaches [26,74,79–89].

My recommended approach is to first perform clinical probability assessment. Subsequently, if patients are less than 80 years old, and are not ICU patients [107], then a D-dimer is performed using a D-dimer test that has been evaluated in VTE patients to have a negative likelihood ratio of ≤ 0.20 . Patients who are PE unlikely or low probability can have PE excluded with a negative D-dimer. The likelihood ratio of 0.06–0.09 with high sensitivity D-dimer test enables PE to be excluded with moderate pretest probability ($\leq 22\%$) when the D-dimer is negative (Figs 3 and 4). However, the high sensitivity D-dimers are limited by very low specificity in the elderly and hospitalized patients and are of little use in these groups. Seven studies using our model reported follow-up data on patients in whom PE was ruled out on the basis of clinical probability (low probability or PE unlikely) and negative D-dimer testing [26,74,79,81,82,85,86]. The VTE event rates in follow-up were less than 0.5%. Studies employing similar strategies but using other clinical assessment tools also reported very low rates of follow-up events [53,98]. Thus, the D-dimer assay can be the first objective test used after clinical assessment with the goal of determining which patients require diagnostic imaging.

If patients are high pretest probability or PE likely or they have a positive D-dimer then imaging (V/Q scan or CTPA) is required. If the imaging test is a V/Q scan then lower extremity venous ultrasound is also recommended when the VQ is non-diagnostic (i.e. neither normal nor high probability, Figs 3 and 5) [26,74,85,86,108]. If low pretest probability patients have a high probability VQ scan it is important to verify the diagnosis




Fig. 3. Strategy for diagnosis of PE using V/Q in patients who are PE unlikely. US, ultrasound; V/Q, ventilation perfusion lung scan; PE, pulmonary embolism; CTPA, computerized tomographic pulmonary angiography.




Fig. 4. Strategy for diagnosis of PE using CTPA in patients who are PE unlikely. US, ultrasound; V/Q, ventilation perfusion lung scan; PE, pulmonary embolism; CTPA, computerized tomographic pulmonary angiography.

with an ultrasound study, pulmonary angiogram or CTPA [62,109].

If the imaging test is CTPA we have conflicting data on the need for ultrasound when the CTPA is negative for PE. Two large prospective CTPA studies combined clinical probability and ultrasound with single row detector CTPA [84,97]. Patients with a negative CTPA, negative ultrasound, and low or moderate pretest probability had PE excluded (follow-up event rates 0.4–1.8%). Importantly, 15–18% of patients had negative CTPA but positive ultrasound studies. However, two more recent large studies in which many patients underwent multidetector row CTPA suggested very little additional yield from ultrasound, with only 0.9–1.4% of patients with a negative CTPA having a positive ultrasound result [53,79]. As further evidence, three large studies totaling over 4600 patients have now demonstrated that the combination of clinical probability, D-dimer testing and CTPA results in a strategy that safely excludes PE without ultrasound imaging [81,82,98]. With the lack of direct comparison of the two strategies it seems reasonable to perform ultrasound if clinical suspicion remains high despite a normal CTPA, if the patient has leg symptoms (indeed US could be performed first because of its high positive predictive value and thereby spare radiation exposure) [110] or if single detector row CTPA is employed (Fig. 6).

Bayes theorem predicts that using a (generous) sensitivity of 92% and a specificity of 94% for CTPA, patients who have low

(5%) pretest probability for PE will have a false positive rate in the order of 55% and 21% false positive rate for moderate (20%) pretest probability [111]. Furthermore, in a large accuracy study of CTPA (PIOPED II) it was reported that the positive predictive values for PE detected by CTPA in the lobar, segmental and subsegmental vessels were 97%, 68% and 25%, respectively [112]. To complicate the issue, the Kappa value for interobserver agreement when evaluating segmental vessels using CTPA with 3 mm collimation was only 0.47 [113] and a recent accuracy study comparing multidetector row CTPA with pulmonary angiography demonstrated a false positive CTPA rate of 30% with most false CT results incorrectly detecting PE in isolated segmental vessels or subsegmental vessels [114]. Clinical probability assessment enables a strategy to deal with potentially false positive results.

High-probability V/Q scans or positive results on CTPA should be considered diagnostic of PE if pretest probability is high or PE likely (Figs 5 and 6) but not when the pretest clinical probability is low (Figs 3 and 4). In this latter case, the results should be reviewed with the radiologist with consideration of a false-positive result. Confirmatory ultrasound or conventional pulmonary angiography should be considered.

It is possible that the use of diagnostic algorithms will increase the number of patients in whom a diagnosis of PE is considered as physicians may 'screen' for PE. Goldstein *et al.* [115] implemented a D-dimer-based screening system and found a 40% increase in the rate of V/Q scanning. However, the percentage of V/Q scans that were read as positive for PE increased and a PE diagnosis almost doubled using a D-dimer algorithm. Kline's group did not find an increase in imaging when D-dimer testing and algorithms were employed [95]. In addition, in hospitals where imaging is not available at night, algorithms may offer a rational method to decide which patients should receive temporary anticoagulation until




Fig. 5. Strategy for diagnosis of PE using V/Q in patients who are PE likely. US, ultrasound; V/Q, ventilation perfusion lung scan; DVT, deep vein thrombosis; PE, pulmonary embolism.




Fig. 6. Strategy for diagnosis of PE using single row detector CTPA in patients who are PE likely. US, ultrasound; PE, pulmonary embolism; CTPA, computerized tomographic pulmonary angiography. *If multidetector CTPA used, in most cases PE ruled out without need for ultrasound, but consider ultrasound if patient has leg symptoms or clinical suspicion is high.

imaging is available. Finally, it is clear that following algorithms improves patient care, as more diagnostic failures occurred if algorithms were not followed [5,7,74].

Conclusions

Recent advances in the management of patients with suspected PE and DVT have both improved diagnostic accuracy and made management algorithms safer and more accessible. Ongoing clinical trials are evaluating whether these diagnostic processes can be made even easier and less expensive. Patients at low risk with a negative D-dimer can avoid imaging tests and those at moderate risk with a negative high sensitivity D-dimer can have VTE excluded without the need for imaging. Diagnostic strategies should include pretest clinical probability, D-dimer assays and non-invasive imaging tests.

Disclosure of Conflict of Interests

PSW has received speaker honoraria from Biomerieux, Dade Behring and Sanofi-Aventis.

References

- Spencer FA, Emery C, Lessard D, Anderson F, Emani S, Aragam J, Becker RC, Goldberg RJ. The Worcester Venous Thromboembolism Study. A population-based study of the clinical epidemiology of venous thromboembolism. *J Gen Intern Med* 2006; **21**: 722–7.
- Heit JA, Silverstein MD, Mohr DN, Petterson TM, O'Fallon WM, Melton LJ III. Risk factors for deep vein thrombosis and pulmonary embolism: a population-based case-control study. *Arch Intern Med* 2000; **160**: 809–15.
- Heit JA, O'Fallon WM, Petterson TM, Lohse CM, Silverstein MD, Mohr DN, Melton LJ III. Relative impact of risk factors for deep vein thrombosis and pulmonary embolism: a population-based study. *Arch Intern Med* 2002; **162**: 1245–8.
- Schlager N, Henschke C, King T, Russo R, Binkert B, Rackson M, Hayt D. Diagnosis of pulmonary embolism at a large teaching hospital. *J Thorac Imaging* 1994; **9**: 180–4.
- Berghout A, Oudkerk M, Hicks SG, Teng TH, Pillay M, Buller HR. Active implementation of a consensus strategy improves diagnosis

- and management in suspected pulmonary embolism. *Q J Med* 2000; **93**: 335–40.
- Horlander KT, Mannino DM, Leeper KV. Pulmonary embolism mortality in the United States, 1979–1998. An analysis using multiple-cause mortality data. *Arch Intern Med* 2003; **163**: 1711–7.
- Roy PM, Meyer G, Vielle B, Le Gall C, Verschuren F, Carpentier F, Leveau P, Furber A. Appropriateness of diagnostic management and outcomes of suspected pulmonary embolism. *Ann Intern Med* 2006; **144**: 157–64.
- Bayes T. An essay towards solving a problem in the doctrine of chances. *Philos Trans R Soc Lon* 1763; **53**: 370–418.
- Heller I, Topilsky M, Shapira I, Isakov A. Graphic representation of sequential Bayesian analysis. *Methods Inf Med* 1999; **38**: 182–6.
- Lensing AW, Prandoni P, Brandjes DPM, Huisman PM, Vigo M, Tomacella G, Krekt J, ten Cate JW, Huisman MV, Buller HR. Detection of deep-vein thrombosis by real-time B-mode ultrasonography. *N Engl J Med* 1989; **320**: 342–5.
- Cogo A, Lensing AWA, Koopman MMW, Piovella F, Siragusa S, Wells PS, Villalta S, Buller HR, Turpie AGG, Prandoni P. Compression ultrasonography for diagnostic management of patients with clinically suspected deep vein thrombosis: prospective cohort study. *Br Med J* 1998; **316**: 17–20.
- Heijboer H, Buller HR, Lensing AWA, Turpie AGG, Colly LP, ten Cate JW. A comparison of real-time compression ultrasonography with impedance plethysmography for the diagnosis of deep-vein thrombosis in symptomatic outpatients. *N Engl J Med* 1993; **329**: 1365–9.
- Anand SS, Wells PS, Hunt D, Brill-Edwards P, Cook D, Ginsberg JS. Does this patient have deep vein thrombosis? *JAMA* 1998; **279**: 1094–9.
- Wells PS, Anderson DR, Bormanis J, Guy F, Mitchell M, Gray L, Clement C, Robinson KS, Lewandowski B. Value of assessment of pretest probability of deep-vein thrombosis in clinical management. *Lancet* 1997; **350**: 1795–8.
- Kearon C, Julian JA, Newman TE, Ginsberg JS. Noninvasive diagnosis of deep vein thrombosis. *Ann Intern Med* 1998; **128**: 663–77.
- Wells PS, Lensing AWA, Davidson BL, Prins MH, Hirsh J. Accuracy of ultrasound for the diagnosis of deep venous thrombosis in asymptomatic patients after orthopedic surgery. A meta-analysis. *Ann Intern Med* 1995; **122**: 47–53.
- Perone N, Bounameaux H, Perrier A. Comparison of four strategies for diagnosing deep vein thrombosis: a cost-effectiveness analysis. *Am J Med* 2000; **110**: 33–40.
- Hillner BE, Philbrick JT, Becker DM. Optimal management of suspected lower-extremity deep vein thrombosis. An evaluation with cost assessment of 24 management strategies. *Arch Intern Med* 1992; **152**: 165–75.
- Stevens SM, Elliot CG, Chan KJ, Egger MJ, Ahmed KM. Withholding anticoagulation after a negative result on duplex ultrasonography for suspected symptomatic deep venous thrombosis. *Ann Intern Med* 2004; **140**: 985–91.
- Schellong SM, Schwarz T, Halbritter K, Beyer J, Siegert G, Oettler W, Schmidt B, Schroeder HE. Complete compression ultrasonography of the leg veins as a single test for the diagnosis of deep vein thrombosis. *Thromb Haemost* 2003; **89**: 228–34.
- Elias A, Mallard L, Elias M, Alquier C, Guidolin F, Gauthier B, Viard A, Mahouin P, Vinel A, Boccalon H. A single complete ultrasound investigation of the venous network for the diagnostic management of patients with a clinically suspected first episode of deep venous thrombosis of the lower limbs. *Thromb Haemost* 2003; **89**: 221–7.
- Wells PS, Anderson DR, Rodger M, Forgie MA, Kearon C, Dreyer JF, Kovacs G, Mitchell M, Lewandowski B, Kovacs MJ. Evaluation of D-Dimer in the diagnosis of suspected deep-vein thrombosis. *N Engl J Med* 2003; **349**: 1227–35.
- Wells PS, Owen C, Doucette S, Fergusson D, Tran H. Does this patient have deep vein thrombosis? *JAMA* 2006; **295**: 199–207.

- 24 Penalzoza A, Laureys M, Wautrecht JC, Lheureux Ph, Motte S. Accuracy and safety of pretest probability assessment of deep vein thrombosis by physicians in training using the explicit Wells clinical model. *J Thromb Haemost* 2006; **4**: 278–81.
- 25 Bombeli T, Raddatz MP, Fehr J. Evaluation of an optimal dose of low-molecular-weight heparin for thromboprophylaxis in pregnant women at risk of thrombosis using coagulation activation markers. *Haemostasis* 2001; **31**: 90–8.
- 26 Bosson JL, Barro C, Satger B, Carpentier PH, Polack B, Pernod G. Quantitative high D-dimer value is predictive of pulmonary embolism occurrence independently of clinical score in a well-defined low risk factor population. *J Thromb Haemost* 2005; **3**: 93–9.
- 27 Bosson JL, Labaree J, Sevestre MA, Belmin J, Beyssier L, Elias A, Franco A, Le Roux P. Deep vein thrombosis in elderly patients hospitalized in subacute care facilities. A multicenter cross-sectional study of risk factors, prophylaxis and prevalence. *Arch Intern Med* 2003; **163**: 2613–8.
- 28 Righini M, Goehring C, Bounameaux H, Perrier A. Effects of age on the performance of common diagnostic tests for pulmonary embolism. *Am J Med* 2000; **109**: 357–61.
- 29 Philbrick JT, Heim SW. The D-dimer test for deep venous thrombosis: gold standards and bias in negative predictive value. *Clin Chem* 2003; **49**: 570–4.
- 30 Stein PD, Hull RD, Patel KC, Olson RE, Ghali WA, Brant RF, Biel RK, Bharadia V, Kalra NK. D-Dimer for the exclusion of acute venous thrombosis and pulmonary embolism. A systematic review. *Ann Intern Med* 2004; **140**: 589–602.
- 31 Kraaijenhagen RA, Lijmer JG, Bossuyt PMM, Prins MH, Heisterkamp SH, Buller HR. The accuracy of D-dimer in the diagnosis of venous thromboembolism: a meta-analysis. In: Kraaijenhagen RA, ed. *The Etiology, Diagnosis and Treatment of Venous Thromboembolism*. The Netherlands: University of Amsterdam, 2000:159–83.
- 32 Heim SW, Schechtman JM, Siadaty MS, Philbrick JT. D-dimer testing for deep venous thrombosis: a metaanalysis. *Clin Chem* 2004; **50**: 1136–47.
- 33 Kovacs MJ, MacKinnon KM, Anderson DR, O'Rourke K, Keeney M, Kearon C, Ginsberg JS, Wells PS. A comparison of three rapid D-dimer methods for the diagnosis of venous thromboembolism. *Br J Haematol* 2001; **115**: 140–4.
- 34 Goodacre S, Sutton AJ, Sampson FC. Meta-analysis: the value of clinical assessment in the diagnosis of deep venous thrombosis. *Ann Intern Med* 2005; **143**: 129–39.
- 35 British Society for Haematology. The diagnosis of deep vein thrombosis in symptomatic outpatients and the potential for clinical assessment and D-dimer assays to reduce the need for diagnostic imaging. *Br J Haematol* 2004; **124**: 15–25.
- 36 Bauld DL, Kovacs MJ. Dalteparin in emergency patients to prevent admission prior to investigation for venous thromboembolism. *Am J Emerg Med* 1999; **17**: 11–4.
- 37 Anderson DR, Kovacs MJ, Kovacs G, Stiell I, Mitchell M, Khoury V, Dreyer JF, Ward J, Wells PS. Combined use of clinical assessment and D-dimer to improve the management of patients presenting to the emergency department with suspected deep-vein thrombosis (the EDITED Study). *J Thromb Haemost* 2003; **1**: 645–51.
- 38 Bates SM, Kearon C, Crowther MA, Linkins L, O'Donnell M, Douketis JD, Lee AYY, Weitz JI, Johnston M, Ginsberg JS. A diagnostic strategy involving a quantitative latex D-dimer assay reliably excludes deep venous thrombosis. *Ann Intern Med* 2003; **138**: 787–94.
- 39 Schutgens REG, Ackermack P, Haas FJLM, Nieuwenhuis HK, Peltenburg HG, Pijlman AH, Pruijm M, Oltmans R, Kelder JC, Biesma DH. Combination of a normal D-dimer concentration and a non-high pretest clinical probability score is a safe strategy to exclude deep venous thrombosis. *Circulation* 2003; **107**: 593–7.
- 40 Kraaijenhagen RA, Piovello F, Bernardi E, Verlato F, Beckers EA, Koopman MM, Barone M, Camporese G, Potter Van Loon BJ, Prins MH, Prandoni P, Buller HR. Simplification of the diagnostic management of suspected deep vein thrombosis. *Arch Intern Med* 2002; **162**: 907–11.
- 41 Bernardi E, Prandoni P, Lensing AW, Agnelli G, Guazzaloca G. D-dimer testing as an adjunct to ultrasonography in patients with clinically suspected deep vein thrombosis: prospective cohort study. The Multicentre Italian D-dimer Ultrasound Study Investigators Group. *BMJ* 1998; **317**: 1037–40.
- 42 Stricker H, Marchetti O, Haeberli A, Mombelli G. Hemostatic activation under anticoagulant treatment: a comparison of unfractionated heparin vs. nadroparin in the treatment of proximal deep vein thrombosis. *Thromb Haemost* 1999; **82**: 1227–31.
- 43 Heijboer H, Jongbloets LMM, Buller HR, Lensing AWA, ten Cate JW. Clinical utility of real-time compression ultrasonography for diagnostic management of patients with recurrent venous thrombosis. *Acta Rad* 1992; **33**: 297–300.
- 44 Rathbun SW, Whitsett TL, Raskob GE. Negative D-dimer result to exclude recurrent deep venous thrombosis: a management trial. *Ann Intern Med* 2004; **141**: 839–45.
- 45 Hudson ER, Smith TP, McDermott VG, Newman GE, Suhocki PV, Payne CS, Stackhouse DJ. Pulmonary angiography performed with iopamidol: complications in 1,434 patients. *Radiology* 1996; **198**: 61–5.
- 46 Stein PD, Athanasoulis C, Alavi A, Greenspan RH, Hales CA, Saltzman HA, Vreim CE, Terrin ML, Weg JG. Complications and validity of pulmonary angiography in acute pulmonary embolism. *Circulation* 1992; **85**: 462–8.
- 47 Henry JW, Relyea B, Stein PD. Continuing risk of thromboemboli among patients with normal pulmonary angiograms. *Chest* 1995; **107**: 1375–8.
- 48 PIOPED Investigators. Value of the ventilation/perfusion scan in acute pulmonary embolism. Results of the prospective investigation of pulmonary embolism diagnosis (PIOPED). *JAMA* 1990; **263**: 2753–9.
- 49 Hull RD, Hirsh J, Carter CJ, Jay RM, Dodd PE, Ockelford PA, Coates G, Gill GJ, Turpie AGG, Doyle DJ, Buller HR, Raskob GE. Pulmonary angiography, ventilation lung scanning, and venography for clinically suspected pulmonary embolism with abnormal perfusion lung scan. *Ann Intern Med* 1983; **98**: 891–9.
- 50 Remy-Jardin M, Remy J, Artraud D, Fribourg M, Beregi JP. Spiral CT of pulmonary embolism: diagnostic approach, interpretive pitfalls and current indications. *Eur Radiol* 1998; **8**: 1376–90.
- 51 Rathbun SW, Raskob G, Whitsett TL. Sensitivity and specificity of helical computed tomography in the diagnosis of pulmonary embolism: a systematic review. *Ann Intern Med* 2000; **132**: 227–32.
- 52 Hayashino Y, Goto M, Noguchi Y, Fukui T. Ventilation-perfusion scanning and helical CT in suspected pulmonary embolism: meta-analysis of diagnostic performance. *Radiology* 2005; **234**: 740–8.
- 53 Perrier A, Roy PM, Sanchez O, Le Gal G, Meyer G, Gourdier AL, Furber A, Revel MP, Howarth N, Davido A, Bounameaux H. Multidetector-row computed tomography in suspected pulmonary embolism. *N Engl J Med* 2005; **352**: 1760–8.
- 54 Kluge A, Mueller C, Strunk J, Lange U, Bachmann G. Experience in 207 combined MRI examinations for acute pulmonary embolism and deep vein thrombosis. *AJR Am J Roentgenol* 2006; **186**: 1686–96.
- 55 Oudkerk M, van Beek EJ, Wielopolski P, van Ooijen PM, Brouwers-Kuyper EM, Bongaerts AHH, Berghout A. Comparison of contrast-enhanced magnetic resonance angiography and conventional pulmonary angiography for the diagnosis of pulmonary embolism: a prospective study. *Lancet* 2002; **359**: 1643–7.
- 56 van Beek EJ, Wild JM, Fink C, Moody AR, Kauczor HU, Oudkerk M. MRI for the diagnosis of pulmonary embolism. *J Magn Reson Imaging* 2003; **18**: 627–40.
- 57 Stein PD, Saltzman HA, Weg JG. Clinical characteristics of patients with acute pulmonary embolism. *Am J Cardiol* 1991; **68**: 1723–4.
- 58 Stein PD, Terrin ML, Hales CA, Palevsky HI, Saltzman HA, Thompson BT, Weg JG. Clinical, laboratory, roentgenographic, and

- electrocardiographic findings in patients with acute pulmonary embolism and no pre-existing cardiac or pulmonary disease. *Chest* 1991; **100**: 598–603.
- 59 Susac O Jr, Boudrow D, Kline JA. The clinical features of acute pulmonary embolism in ambulatory patients. *Acad Emerg Med* 1997; **4**: 891–7.
 - 60 Manganello D, Palla A, Donnamaria V, Giuntini C. Clinical features of pulmonary embolism. Doubts and certainties. *Chest* 1995; **107**(1 Suppl.): 25S–32S.
 - 61 Anderson FA Jr, Wheeler HB. Venous thromboembolism. Risk factors and prophylaxis. *Clin Chest Med* 1995; **16**: 235–51.
 - 62 Wells PS, Ginsberg JS, Anderson DR, Kearon C, Gent M, Turpie AGG, Bormanis J, Weitz J, Chamberlain M, Bowie D, Barnes D, Hirsh J. Use of a clinical model for safe management of patients with suspected pulmonary embolism. *Ann Intern Med* 1998; **129**: 997–1005.
 - 63 Hull RD, Raskob G, Carter CJ, Coates G, Gill GJ, Sackett DL, Hirsh J, Thompson M. Pulmonary embolism in outpatients with pleuritic chest pain. *Arch Intern Med* 1988; **148**: 838–44.
 - 64 Ferrari E, Imbert A, Chevalier T, Mihoubi A, Morand P, Baudouy M. The ECG in pulmonary embolism. Predictive value of negative T waves in precordial leads—80 case reports. *Chest* 1997; **111**: 537–43.
 - 65 McFarlane MJ, Imperiale TF. Use of the alveolar – arterial oxygen gradient in the diagnosis of pulmonary embolism. *Am J Med* 1994; **96**: 57–62.
 - 66 Rodger MA, Carrier M, Jones GN, Rasuli P, Raymond F, Djunaedi H, Wells PS. Diagnostic value of arterial blood gas measurement in suspected pulmonary embolism. *Am J Respir Crit Care Med* 2000; **162**: 2105–8.
 - 67 Rodger MA, Makropoulos D, Turek M, Quevillon J, Raymond F, Rasuli P, Wells PS. Diagnostic value of the electrocardiogram in suspected pulmonary embolism. *Am J Cardiol* 2000; **86**: 807–9.
 - 68 Perrier A, Desmarais S, Miron MJ, de Moerloose P, Lepage R, Slosman D, Didier D, Unger PF, Patenaude JV, Bounameaux H. Non-invasive diagnosis of venous thromboembolism in outpatients. *Lancet* 1999; **353**: 190–5.
 - 69 Chagnon I, Bounameaux H, Aujesky D, Roy P, Gourdiere A, Cornuz J, Perneger T, Perrier A. Comparison of two clinical prediction rules and implicit assessment among patients with suspected pulmonary embolism. *Am J Med* 2002; **113**: 269–75.
 - 70 Runyon MS, Webb WB, Jones AE, Kline JA. Comparison of the unstructured clinician estimate of pretest probability for pulmonary embolism to the Canadian score and the Charlotte rule: a prospective observational study. *Acad Emerg Med* 2005; **12**: 587–93.
 - 71 Richardson WS. Where do pretest probabilities come from? *ACP J Club* 1999; **4**: 68–9.
 - 72 Jackson RE, Rudoni RR, Pascual R. Emergency physician assessment of the pretest probability of pulmonary embolism. *Acad Emerg Med* 1999; **4**: 891–7.
 - 73 Rosen MP, Sands DZ, Morris J, Drake W, Davis RB. Does a physician's ability to accurately assess the likelihood of pulmonary embolism increase with training? *Acad Med* 2000; **75**: 1199–205.
 - 74 Wells PS, Anderson DR, Rodger MA, Stiell I, Dreyer JF, Barnes D, Forgie MA, Kovacs G, Ward J, Kovacs MJ. Excluding pulmonary embolism at the bedside without diagnostic imaging: management of patients with suspected pulmonary embolism presenting to the emergency department by using a simple clinical model and D-dimer. *Ann Intern Med* 2001; **135**: 98–107.
 - 75 Wicki J, Perneger TV, Junod AF, Bounameaux H, Perrier A. Assessing clinical probability of pulmonary embolism in the emergency ward: a simple score. *Arch Intern Med* 2001; **161**: 92–7.
 - 76 Le Gal G, Righini M, Roy PM, Sanchez O, Aujesky D, Bounameaux H, Perrier A. Prediction of pulmonary embolism in the emergency department: the revised Geneva score. *Ann Intern Med* 2006; **144**: 165–71.
 - 77 Kline JA, Mitchell AM, Kabrhel C, Richman PB, Courtney DM. Clinical criteria to prevent unnecessary diagnostic testing in emergency department patients with suspected pulmonary embolism. *J Thromb Haemost* 2004; **2**: 1247–55.
 - 78 Wells PS, Anderson DR, Rodger MA, Ginsberg JS, Kearon C, Gent M, Turpie AGG, Bormanis J, Weitz J, Chamberlain M, Bowie D, Barnes D, Hirsh J. Derivation of a simple clinical model to categorize patients probability of pulmonary embolism: increasing the models utility with the SimpliRED D-dimer. *Thromb Haemost* 2000; **83**: 416–20.
 - 79 Anderson DR, Wells PS, Kahn SR, Rodger M, Kovacs MJ, Morris T, Pleasance S. Computerized-tomographic pulmonary angiography compared with ventilation-perfusion lung scanning as initial diagnostic modality for patients with suspected pulmonary embolism: a randomized controlled trial. *Blood* 2005; **106**: 463 (Abstract).
 - 80 Sohne M, Kamphuisen PW, van Mierlo PJ, Buller HR. Diagnostic strategy using a modified clinical decision rule and D-dimer test to rule out pulmonary embolism in elderly in- and outpatients. *Thromb Haemost* 2005; **94**: 206–10.
 - 81 van Belle A, Buller HR, Huisman MV, Huisman PM, Kaasjager K, Kamphuisen PW, Kramer MH, Kruip MJ, Kwakkel-van Erp JM, Leebeek FW, Nijkeuter M, Prins MH, Sohne M, Tick LW. Effectiveness of managing suspected pulmonary embolism using an algorithm combining clinical probability, D-dimer testing, and computed tomography. *JAMA* 2006; **295**: 172–9.
 - 82 Goekoop RJ, Steeghs N, Niessen RW, Jonkers GJ, Dik H, Castel A, Werker-van Gelder L, Tom VL, van Klink RC, Planken EV, Huisman MV. Simple and safe exclusion of pulmonary embolism in outpatients using quantitative D-dimer and Wells' simplified decision rule. *Thromb Haemost* 2007; **97**: 146–50.
 - 83 Penalzoza A, Melot C, Dochy E, Blocklet D, Gevenois PA, Wautrecht JC, Lheureux P, Motte S. Assessment of pretest probability of pulmonary embolism in the emergency department by physicians in training using the Wells model. *Thromb Res* 2007 (in press).
 - 84 Anderson DR, Kovacs MJ, Dennie C, Kovacs G, Stiell I, Dreyer J, McCarron B, Pleasance S, Burton E, Cartier Y, Wells PS. Use of spiral computed tomography contrast angiography and ultrasonography to exclude the diagnosis of pulmonary embolism in the emergency department. *J Emerg Med* 2005; **29**: 399–404.
 - 85 Kearon C, Ginsberg JS, Douketis J, Turpie AG, Bates SM, Lee AY, Crowther MA, Weitz JI, Brill-Edwards P, Wells PS, Anderson DR, Kovacs MJ, Linkins LA, Julian JA, Bonilla LR, Gent M. An evaluation of D-dimer in the diagnosis of pulmonary embolism: a randomized trial. *Ann Intern Med* 2006; **144**: 812–21.
 - 86 Leclercq MGL, Lutsan JG, van Marwijk Kooy M, Kuipers BF, Oostdijk AHJ, van der Leur JJCM, Buller HR. Ruling out clinically suspected pulmonary embolism by assessment of clinical probability and D-dimer levels: a management study. *Thromb Haemost* 2003; **89**: 97–103.
 - 87 Kruip MJ, Slob MJ, Schijen JH, van der Heul C, Buller HR. Use of a clinical decision rule in combination with D-dimer concentration in diagnostic workup of patients with suspected pulmonary embolism: a prospective management study. *Arch Intern Med* 2002; **162**: 1631–5.
 - 88 Rodger MA, Maser E, Stiell I, Howley HE, Wells PS. The interobserver reliability of pretest probability assessment in patients with suspected pulmonary embolism. *Thromb Res* 2005; **116**: 101–7.
 - 89 Wolf SJ, McCubbin TR, Feldhaus KM, Faragher JP, Adcock DM. Prospective validation of Wells criteria in the evaluation of patients with suspected pulmonary embolism. *Ann Emerg Med* 2004; **44**: 503–10.
 - 90 Miniati M, Prediletto R, Formichi B, Marini C, Di Riccio G, Tonelli L, Allescia G, Pistolesi M. Accuracy of clinical assessment in the diagnosis of pulmonary embolism. *Am J Respir Crit Care Med* 1999; **159**: 864–71.
 - 91 Wicki J, Perrier A, Perneger TV, Bounameaux H, Junod AF. Predicting adverse outcome in patients with acute pulmonary embolism: a risk score. *Thromb Haemost* 2000; **84**: 548–52.

- 92 Kline JA, Nelson RD, Jackson RE, Courtney DM. Criteria for the safe use of D-dimer testing in emergency department patients with suspected pulmonary embolism: a multicenter US study. *Ann Emerg Med* 2002; **39**: 144–52.
- 93 Kline JA, Mitchell AM, Kabrheh C, Richman PB, Courtney DM. More on: clinical criteria to prevent unnecessary diagnostic testing in emergency department patients with suspected pulmonary embolism. *J Thromb Haemost* 2005; **3**: 190–1.
- 94 Kline JA, Runyon MS, Webb WB, Jones AE, Mitchell AM. Prospective study of the diagnostic accuracy of the Simplify D-dimer assay for pulmonary embolism in emergency department patients. *Chest* 2006; **129**: 1417–23.
- 95 Kline JA, Webb WB, Jones AE, Hernandez-Nino J. Impact of a rapid rule-out protocol for pulmonary embolism on the rate of screening, missed cases, and pulmonary vascular imaging in an urban US emergency department. *Ann Emerg Med* 2004; **44**: 490–502.
- 96 Hyers TM. Venous thromboembolism. *Am J Respir Crit Care Med* 1999; **159**: 1–14.
- 97 Musset D, Parent F, Meyer G, Maitre S, Girard P, Leroyer C, Revel MP, Carette MF, Laurent M, Charbonnier B, Laurent F, Mal H, Nonent M, Lancar R, Grenier P, Simonneau G. Diagnostic strategy for patients with suspected pulmonary embolism: a prospective multicentre outcome study. *Lancet* 2002; **360**: 1914–20.
- 98 Ghanima W, Almaas V, Aballi S, Dorje C, Nielssen BE, Holmen LO, Almaas R, Abdelnoor M, Sandset PM. Management of suspected pulmonary embolism (PE) by D-dimer and multi-slice computed tomography in outpatients: an outcome study. *J Thromb Haemost* 2005; **3**: 1926–32.
- 99 Rodger MA, Raymond F, Stiell I, Nair R, Bredeson CN, Jones G, Rasuli P, Wells PS. Development and validation of a simple clinical prediction rule to exclude Pulmonary Embolism. *Blood* 1999; **94**(10 Suppl. 1): 27 (Abstract).
- 100 Gibson NS, Sohne M, Gerdes VE, Bossuyt PM, Wells PS, Buller HR. Further validation and simplification of the Wells clinical decision rule in pulmonary embolism. *Thromb Haemost* 2007; Submitted.
- 101 Tillie-Leblond I, Mastora I, Radenne F, Paillard S, Tonnel AB, Remy J, Remy-Jardin M. Risk of pulmonary embolism after a negative spiral CT angiogram in patients with pulmonary disease: 1-year clinical follow-up study. *Radiology* 2002; **223**: 461–7.
- 102 Hull RD, Raskob GE, Coates G, Panju AA. Clinical validity of a normal perfusion lung scan on patients with suspected pulmonary embolism. *Chest* 1990; **97**: 23–6.
- 103 van Beek EJ, Kuyper PMM, Schenk BE, Brandjes DPM, ten Cate JW, Buller HR. A normal perfusion lung scan in patients with clinically suspected pulmonary embolism: frequency and clinical validity. *Chest* 1995; **108**: 170–3.
- 104 Kipper MS, Moser KM, Kortman KE, Ashburn WL. Longterm follow-up of patients with suspected pulmonary embolism and a normal lung scan. Perfusion scans in embolic suspects. *Chest* 1982; **82**: 411–5.
- 105 Gosselin MV, Rubin GD, Leung AN, Huang J, Rizk NW. Unsuspected pulmonary embolism: prospective detection on routine helical CT scans. *Radiology* 1998; **208**: 209–15.
- 106 Storto ML, Di Credico A, Guido F, Larici AR, Bonomo L. Incidental detection of pulmonary emboli on routine MDCT of the chest. *AJR Am J Roentgenol* 2005; **184**: 264–7.
- 107 Crowther MA, Cook DJ, Griffith LE, Meade M, Hanna S, Rabbat C, Bates SM, Geerts W, Johnston M, Guyatt G. Neither baseline tests of molecular hypercoagulability nor D-dimer levels predict deep venous thrombosis in critically ill medical-surgical patients. *Intensive Care Med* 2005; **31**: 48–55.
- 108 Perrier A, Bounameaux H, Morabia A, de Moerloose P, Slosman D, Didier D, Unger PF, Junod A. Diagnosis of pulmonary embolism by a decision analysis-based strategy including clinical probability, D-dimer levels, and ultrasonography: a management study. *Arch Intern Med* 1996; **156**: 531–6.
- 109 Mayo JR, Remy-Jardin M, Muller NL, Remy J, Worsley DF, Houssein-Foucher C, Kwong JS, Brown MJ. Pulmonary embolism: prospective comparison of spiral CT with ventilation-perfusion scintigraphy. *Radiology* 1997; **205**: 447–52.
- 110 Le Gal G, Righini M, Sanchez O, Roy PM, Baba-Ahmed M, Perrier A, Bounameaux H. A positive compression ultrasonography of the lower limb veins is highly predictive of pulmonary embolism on computed tomography in suspected patients. *Thromb Haemost* 2006; **95**: 963–6.
- 111 Ranji SR, Shojania KG, Trowbridge RL, Auerbach AD. Impact of reliance on CT Pulmonary Angiography on diagnosis of pulmonary embolism: a Bayesian analysis. *J Hosp Med* 2006; **1**: 81–7.
- 112 Stein PD, Fowler SE, Goodman LR, Gottschalk A, Hales CA, Hull RD, Leeper KV, Popovich J Jr, Quinn DA, Sos TA, Sostman HD, Tapson VF, Wakefield TW, Weg JG, Woodard PK, for the PIOPE II Investigators. Multidetector computed tomography for acute pulmonary embolism. *N Engl J Med* 2006; **354**: 2317–27.
- 113 Chartrand-Lefebvre C, Howarth N, Lucidarme O, Beigelman C, Cluzel P, Mourey-Gerosa I, Cadi M, Grenier P. Contrast-enhanced helical CT for pulmonary embolism detection: inter and intraobserver agreement among radiologists with variable experience. *AJR Am J Roentgenol* 1999; **172**: 107–12.
- 114 Winer-Muram HT, Rydberg J, Johnson MS, Tarver RD, Williams MD, Shah H, Namyslowski J, Conces D, Jennings SG, Ying J, Trerotola SO, Kopecky KK. Suspected acute pulmonary embolism: evaluation with multi-detector row CT vs. digital subtraction pulmonary arteriography. *Radiology* 2004; **233**: 806–15.
- 115 Goldstein NM, Kollef MH, Ward S, Gage BF. The impact of the introduction of a rapid D-dimer assay on the diagnostic evaluation of suspected Pulmonary Embolism. *Arch Intern Med* 2001; **161**: 567–71.