1. Use Euler method to solve $\frac{dy}{dx} = 7x^3y, y(0) = 2, \Delta x = 0.1$ at $x = 0.5$ in 4 decimal points.

2. Use Euler method to solve $\frac{dy}{dx} = 2x^4 + y, y(0) = 1, \Delta x = 0.2$ at $x = 1$ in 4 decimal points.

3. Use Euler method to solve $\frac{dy}{dx} = 2y + e^{8x}, y(0.2) = 4, \Delta x = 0.05$ at $x = 0.8$ in 4 decimal points.

4. Use Euler method to $\frac{dy}{dx} - 3y - \sin(x) = 0, y(0.4) = 0.75, \Delta x = 0.025$ at $x = 1$ in 4 decimal points.

5. Use Euler method to solve $\frac{dy}{dx} - \frac{5y}{x^2+9} = 0, y(0.01) = 2.36, \Delta x = 0.1$ at $x = 0.41$in 4 decimal points.

6. Use Euler method to solve $\frac{dy}{dx} - 4.2\sin(x+1) - \frac{1}{e^{x+1}} = 0, y(-0.2) = 4.11, \Delta x = 0.1$ at $x = 0.7$ in 4 decimal points.

7. Use Euler method to solve $2\frac{dy}{dx} - 3y + \frac{1}{\cosh(3x+7)} = 0, y(1) = 6.731, \Delta x = 0.1$ at $x = 2.9$ in 6 decimal points.

8. Use Euler method to $\frac{1}{4}\frac{dy}{dx} - \frac{5}{ye^{5(x-2)}} = 0, y(1) = 0.7118, \Delta x = 0.01$ at $x = 2$ in 6 decimal points.

9. Use Euler method to solve $e^{x+y}\frac{dy}{dx} = 7x^3y, y(0) = 2, \Delta x = 0.01$ at $x = 0.5$ in 6 decimal points.

10. Use Euler method to solve $\cosh^{-1}(x)\frac{dy}{dx} - 7x^3y = 0, y\left(\frac{\pi}{2}\right) = 2, \Delta x = \frac{\pi}{20}$ at $x = \pi$ in 6 decimal points.

Assoc. Prof. Dr. Noor Fadiya Mohd Noor