

Fungicide

JAGER®

*A fungicide for the control of
foliar and ear disease in winter
and spring wheat, durum wheat,
rye, triticale and winter and
spring barley.*

3 L e


GROUP 11 | 3 FUNGICIDES
MAPP 18942
F


**A suspension concentrate formulation
containing 175 g/L prothioconazole and
150 g/L trifloxystrobin**

The (COSHH) Control of Substances
Hazardous to Health Regulations may
apply to the use of this product at work.

Bayer CropScience Ltd
PO Box 1582
Cambridge
CB1 0FE

Telephone: 01223 226500

**For 24 hour emergency information
contact Bayer CropScience Ltd
Telephone: 0330 678 3382 (24 hr)**

**National Poisons Information Centre
UK: 0344 892 0111**
(medical professionals only)

**National Poisons Information Centre
Dublin: +353 1 809 2166**

JAGER®

UFI: 2AJ3-J0Q5-600X-UJV9

Contains 175 g/L prothioconazole
and 150 g/L trifloxystrobin.



Warning

May cause an allergic skin reaction.

May cause harm to breast-fed children.

Very toxic to aquatic life with long lasting effects.

Wear protective gloves/protective clothing/eye protection/face protection.
Avoid contact during pregnancy/ while nursing.

IF exposed or concerned: Call a
POISON CENTER/ doctor/ physician.
Collect spillage.

Protect from sunlight.

Dispose of contents/container to a
licensed hazardous-waste disposal
contractor or collection site except for
empty clean containers which can be
disposed of as non-hazardous waste.

**To avoid risks to human health and
the environment, comply with the
instructions for use.**

IMPORTANT INFORMATION FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crops:	Winter and spring wheat, durum wheat, rye, triticale and winter and spring barley.
Maximum individual dose:	Wheat, durum wheat, rye and triticale: 1.0 litres of product per hectare per crop. Barley: 0.75 litres of product per hectare per crop.
Maximum total dose:	Wheat, durum wheat, rye and triticale: 2.0 litres of product per hectare per crop. Barley: 1.5 litres of product per hectare per crop.
Latest time of application:	Wheat, durum wheat, rye and triticale: Before grain milky ripe stage (GS 71) Barley: up to beginning of anthesis (GS 61)
Other specific restrictions:	To reduce the risk of resistance developing in target diseases the total number of applications of products containing Qol fungicides made to any cereal crop must not exceed two.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate or when handling contaminated surfaces.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when applying the product.

However, engineering controls may replace personal protective equipment

if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT DRINK OR SMOKE.

WASH ANY CONTAMINATION from eyes immediately.

IF YOU FEEL UNWELL, seek medical advice (show label where possible).

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

Environmental Protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water.



This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation

from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone

must be maintained. The results of the LERAP must be recorded and kept available for three years.

Storage and Disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an intergrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling, and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Jager is a mixture of a triazole and strobilurin fungicide recommended for control of a wide range of diseases on winter and spring wheat, durum wheat, rye, triticale and winter and spring barley.

RESTRICTIONS

PROTECT FROM FROST.

DISEASES CONTROLLED

Wheat, Durum Wheat, triticale

Eyespot, *Septoria* (leaf and glume blotch), powdery mildew, yellow rust, brown rust, ear disease complex* – *Fusarium* ear blight.

* good reduction

Barley

Eyespot*, powdery mildew, yellow rust, brown rust, *Rhynchosporium* and net blotch.

* reduces the incidence and severity

Rye

Eyespot, powdery mildew, brown rust and *Rhynchosporium*.

Eyespot (*Oculimacula* spp.)

Jager controls eyespot in wheat, triticale and rye and reduces the incidence and severity in barley. Spray in the spring at the first sign of disease, from when the leaf sheaths begin to become erect until the 2nd node is detectable (GS 30-32).

Septoria Leaf Blotch and Glume Blotch (*Septoria tritici* and *Stagonospora nodorum*)

Apply before disease is established in the crop. To protect the upper leaves and ear apply Jager at full flag leaf emergence (GS 39) up to mid-flowering (GS 65). Where disease pressure remains high application may be repeated.

Applications to upper leaves where *S. tritici* symptoms are present are likely to be less effective.

Jager contains a DMI fungicide.

Resistance to some DMI fungicides has been identified in *Septoria* leaf blotch (*Mycosphaerella graminicola*) which may seriously affect the performance of some products. For further advice on resistance management in DMI's contact your agronomist or specialist advisor, and visit the FRAG-UK website.

Powdery Mildew (*Blumeria graminis*)

Apply Jager at the first signs of disease. Jager controls mildew in barley and when used for control of other diseases will also give control of powdery mildew in wheat, durum wheat, triticale and rye. Where disease pressure remains high application may be repeated.

Strains of wheat and barley powdery mildew resistant to Qol fungicides are common in the UK. Where specific control of wheat mildew is required, this should be achieved through a programme of measures, including products with recommendations for control of mildew that contain different active substances used in mixture or sequence.

Yellow Rust (*Puccinia striiformis*)

Apply Jager at the first signs of disease. A second application may be made 2-3 weeks later if re-infection occurs. Applications made to established infections are likely to be less effective.

Brown Rust (*Puccinia hordei*, *P. recondita*)

Apply Jager at the first signs of disease. A second application may be made 2-3 weeks later if re-infection occurs. Applications made to established infections are likely to be less effective.

Ear Disease Complex

Apply Jager soon after ear emergence until the end of flowering (GS 59-69) for good reduction of *Fusarium* ear blight and reduction of sooty moulds in wheat. Control of ear diseases can result in cleaner, brighter ears.

Leaf Blotch (*Rhynchosporium secalis*)

Jager gives high levels of *Rhynchosporium* control. Apply Jager in spring at the first signs of disease. For

severe infections a second application may be necessary 2-3 weeks later.

Net Blotch (*Pyrenophora teres*)

Apply Jager at the first signs of disease in spring/early summer. For severe infections, a second application 2-3 weeks later will give most effective control when conditions remain favourable for disease development.

CROP SPECIFIC INFORMATION

Jager may be used on all commercial varieties of winter and spring wheat, durum wheat, rye, triticale and winter and spring barley.

Apply Jager at 1.0 litres per hectare in wheat, durum wheat, triticale and rye and 0.75 litres per hectare in barley.

The maximum total dose per crop is:- 2.0 litres per hectare per crop in wheat, durum wheat, triticale and rye and 1.5 litres per hectare per crop in barley.

Apply Jager in 100-300 litres of water per hectare. The higher spray volumes are recommended where the crop is dense or disease pressure / risk is high to ensure good penetration to the lower leaves and stem bases. Disease control maybe compromised by reducing water volumes, where good spray coverage is difficult to achieve.

A spray pressure of 2-3 bar is recommended.

Apply as a **MEDIUM** spray quality (as defined by BCPC).

Latest Permitted Timing

Jager may be applied at any stage before grain milky ripe stage (GS 71) in wheat, durum wheat, triticale and rye and up to beginning of anthesis (GS 61) in barley.

MIXING AND SPRAYING

Thoroughly shake the pack before use.

Add the required quantity of Jager to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

Sprayers should be thoroughly cleaned before use, and filters and jets checked

for damage and blockages.

Boom height should be adjusted to ensure even coverage of the crop, particularly at later growth stages.

The correct height is one at which the spray from alternate nozzles meets just above the crop, in dense crops, at later growth stages, higher water volumes should be used.

Spray equipment should be thoroughly cleaned with water and detergent after use.

RESISTANCE

Jager contains trifloxystrobin, a member of the QoI cross resistance group. Jager should be used preventatively or as early as possible

in the disease cycle and should not be relied upon for its curative potential. It should not be used where disease is already established. Use Jager as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action. You must not apply more than two foliar applications of Qol containing products to any cereal crop. Strains of wheat and barley powdery mildew resistant to Qol's are common in the UK. Limiting the number of sprays is an important factor in delaying the build-up of resistant pathogen populations. There is a significant risk of widespread resistance occurring in Septoria

tritici populations in the UK. Failure to follow resistance management action may result in reduced levels of disease control. Appropriate disease risk assessment or decision support systems must be used when deciding whether to treat with Jager.

CAUTION

The possible development of resistance to Jager cannot be predicted. Should it occur Jager may not give satisfactory control and no responsibility can be accepted for any loss incurred.

® Jager and DFF are registered Trade Marks of Bayer

© Bayer CropScience Limited 2025



To access the **Safety Data Sheet** for this product scan the code or use the link below:

<https://cropscience.bayer.co.uk/our-products/fungicides/jager>
or alternatively contact your supplier