

Fungicide

iblon
TECHNOLOGY

Inatreq™ active

JESSICO[®] fusion

A fungicide for the control of foliar diseases in wheat, rye, triticale and spelt.

An emulsifiable concentrate formulation containing 50 g/L Fenpicoxamid and 37.5 g/L Isoflucypram.

This product must be sold and used only in England, Scotland and Wales.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Bayer CropScience Ltd, PO Box 1582, Cambridge, CB1 0FE
Telephone: 01223 226500

**For 24 hour emergency
information contact
Bayer CropScience Ltd**

**Telephone:
0330 678 3382 (24 hr)**

**National Poisons Information
Centre UK: 0344 892 0111
(medical professionals only)**

5 L e



GROUP 21 | 7 FUNGICIDES

MAPP 21430

F



JESSICO® FUSION

Contains 50 g/L Fenpicoxamid and
37.5 g/L Isoflucypram



Danger

Causes skin irritation.

May cause an allergic skin reaction.

Causes serious eye damage.

May cause respiratory irritation.

Suspected of damaging fertility

**Very toxic to aquatic life with long
lasting effects.**

Obtain special instructions before use.

Wear protective gloves/protective clothing/eye
protection/face protection.

IF IN EYES; Rinse cautiously with water for several
minutes. Remove contact lenses, if present and easily
to do. Continue rinsing.

IF exposed or concerned: Call a POISON CENTRE/
doctor/ physician.

Immediately call a POISON CENTER/doctor/physician.
Collect spillage.

Dispose of contents/container to a licensed hazardous-
waste disposal contractor or collection site except for
empty clean containers which can be disposed of as
non-hazardous waste.

**To avoid risks to human health and the environment,
comply with the instructions for use.**

IMPORTANT INFORMATION FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crops:	Wheat, rye, triticale and spelt
Maximum individual dose:	1.5 litres product per hectare
Latest time of application:	BBCH 59 (inflorescence fully emerged)
Aquatic buffer zone (metres):	12
Other specific restrictions:	Do not apply via hand-held equipment.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER
THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.
FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION
PRODUCTS.**



To access the **Safety Data Sheet** for this
product scan the code or use the link below:
[https://cropsscience.bayer.co.uk/our-products/
fungicides/jessico-fusion](https://cropsscience.bayer.co.uk/our-products/fungicides/jessico-fusion)
or alternatively contact your supplier

Bayer

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

	 GLOVES	 COVERALLS	 PROTECTIVE EYEWEAR
HANDLING THE CONCENTRATE	Chemical resistant	Yes	Faceshield

- WHEN USING DO NOT EAT, DRINK OR SMOKE.
- WASH ANY CONTAMINATION from eyes immediately.
- WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental Protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water/ Avoid contamination via drains from farmyards and roads)

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements. HORIZONTAL BOOM SPRAYERS MUST BE FITTED WITH THREE STAR DRIFT REDUCTION TECHNOLOGY. Low drift

spraying equipment must be operated according to the specific conditions stated in the official three star rating for that equipment as published on HSE Chemicals Regulation Divisions website. Maintain three star operating conditions until 30 m from the top of the bank of any surface water bodies.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within the distance specified for the crop to the top of the bank of a static or flowing water, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water. NOTE: BUFFER ZONES OF MORE THAN 5 m CANNOT BE REDUCED UNDER THE LOCAL ENVIRONMENT RISK ASSESSMENT FOR PESTICIDES (LERAP) SCHEME.

The statutory buffer zone must be maintained and the distance recorded in the LERAP record form. The LERAP record form must be kept available for three years.

Storage and Disposal

- KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.
- KEEP OUT OF REACH OF CHILDREN
- KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.
- WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.
- DO NOT RE-USE CONTAINER for any purpose.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Jessico Fusion is a mixture of a isoflucypram (a SDHI carboxamide fungicide) and fenpicoxamid (a picolinamide fungicide) recommended for control of a wide range of diseases on wheat,rye, triticale and spelt.

PROTECT FROM FROST

APPLICATION

	CROPS	All commercial varieties of winter and spring wheat, rye, triticale and spelt.
	SPRAY QUALITY	Apply as a MEDIUM spray quality. <i>(as defined by BCPC).</i>
	VOLUME AND PRESSURE	Apply in 150–300 litres per hectare water. A spray pressure of 2-3 bar is recommended. The higher spray volumes are recommended where the crop is dense to ensure good penetration to the lower leaves and stem bases. Disease control may be compromised by reducing water volumes, where good spray coverage is difficult to achieve.
	LATEST PERMITTED TIMING	Wheat, Rye, Triticale and Spelt From BBCH 30 to 59
	RATE	1.5 Litres per hectare. Horizontal boom sprayers must be fitted with three star drift reduction technology for all uses

DISEASES CONTROLLED

For foliar disease apply Jessico Fusion at first signs of disease before disease is established and use as part of an integrated crop management strategy.

TARGET DISEASE	CONTROL CLAIM (C=CONTROL)				APPLICATION COMMENTS
	WHEAT (Winter and Spring)	TRITICALE (Winter and Spring)	RYE (Winter and Spring)	SPELT	
Septoria leaf blotch	C	C	C	C	Apply as part of integrated crop management programme preferably in a protective situation.
Yellow Rust	C	C			Applications made to established infections are likely to be less effective. Apply in a programme with optimum timing for your conditions.
Brown rust	C	C		C	Applications made to established infections are likely to be less effective Apply in a programme with optimum timing for your conditions.

MIXING

STEP 1	STEP 2	STEP 3	STEP 4
Thoroughly shake the pack before use.	Add the required quantity of Jessico Fusion to the half-filled spray tank with the agitation system in operation and then fill to the required level.	Spray immediately after mixing.	Continue agitation at all times during spraying and stoppages until the tank is completely empty.

CLEANING AND BOOM HEIGHT

- Sprayers should be thoroughly cleaned with water and detergent after use, and filters and jets checked for damage and blockages.
- Boom height should be adjusted to ensure even coverage of the crop, particularly at later growth stages.
- The correct height is one at which the spray from alternate nozzles meets just above the crop, in dense crops, at later growth stages, higher water volumes should be used.

RESISTANCE STRATEGY

PATHOGENS	APPLICATION LIMITS	ICM STRATEGY	TANK MIX/ALTERNATE
Jessico Fusion is a formulated mixture of a Qil fungicide and an SDHI from different chemical groups. Resistance to some SDHI fungicides has been identified in some pathogens which may seriously affect the performance of some products.	No more than two application timings of SDH inhibitors must be applied to the same cereal crop. No more than one foliar application of products containing Qil to any cereal crop	Use Jessico Fusion as part of an integrated crop management strategy incorporating use of different modes of action within the programme and other non-chemical methods of disease reduction such as use of resistant varieties.	Tank mixtures or alternation with fungicides having a different mode of action against the diseases present have been shown to protect against the development of resistant forms of disease.

- There is a risk of widespread resistance occurring on *Zymoseptoria tritici* in the UK. Failure to follow resistance management action may result in reduced levels of disease control.
- Jessico Fusion should be used preventively and should not be relied on for its curative potential.
- Strongly reduced rate programs including multiple applications must not be used.



For further advice on resistance management in Qil's and SDHI's contact your agronomist or specialist advisor, and visit the FRAG-UK website.

Company advisory information:

The possible development of disease strains resistant to Jessico Fusion cannot be excluded or predicted. Where such resistant strains occur, Jessico Fusion is unlikely to give satisfactory control.

Jessico® Fusion contains isoflucypram and fenpicoxamid.

Iblon® is the commonly used name of isoflucypram active substance.

Inatreq™ active is the commonly used name of fenpicoxamid active substance.

Iblon® and Jessico® are Registered Trademarks of Bayer.

Inatreq™ is a trademark of Corteva Agriscience and its affiliated companies.

© Bayer CropScience Limited 2025