

Fungicide

RECITAL

A mixture of a carboxamide and triazolinthione fungicide for disease control in oilseed rape.

**A suspo emulsion formulation containing
125 g/L (12% w/w) fluopyram and
125 g/L (12% w/w) prothioconazole.**

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Bayer CropScience Ltd, PO Box 1582, Cambridge, CB1 0FE
Telephone: 01223 226500

**For 24 hour emergency
information contact
Bayer CropScience Ltd
Telephone:
0330 678 3382 (24 hr)**

National Poisons Information
Centre UK: 0344 892 0111
(medical professionals only)

National Poisons Information
Centre Dublin: +353 1 809 2166

5 L e



GROUP 7 | 3 FUNGICIDES

MAPP 17909

F



RECITAL®

UFI: 6E80-S0DX-N00H-NC5Q

Contains 125 g/L (12% w/w) fluopyram
and 125 g/L (12% w/w) prothioconazole

**Warning****Very toxic to aquatic life with long lasting effects.**

Wear protective gloves/protective clothing/
eye protection/face protection.

Protect from sunlight.

Dispose of contents/container to a licensed
hazardous-waste disposal contractor or
collection site except for empty clean
containers which can be disposed of as
non-hazardous waste.

Contains 1,2-benzisothiazolin-3-one, reaction mass of: 5-chloro-2-methyl-4-isothiazolin-3-one and 2-methyl-4-isothiazolin-3-one (3:1), 2-[2-(1-chlorocyclopropyl)-2-hydroxy-3-phenylpropyl]-2,4-dihydro-3H-1,2,4-triazole-3-thione. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crops:	Oilseed rape
Maximum individual dose:	1.0 litres product per hectare
Maximum total dose:	2.0 litres product per hectare
Latest time of application:	56 days before harvest
Other Specific Restrictions:	A minimum interval of 14 days must be observed between applications.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

PROTECT
FROM FROST



To access the **Safety Data Sheet** for this product scan the code or use the link below:

<https://cropscience.bayer.co.uk/our-products/fungicides/recital>

or alternatively contact your supplier

Bayer

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate, handling contaminated surfaces or applying the product. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH ANY CONTAMINATION from eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

Environmental Protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme.

Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely. DO NOT RE-USE CONTAINER for any purpose.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Recital is a mixture of a carboxamide and triazolinthione fungicide recommended for the control of a wide range of diseases on oilseed rape.

PROTECT FROM FROST

CROPS

Recital may be used on all commercial varieties of oilseed rape.

RATE OF USE

Apply Recital at 1.0 litre per hectare.
The maximum total dose per crop is:-
2.0 litre per hectare.

APPLICATION

Water volume

Apply Recital in 200-300 litres per hectare water. The higher spray volumes are recommended where the crop is dense or disease pressure / risk is high to ensure good penetration to the lower leaves and stem bases. Disease control may be compromised by reducing water volumes, where good spray coverage is difficult to achieve.

Qualified use recommendation

If required, water volumes down to 100 litres per hectare may be used; however

these have not been supported by effectiveness or crop safety data.

A spray pressure of 2-3 bar is recommended.

Spray quality

Apply as a MEDIUM spray quality (as defined by BCPC).

Latest Permitted Timing

Recital may be applied at any stage up to 56 days before harvest.

Mixing

Thoroughly shake the pack before use.

Add the required quantity of Recital to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

General

Sprayers should be thoroughly cleaned with water and detergent after use, and filters and jets checked for damage and blockages.

Boom height should be adjusted to ensure even coverage of the crop, particularly at later growth stages. The correct height is one at which the spray from alternate nozzles meets just above the crop, in dense crops, at later growth stages, higher water volumes should be used.

DISEASES CONTROLLED

Winter oilseed rape

Light leaf spot, *Phoma* leaf spot and stem canker*, *Sclerotinia* stem rot and powdery mildew.

* Moderate control.

APPLICATION TIMING

Depending upon disease pressure and choice of management programme one application of Recital may be made in the autumn followed by a second in the spring/summer. Alternatively two applications may be made in the spring/summer.

WINTER OILSEED RAPE

Light Leaf Spot

Apply Recital protectively in the autumn/winter (usually late October to early December) after the first two leaves have unfolded. Follow up spray(s) may be required in early spring from the onset of stem elongation, depending on disease development.

Phoma Leaf spot/Stem Canker

Apply Recital in autumn at the first sign of disease (after the first two leaves have unfolded). A repeat application with another effective product may be necessary in late autumn / early winter should disease symptoms reoccur. Recital will provide moderate control of *phoma* stem canker.

Sclerotinia stem rot (*Sclerotinia sclerotiorum*).

Apply Recital at early to full flower.

Powdery Mildew

Apply Recital at first signs of disease (after first two leaves unfolded stage) if infection occurs in autumn, or at early to mid flower in spring. Where disease pressure remains high application with an alternative effective product may be repeated.

Spring oilseed rape

(QUALIFIED MINOR USE RECOMMENDATION)

Recital can also be used on varieties of spring oilseed rape but crop safety has not been fully established.

RESISTANCE STRATEGY

Strains of light leaf spot resistant to azole fungicides are known to exist. To avoid development of resistance apply product protectively in response to disease forecasts. Where possible, when light leaf spot is present, avoid the use of azole based fungicides when targeting other diseases such as *Sclerotinia* at mid flowering.

For further advice on resistance management in oilseed rape contact your agronomist, or specialist advisor, and visit the FRAG-UK website.

CAUTION: The possible development of disease strains resistant to Recital cannot be excluded or predicted. Where such resistant strains occur, Recital is unlikely to give satisfactory control.

© Recital is a registered Trade Mark of Bayer

© Bayer CropScience Limited 2025