

# OVATION<sup>®</sup>

(H) – Herbicide

A foliar applied translocated herbicide for the control of annual and perennial grass and broad-leaved weeds before sowing or planting of all crops.

For use pre-emergence and pre-harvest in cereals and certain other crops, for destruction of grassland, and use in stubbles and orchards, and non-crop areas.  
For control of emerged weeds in amenity and forestry situations.

Degraded by micro-organisms/microbes in the soil.

A soluble concentrate containing 360 g/L glyphosate, present as 441 g/L (35.3% w/w) of the potassium salt of glyphosate

**This product must only be used in England, Scotland and Wales.**

GROUP 9 HERBICIDE

**MAPP Number 21195**

PROTECT FROM FROST

## Ovation

UFI: KPF1-E0Y7-K009-RH2Q

Contains 360 g/L glyphosate, present as 441g/L (35.3% w/w) of the potassium salt of glyphosate.

### WARNING

Causes serious eye irritation.

Toxic to aquatic life with long lasting effects.

Wash hands thoroughly after handling.

Wear protective gloves/protective clothing/eye protection/face protection.

Keep only in original container.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If eye irritation persists: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**



The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work

**CONTENTS:**

**20L**itres





RECOMMENDATION TABLES

| AREA OF USE                                                           | TARGET WEEDS/USAGE                                           | CROP/SITUATION                                                                                                 | WEED INFESTATION                                                                                 | APPLICATION RATE L/ha | WATER VOLUME                                | APPLICATION TIMING AND GUIDANCE                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|-----------------------------------------------------------------------|--------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------|-----------------------|---------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PRE-HARVEST ARABLE CROPS                                              | Common Couch                                                 | WINTER and SPRING WHEAT, DURUM WHEAT, WINTER and SPRING BARLEY and WINTER and SPRING OATS                      | 1 to 25 shoots/m <sup>2</sup><br>Up to 75 shoots/m <sup>2</sup><br>Over 75 shoots/m <sup>2</sup> | 2.0<br>3.0<br>4.0     | 80-250 L/ha*                                | Grain/seed moisture must not exceed 30% at spraying.<br><br>Harvest intervals:<br>CEREALS, PEAS, BEANS 7+ days<br>OILSEED RAPE 14-21 days<br>LINSEED 14-28 days<br>MUSTARDS 8-10 days<br><br>Use high clearance, narrow wheeled tractors, wide booms and crop dividers.<br>Where desiccating crops, check susceptibility of any weeds present.<br><br>Do not attempt to desiccate OILSEED RAPE or MUSTARD crops with significant secondary growth, uneven maturity, disease or stress.<br><br>Desiccate LINSEED when seeds are light brown and capsules brown; stems/leaves may be yellow/green.<br>Effects on brewing and baking have not been tested. Consult grain merchant or processor before use. |
|                                                                       |                                                              | OILSEED RAPE AND MUSTARDS                                                                                      | Up to 75 shoots/m <sup>2</sup><br>Over 75 shoots/m <sup>2</sup>                                  | 3.0<br>4.0            | 100-250 L/ha#                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|                                                                       |                                                              | COMBINING PEAS FIELD BEANS                                                                                     | Up to 75 shoots/m <sup>2</sup><br>Over 75 shoots/m <sup>2</sup>                                  | 3.0<br>4.0            | 80-250 L/ha*                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|                                                                       |                                                              | LINSEED                                                                                                        | Up to 75 shoots/m <sup>2</sup><br>Over 75 shoots/m <sup>2</sup>                                  | 3.0<br>4.0            | 80-250 L/ha*                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|                                                                       |                                                              |                                                                                                                |                                                                                                  |                       |                                             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|                                                                       | Perennial broad-leaved weeds and other perennial grasses     | WINTER and SPRING WHEAT, DURUM WHEAT, WINTER and SPRING BARLEY and WINTER and SPRING OATS                      | All levels of all species                                                                        | 4.0                   | 80-250 L/ha*                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|                                                                       |                                                              | OILSEED RAPE AND MUSTARDS                                                                                      | All levels of all species                                                                        | 4.0                   | 100-250 L/ha#                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|                                                                       |                                                              | COMBINING PEAS AND FIELD BEANS                                                                                 | All levels of all species                                                                        | 4.0                   | 80-250 L/ha*                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|                                                                       |                                                              | LINSEED                                                                                                        | All levels of all species                                                                        | 4.0                   | 80-250 L/ha                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|                                                                       | Harvest management                                           | WINTER and SPRING WHEAT, DURUM WHEAT, WINTER and SPRING BARLEY and WINTER and SPRING OATS                      | Annual grasses, crop stems and leaves<br>Annual broad-leaved weeds                               | 1.0 (+)<br>1.5 (+)    | 80-250 L/ha*                                | DO NOT TREAT CROPS GROWN FOR SEED.<br><br>* Rotary atomisers may be used at a water volume of 40 L/ha. Ensure droplet diameter falls within the range 200-300 microns.<br># Use higher volumes for dense canopies.<br>(+) For optimum results use an approved adjuvant at 0.5% spray solution as described in 'Compatability' section.                                                                                                                                                                                                                                                                                                                                                                  |
|                                                                       | Crop desiccation and annual weeds, prior to direct combining | OILSEED RAPE MUSTARDS                                                                                          | All levels/species                                                                               | 3.0                   | 100-250 L/ha#                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|                                                                       |                                                              | LINSEED                                                                                                        | All levels/species                                                                               | 3.0                   | 80-250 L/ha                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| ALL EDIBLE AND NON-EDIBLE CROPS (DESTRUCTION, BEFORE SOWING/PLANTING) |                                                              | Vegetation management                                                                                          | Annual weeds<br>Perennial grasses<br>Perennial broad-leaved weeds                                | 1.5<br>4.0<br>5.0     | 80-250 L/ha* or hand-held equipment (p. 13) | Do not use in or alongside hedgerows<br>Do not use under polythene or glass.<br>Apply the annual weed dose at least 2 days before sowing/planting.<br>Apply at perennial weed doses at least 5 days before sowing/planting.<br>* Rotary atomisers may be used at a water volume of 40 L/ha. Ensure droplet diameter falls within the range 200-300 microns                                                                                                                                                                                                                                                                                                                                              |
| POST SOWING/PLANTING, PRE-EMERGENCE OF THE CROP                       | Volunteer cereals and annual weeds                           | LISTED CEREALS<br>OILSEED RAPE, MUSTARD, LINSEED, PEAS, FIELD BEANS, SUGAR BEET, SWEDE, TURNIP, ONION and LEEK | All levels/species                                                                               | 1.5                   | 80-250 L/ha*                                | CAUTION - Ensure that spraying precedes ANY crop emergence.<br>* Rotary atomisers may be used at a water volume of 40 L/ha. Ensure droplet diameter falls within the range 200-300 microns.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |

| AREA OF USE                                                             | TARGET WEEDS/USAGE                                             | CROP/SITUATION                                                                     | WEED INFESTATION                                                                                                                                                                                                | APPLICATION RATE L/ha                        | WATER VOLUME                                                                      | APPLICATION TIMING AND GUIDANCE                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|-------------------------------------------------------------------------|----------------------------------------------------------------|------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|-----------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| STUBBLES, PRE-SOWING and PRE-PLANTING                                   | Common Couch                                                   | BEFORE ALL CROPS EXCEPT ORCHARDS                                                   | Up to 75 shoots/m <sup>2</sup><br>Over 75 shoots/m <sup>2</sup>                                                                                                                                                 | 3.0<br>4.0                                   | 80-250 L/ha*                                                                      | Do not cultivate immediately before spraying.<br>For PERENNIAL weed control, allow:<br>– 21+ days growth before spraying in spring<br>– VOLUNTEER POTATOES to make ample top growth<br>– 5 days before cultivating or drilling<br><br>For ANNUAL weed control, allow:<br>– 24 hours before cultivating<br>– 48 hours before drilling<br><br>Allow 7 days before planting trees<br>* Rotary atomisers may be used at a water volume of 40 L/ha. Ensure droplet diameter falls within the range 200-300 microns.<br>(+) For optimum results use an approved adjuvant at 0.5% spray solution as described in 'Compatibility' section.                                                                                                                                                                                                                    |
|                                                                         | Other perennial grasses; volunteer potatoes (autumn only)      |                                                                                    | All levels of all species                                                                                                                                                                                       | 4.0                                          |                                                                                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|                                                                         | Volunteer cereals and annual weeds                             |                                                                                    | All levels of all species                                                                                                                                                                                       | 1.5 (+)                                      |                                                                                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|                                                                         | Perennial broad-leaved weeds                                   |                                                                                    | All levels of all species                                                                                                                                                                                       | 5.0                                          |                                                                                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|                                                                         | Perennial grasses and broad-leaved weeds                       | BEFORE ORCHARD PLANTING                                                            | Arable weeds<br>Pasture weeds                                                                                                                                                                                   | 4.0<br>5.0                                   |                                                                                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| GREEN COVER ON LAND NOT BEING USED FOR CROP PRODUCTION E.G. 'SET ASIDE' | Common Couch                                                   | BEFORE or DURING REMOVAL FROM PRODUCTION e.g. prior to growing a set aside mixture | Up to 75 shoots/m <sup>2</sup><br>Over 75 shoots/m <sup>2</sup>                                                                                                                                                 | 3.0<br>4.0                                   | 80-250 L/ha* or hand-held equipment (p. 13) or tractor mounted weed wiper (p. 13) | Before using on land taken out of production as part of a grant aided scheme, ensure compliance with the management rules of that scheme.<br>Do not 'top' or cultivate immediately before application.<br>For PERENNIAL weed control, allow:-<br>– 21+ days growth before spraying in spring<br>– 5 days before cultivating or drilling.<br>For ANNUAL weed control, allow:-<br>– 24 hours before cultivating.<br>Do not direct drill after set aside.<br>Avoid applications during stem elongation as reduced control and re-spray is likely<br>* Rotary atomisers may be used at a water volume of 40 L/ha. Ensure droplet diameter falls within the range 200-300 microns.<br>Best control of annual grasses is achieved between full ear emergence and senescence.<br>+Only for weeds listed as per grassland destruction application rate table. |
|                                                                         | Perennial broad-leaved weeds and other perennial grasses       |                                                                                    | All levels/species                                                                                                                                                                                              | 4.0                                          |                                                                                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|                                                                         | Annual weeds:<br>• Early autumn/spring<br>• Late spring/summer |                                                                                    | All levels/species<br>All levels/species                                                                                                                                                                        | 1.5<br>3.0                                   |                                                                                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|                                                                         | Natural regeneration and cover crop destruction                | AFTER SHORT ROTATION or LONG TERM REMOVAL FROM PRODUCTION                          | Annual weeds only<br>Perennial grasses<br>Perennial broad-leaved weeds<br>Perennial broad-leaved weeds as listed below.                                                                                         | 3.0<br>4.0<br>5.0<br>6.0+                    | 150-250 L/ha*                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| PERMANENT GRASSLAND (DESTRUCTION)                                       | Short rotation Ryegrass, longer leys and permanent pasture     | GRASS                                                                              | Short rotation Ryegrass with annual weeds<br>Leys 2-4 years old with perennial grass weeds<br>Long leys 4-7 years old with perennial broad-leaved weeds<br><br>Permanent pasture<br>See Weed Table on p. 9 - 10 | 3.0 L/ha<br>4.0 L/ha<br>5.0 L/ha<br>6.0 L/ha | 150-250 L/ha                                                                      | Treat EITHER before grazing/mowing in June-Oct, when growth is 30-60 cm, not dense and lacking mature seeds, OR re-growth after grazing/mowing.<br>Select the application rate which controls/destroys the least susceptible weed and grass species present in the sward.<br>Grass may be conserved or grazed by cattle, dairy cows or sheep 5+ days after spraying. REMOVE POISONOUS PLANTS BEFORE GRAZING/MOWING.<br>If Ragwort is present, the guidance in the 'DIRECTIONS FOR USE' must be followed.<br>ONLY direct drill grass and clover EITHER into 1-2 year leys without mat, 5+ days after spraying, OR long leys with some mat, in the spring following autumn application.                                                                                                                                                                 |

| APPLICATION RATE FOR GRASSLAND DESTRUCTION                                                                                          |                                                                                              |                                                                                                       |                                                                                                |                                                                                                                            |                                                                                                                                                                           |                                                                                                                                                                                                                                                                                                                                                                                                |
|-------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 3.0 L/ha                                                                                                                            |                                                                                              | 4.0 L/ha                                                                                              |                                                                                                | 5.0 L/ha                                                                                                                   |                                                                                                                                                                           | 6.0 L/ha                                                                                                                                                                                                                                                                                                                                                                                       |
| Annual Meadow-grass<br>Common Mouse-ear<br>Italian Rye-grass<br>Meadow Fescue<br>Rough Meadow-grass<br>Timothy                      | Common Chickweed<br>Dock Seedlings<br>Mayweed species<br>Meadow Foxtail<br>Speedwell species | Black-bent<br>Cock's-foot<br>Common Couch<br>Creeping Soft-grass<br>Perennial Rye-grass<br>Soft Brome | Broad-leaved Dock<br>Common Bent<br>Creeping Bent<br>Curled Dock<br>Plantains<br>Yorkshire Fog | Bracken**<br>Common Sorrel<br>Creeping Thistle<br>Dwarf Thistle<br>Red Clover<br>Sheep's Sorrel<br>Spear Thistle<br>Yarrow | Common Nettle<br>Creeping Buttercup<br>Daisy<br>Perennial Sow-thistle<br>Sedges<br>Soft Rush<br>Tufted Hairgrass                                                          | Common Ragwort<br>Heath Rush<br><i>Molinia</i> (Purple Moor-grass)<br>Red Fescue<br>White Clover*                                                                                                                                                                                                                                                                                              |
| * White Clover is best cut in June and sprayed one month later. ** At full frond expansion                                          |                                                                                              |                                                                                                       |                                                                                                |                                                                                                                            |                                                                                                                                                                           |                                                                                                                                                                                                                                                                                                                                                                                                |
| AREA OF USE                                                                                                                         | TARGET WEEDS/USAGE                                                                           | CROP/SITUATION                                                                                        | WEED INFESTATION                                                                               | APPLICATION RATE L/ha                                                                                                      | WATER VOLUME                                                                                                                                                              | APPLICATION TIMING AND GUIDANCE                                                                                                                                                                                                                                                                                                                                                                |
| ORCHARDS                                                                                                                            | Perennial grasses and broad-leaved weeds<br>Root suckers                                     | WITHIN ORCHARDS OF APPLE, PEAR, PLUM, CHERRY, DAMSON                                                  | All levels of most species<br><br>All species                                                  | 5.0 L/ha<br><br>5.0 L/ha                                                                                                   |                                                                                                                                                                           | Trees must have been established for 2 years before spraying.<br>Spray AFTER autumn leaf-fall and BEFORE:<br>Apples, pears – green cluster stage<br>Stone fruit – white bud stage<br><br>Avoid contact with tree branches and trunks above 30 cm from the ground.<br>Treat suckers in late spring only.                                                                                        |
| IN-CROP (TRACTOR-MOUNTED WEED WIPER) APPLICATION                                                                                    | Bolters, weed beet, other weeds                                                              | ARABLE CROPS AND GRASSLAND SET ASIDE                                                                  | All levels                                                                                     |                                                                                                                            | 1:1 dilution with water OR<br>1:2 dilution with water in hot, dry conditions.<br>For 'new generation' wipers consult the manufacturer for guidance.                       | Weeds must be 10+ cm taller, and wiper 5+ cm higher, than desired vegetation.<br>Wipe dense populations twice, in opposite directions. BOLTING BEET requires three applications, 2 weeks apart, from early July to early August.<br>Contact Bayer or your distributor for specific recommended weed wiper applicators.<br>POISONOUS WEEDS and grazing/mowing interval – See GRASSLAND section. |
| NATURAL SURFACES NOT INTENDED TO BEAR VEGETATION, PERMEABLE SURFACES OVERLYING SOIL. ALL SITUATIONS (DESTRUCTION, BEFORE PLANTING). | Vegetation management                                                                        | Including farmyards roadsides, paths, and along fences and walls                                      | Annual weeds<br>Perennial grasses and broad-leaved weeds                                       | 4.0-5.0                                                                                                                    | Hydraulic sprayers (boom and knapsack) at water volumes 80-400 L/ha or rotary atomisers* at water volumes 40 L/ha or hand-held equip-ment. See Mixing & Spraying section. | Do not use under polythene or glass.<br><br>* Where rotary atomisers are used their droplet diameter must fall within the range 200-300 µm.                                                                                                                                                                                                                                                    |
| HARD SURFACES                                                                                                                       | Vegetation management                                                                        | Including farmyards roadsides, paths, hard surfaces and along fences and walls                        | Annual weeds<br>Perennial grasses and broad-leaved weeds                                       | 1.5<br>4.0-5.0                                                                                                             | Hydraulic sprayers (boom and knapsack) at water volumes 80-400 L/ha or rotary atomisers* at water volumes 40 L/ha or hand-held equip-ment. See Mixing & Spraying section. | Apply this product carefully. Ensure spraying takes place only when weeds are actively growing (normally March to October) and is confined to only visible weeds including those in the 30cm swath covering the kerb edge and road gully – do not overspray drains.                                                                                                                            |

FORESTRY/FARM FORESTRY WEED CONTROL

Ovation can be used for site preparation and for weed control in planted out trees

| AREA OF USE                                                                                | TARGET WEEDS/USAGE                                   | WEED INFESTATION                                                                                                                                                                                                                  | APPLICATION RATE L/ha                                                | WATER VOLUME                                                                                                                                                                     | APPLICATION TIMING AND GUIDANCE                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|--------------------------------------------------------------------------------------------|------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| FORESTRY:<br>- PRE-PLANTING                                                                | Arable land, planting, replanting, & grassland areas | Arable weeds<br>Grassland weeds                                                                                                                                                                                                   | 4.0<br>5.0                                                           | Hydraulic sprayers:<br>80-250 L/ha<br>or<br>rotary atomisers: 40 L/ha*                                                                                                           | All tree species may be planted 7 days or more after treatment.<br>*Where rotary atomisers are used their droplet diameter must fall within the range 200-300µm.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| FORESTRY:<br>- POST-PLANTING (DIRECTED) IN CONIFERS & BROAD-LEAVED TREES                   | Clean-up around trees with knapsack applicators      | Annual/perennial grasses and broad-leaved weeds                                                                                                                                                                                   | 4.0                                                                  | Hand held equipment.<br>Knapsack: Apply as a 2% concentration or<br>Weed wiper mini: apply as a concentration of 1 part Ovation to 2 parts water (see Mixing & Spraying section) | It is ESSENTIAL to use a TREE GUARD for all applications made in the growing season.<br>Treat bracken after frond tips are unfurled but before senescence.<br>Treat heather late August to end September.<br>All other woody weeds are treated June to August, before leaf senescence but after new growth of crop has hardened.<br>Application using a weed wiper is not suitable.<br>(*) For improved control of Rhododendron apply 8.0 L/ha OVATION, adding Mixture B (ADJ 0570) at 2% of spray volume.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|                                                                                            |                                                      | Woody weeds:<br>Bracken/Beech<br>Brush/Brambles<br>Sycamore/Oak/Hazel/Willow/ Ash (excluding Rhododendron)                                                                                                                        | 3.0                                                                  |                                                                                                                                                                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|                                                                                            |                                                      | Heather (peat soils)<br>Heather (mineral soils)<br>Rhododendron                                                                                                                                                                   | 4.0<br>6.0<br>10.0                                                   |                                                                                                                                                                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| FORESTRY:<br>- POST-PLANTING (OVERALL DORMANT SEASON IN CERTAIN CONIFERS – CONIFER RELEASE | Grass weeds:<br>- Lowland areas<br>- Upland areas    | Black Bent, Cock's-foot, Common Couch, Creeping Soft-grass, False Oat-grass, Fescues, Meadow-grasses, other Bent species, Purple Moor-grass, Sweet Vernal-grass, Tufted Hair-grass, Wavy Hair-grass, Wood Small-reed (Bush grass) | 1.5<br>2.0                                                           | Hydraulic sprayers:<br>200-250 L/ha<br>or<br>hand-held equipment – see 'Mixing and Spraying' section                                                                             | DO NOT OVERALL SPRAY trees being grown for ORNAMENTAL PURPOSES, including CHRISTMAS TREES. Species safe to spray when fully dormant and leader growth has hardened: Corsican, Lodgepole and Scots Pine, Norway Spruce, Sitka Spruce, Lawson Cypress, Western Red Cedar, Douglas Fir and Noble Fir – safe to spray when fully dormant and leader growth has hardened but NOT in spring. If overall application takes place after the optimum timing weed control may be reduced. It is advisable to spray a limited area of forest to test crop safety under local conditions before widespread overall application in subsequent years. These recommended application rates refer to forestry usage only.<br>Inadequate control may result if used in other areas. Caution: The timing of hardening of leader growth varies considerably between locations and between seasons. It may occur as early as the end of July or be delayed to October or later. To avoid damage to Lammas growth, sprays should be directed away from leaders. |
| FORESTRY:<br>- STUMP APPLICATION FOR CHEMICAL THINNING                                     | Deciduous trees<br>Coniferous trees                  | All species<br>All species                                                                                                                                                                                                        | 10% solution of OVATION in water<br>20% solution of OVATION in water | Hydraulic sprayers:<br>200-250 L/ha<br>or<br>hand-held equipment – see 'Mixing and Spraying' section                                                                             | Apply the solution to saturate the rim of the newly cut surface, with a suitably adapted clearing saw, spot gun or paintbrush. Treat as soon as possible after felling, in the period November to March/April. Do not apply in the period of active sap flow in the spring/early summer. Do not cut trenches or drill holes and fill with the solution or use undiluted product.<br>Note: for ease of identification of treated areas a suitable, commercially available, water-soluble dye may be added to the prepared spray solution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| FORESTRY:<br>- CHEMICAL THINNING BY INJECTION OF TREE STEMS                                | Coniferous and deciduous species                     | -                                                                                                                                                                                                                                 | 2 ml neat OVATION per cut per 10 cm diameter (or less) of tree       |                                                                                                                                                                                  | Use a hatchet to cut one notch in trees up to 10cm diameter and apply 2 ml of the solution to each cut. Use two or three notches in trees over 10cm diameter. Do not treat in the period of active sap flow in the spring/early summer.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |