

Fungicide

Folicur



A broad spectrum systemic fungicide for wheat (excluding durum), barley, oats, rye (

Safety Information

FOLICUR®

UFI: NX80-T0JA-G00G-MDP3

IMPORTANT INFORMATION FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL FUNGICIDE

Crops:	All varieties of wheat (excluding Durum), barley, rye (winter), oats, oilseed rape, field beans and linseed.
---------------	--

WHEAT

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Linseed

Powdery mildew, *Botrytis*.

high repeat applications with alternative effective product – see “Resistance Management”.

Diseases Controlled

Folicur may be applied at any time before brown capsule stage or 35 days before harvest whichever is sooner.

Diseases Controlled – Application Timing

Powdery Mildew

Folicur applied at first signs of disease will give control.

Botrytis

Folicur applied at first signs of disease can give a reduction in this disease.

MIXING

Thoroughly shake the pack before use.

Add the required quantity of Folicur to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

Where tank mixes are used Folicur should be added to the spray tank last, after first dispersing the other product(s),

© Folicur is a registered Trade Mark of Bayer.

© Bayer CropScience Limited 2025