

P.O. Box 5569 A Yuma, AZ 85366-5569 A Phone (928) 783-8844 A FAX (928) 343-9255

FIFRA SECTION 24(C) SPECIAL LOCAL NEED LABEL FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OREGON

PYRAFLUFEN-ETHYL GROUP

Ida

EPA Reg. No. 10163-314, EPA SLN No.: OR-180017

For Control of Broadleaf Weeds on Grasses Grown for Seed.

This label is valid until December 31, 2028 or until withdrawn, cancelled, suspended, or otherwise amended.

ACTIVE INGREDIENT:

Contains 0.208 lb. pyraflufen ethyl per gallon (25 grams per liter)
*contains petroleum distillates

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

ENVIRONMENTAL HAZARDS

Follow all applicable directions, restrictions, and precautions in the Environmental Hazards section of the label affixed to the main container of Vida Herbicide (EPA Reg. No. 10163-314).

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the label affixed to the main container of Vida Herbicide must be in the possession of the user at the time of pesticide
 application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on this SLN label and the label affixed to the main container of Vida Herbicide.

CROP		container of Vida Herbicide.
CROP	FL OZ/ACDE	USE INSTRUCTIONS
	OZ/ACRE	
GRASSES GROWN FOR SEED	1 - 5.5	 RESTRICTIONS Do not make more than 2 applications per year. Do not apply more than 5.5 fl oz Vida Herbicide (0.00894 lb pyraflufen-ethyl) per acre per year. Allow a minimum of 30 days between applications. Apply in a minimum of 10 gallons spray solution per acre by ground. Addition of a non-ionic surfactant (NIS) at 0.25% v/v is recommended for optimum weed control. Do not apply this product by aerial application. Apply to emerged actively growing weeds before seeding up to the jointing stage of the crop. Susceptible broadleaf weeds are controlled or suppressed up to 4 inches in height or less, or rosettes of 3 inches in diameter or less. Use the higher labeled rate for hard-to-control weeds. Refer to the Weeds Controlled section of the label affixed to the main container of Vida Herbicide for more information. Refer to the Tank Mixtures section of the label affixed to the main container of Vida Herbicide when tank mixing with other pesticides labeled for use on grasses grown for seed. PRECAUTIONS Cool season grasses, both newly seeded and established, are generally tolerant of Vida herbicide at labeled rates. To evaluate tolerance of species/varieties in question, apply to a small test plot before treating large areas. Yellowing/discoloration or leaf speckling may occur on some varieties 3 to 5 days following application. Full recovery is typically seen 4 to 7 days after application.

Vida® is a registered trademark of Gowan Company, LLC.

Section 24(c) Registrant:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569
Rev. 09-19-23