

SUPPLEMENTAL LABELING

Buckle® Microactiv Herbicide

EPA Reg. No. 10163-289

**GRANULAR SELECTIVE HERBICIDE FOR SPRING PREPLANT APPLICATION TO CONTROL WILD OATS AND
PIGEONGRASS IN SPRING WHEAT.**

**READ AND FOLLOW THE ENTIRE LABEL FOR BUCKLE® MICROACTIV HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS
CONTAINED IN THIS SUPPLEMENTAL LABELING.**

When using Buckle Microactiv Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

FOR USE WITHIN NORTH DAKOTA EXCLUDING THE FOLLOWING COUNTIES: CASS, GRAND FORKS, PEMBINA, RICHLAND, TRAILL, AND WALSH.

THIS PRODUCT MAY ALSO BE USED IN THE FOLLOWING SOUTH DAKOTA COUNTIES: CORSON, DEWEY, HARDING, PERKINS, AND ZIEBACH.

KEEP OUT OF REACH OF CHILDREN CAUTION

In case of a medical emergency involving this product, call toll free 1-888-487-0798.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with the labeling.

This label must be in possession of the user at the time of herbicide application.

Apply Buckle Microactiv Herbicide at a rate of 10 pounds per acre in the spring prior to planting Spring Wheat. Application is limited to one per growing season and must not exceed 10 lbs. of this product per acre.

Incorporate spring Buckle Microactiv Herbicide applications within 24 hours with a field cultivator with harrow attachment. A second incorporation at a right angle to the first will provide the best results. Delay the second incorporation or spring wheat planting for a minimum of 5 days following the first incorporation. Do not set tillage equipment to incorporate soil deeper than the first incorporation.

Do not apply Buckle Microactiv Herbicide for spring wheat on soil that has been treated with 0.5 lbs ai/acre or greater of trifluralin, or other dinitroaniline herbicide, the previous year.

NOTE: Spring wheat varieties identified as susceptible to early injury from preplant applications of Buckle Microactiv Herbicide include: Alex, Amidon, Butte 86, Ellar, Era, Erik, Lew and Newana.

To minimize stress and allow for uniform germination, a seeding depth of no more than 1.5 to 2 inches is recommended. Delay spring planting of spring wheat until the average soil temperature at the 2-inch depth is 45 degrees F or greater.

DO NOT APPLY BUCKLE MICROACTIV HERBICIDE TO HILLTOPS.

Seed treatments are recommended to reduce potential stress to germinating seedlings from disease or insects.

Do not use on fields if the field is or will be irrigated in the current growing season.

NOTE: Slight stand reductions do not normally affect yield.

DO NOT APPLY BUCKLE MICROACTIV HERBICIDE TO A STRIP OR AREA IN THE FIELD THAT WILL SERVE AS AN UNTREATED COMPARISON FOR WEED CONTROL AND CROP TOLERANCE.

Read the "NOTICE OF CONDITIONS OF SALE AND WARRANTY LIABILITY LIMITATIONS" in the label for Buckle Microactiv Herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Buckle® is a registered trademark of Gowan Company LLC.