

Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937) Sivanto™ 200 SL

EPA Reg. No. 264-1141

For reduced application rates in leafy vegetables in control of aphids.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ALABAMA, ALASKA, COLORADO, DELAWARE, DISTRICT OF COLUMBIA, FLORIDA, GEORGIA, IDAHO, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, LOUISIANA, MASSACHUSETTS, MAINE, MICHIGAN, MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEVADA, NEW JERSEY, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, OREGON, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, UTAH, VERMONT, VIRGINIA, WASHINGTON, WEST VIRGINIA, WISCONSIN AND WYOMING

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

LEAFY VEGETABLES (except Brassica) - FOLIAR

Crops of Crop Group 4 Including: Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Cardoon, Celery, Celtuce, Chervil, Chinese celery, Chrysanthemum (edible-leaved and garland), Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive, Florence fennel (sweet anise, sweet fennel, Finocchio), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach [including New Zealand and vine (Malabar spinach, Indian spinach)]. Swiss chard. Taro leaves

Spirideri, indian Spirideriji, Owiss chard, Taro leaves	
Pests Controlled	Product Rate (fl oz/A)
Aphids	7.0 - 10.5

Foliar Application Restrictions:

Pre-Harvest Interval (PHI): 1 day

Minimum interval between applications: 7 days

Minimum application volumes: 10 gallons/Acre (Ground); 2 gallons/Acre (Aerial)

Maximum SIVANTO 200 SL allowed per crop season: 28.0 fluid ounces/Acre (0.365 lb Al/Acre).

Maximum crop seasons per year: 3

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577 For <u>PRODUCT USE</u> Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937)

Sivanto is a trademark of Bayer.

4/22/2015 Page 1 of 1