

SPECIMEN LABEL

IPCONAZOLE

GROUP

3

FUNGICIDE

ACE[®] 3.8L ST

ACTIVE INGREDIENT:	% BY WT.
Ipconazole* (CAS No. 125225-28-7).....	40.7%
OTHER INGREDIENTS:	59.3%
TOTAL:	100.0%

*Contains 3.86 pounds of active ingredient per gallon.

*Contains 462.5 grams of active ingredient per liter.

KEEP OUT OF REACH OF CHILDREN
MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

CAUTION / PRECAUCION

See inside booklet for complete First Aid, Precautionary Statements, Directions For Use, Storage and Disposal and Conditions of Sale and Warranty.

For 24-hour chemical spill, leak, fire, or accident response information, call CHEMTREC toll free at 1-800-424-9300.

SEED TREATMENT FUNGICIDE

For use with standard mechanical, slurry, or mist-type commercial seed treating equipment only.

Manufactured by:
ALBAUGH, LLC
1525 NE 36th Street,
Ankeny, IA 50021

 **FUNGICIDE
SEED TREATMENT**

EPA Reg. No. 45002-40
AD092024S

 **ALBAUGH[®]**
your alternative

FIRST AID	
IF SWALLOWED: SI SE INGIERE:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
	<ul style="list-style-type: none"> • Llame de inmediato a un centro de control de envenenamientos o a un médico para consejo de tratamiento. • Si la persona puede tragar, haga que beba un vaso de agua lentamente. • NO induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. • NO administre nada por boca a una persona que haya perdido el conocimiento.
HOTLINE NUMBER: For non-emergency questions regarding exposure to this product, call 1-888-347-6732 (7 days/week, 24-hr/day). For medical emergencies, call the poison control center at 1-800-222-1222. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NÚMERO DE LÍNEA DIRECTA: Para información sobre la exposición a este producto que nos sea emergencia, llame al 1-888-347-6732 (7 días/semana, 24-hr). Para obtener información de emergencia sobre este producto, llame al centro de control de intoxicaciones al 1-800-222-1222. Cuando llame a un centro de control de envenenamiento, o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Socks and shoes,
- Chemical-resistant gloves, such as Barrier laminate, Butyl rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, Polyvinyl Chloride (PVC) ≥ 14 mils, Viton ≥ 14 mils

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

DECLARACIONES PREVENTIVAS

RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS

PRECAUCION. Dañino si se ingiere. Lávese minuciosamente con agua y jabón después de manipularlo y antes de comer, beber, mascar chicle, consumir tabaco o ir al baño.

EQUIPO DE PROTECCIÓN PERSONAL (PPE)

Algunos materiales resistentes a este producto químico son enumerados a continuación.

Los aplicadores y otros manipuladores de pesticidas (plaguicidas) deben usar:

- Camisa de manga larga, pantalones largos,
- Calcetines y zapatos,
- Guantes resistentes a productos químicos (como laminado de barrera, caucho de butilo ≥ 14 mils, caucho de nitrilo ≥ 14 mils, caucho de neopreno ≥ 14 mils, cloruro de polivinilo (PVC) ≥ 14 mils, vitón ≥ 14 mils)

Siga las instrucciones del fabricante para la limpieza/mantenimiento del equipo de protección personal (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE separadamente de otra ropa para lavar.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.

As soon as possible, wash thoroughly and change into clean clothing.

RECOMENDACIONES DE SEGURIDAD PARA EL USUARIO

Los usuarios deben usar:

- Lávese minuciosamente con agua y jabón después de manipularlo y antes de comer, beber, mascar chicle, consumir tabaco o ir al baño.
- Quítese la ropa/EPP inmediatamente si el pesticida entra. Luego lávese bien y póngase ropa limpia.
- Quítese el EPP inmediatamente después de manipular este producto. Lave el exterior de los guantes antes de quitárselos.

Tan pronto como sea posible, lávese bien y cámbiese a ropa limpia.

ENVIRONMENTAL HAZARDS

Treated seed exposed on soil surface is hazardous to wildlife. Cover or collect seeds spilled during loading and planting. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. This fungicide may persist in soil and long-term use in the same field may result in an accumulation of active ingredient. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT mix or allow contact with oxidizing agents. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

This product may be applied with standard mechanical, slurry, or mist-type commercial seed treating equipment only, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. **DO NOT** use this product on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

DO NOT apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

INSTRUCCIONES DE USO

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal. Lea atentamente todas las instrucciones de uso antes de aplicar.

Este producto se puede aplicar únicamente con equipos comerciales estándar de tratamiento de semillas mecánicos, de lechada o de nebulización, siempre que el equipo pueda calibrarse para aplicar el producto a las semillas de manera precisa y uniforme. **NO** use este producto en establecimientos agrícolas en cajas con tolva, cajas para macetas, cajas para lechada u otras aplicaciones de tratamiento de semillas durante o inmediatamente antes de la siembra.

NO aplique este producto de manera que entre en contacto con trabajadores u otras personas, ya sea directamente o por deriva. Solo los manipuladores protegidos pueden estar en el área durante la aplicación. Para cualquier requisito específico de su estado o tribu, consulte a la agencia estatal o tribal responsable de la regulación de pesticidas.

PRODUCT USE INFORMATION

This product is a systemic broad-spectrum fungicide seed treatment specifically for protection against a wide variety of soil borne and seed borne diseases as listed in the Application Instructions table for the specific crops. This product does not provide activity against *Pythium* spp. or *Phytophthora* spp. To provide best seedling protection against these specific fungal pathogens, use this product in combination with a seed treatment fungicide containing mefenoxam at the label rate registered for your crop and disease complex. Consult the label for appropriate use rate and follow all use instructions and precautions for both products. For broader spectrum control, this product may be mixed in a water-based slurry with other registered seed treatment insecticides and fungicides, however, **do not** use with other seed treatment products unless previous experience assures compatibility.

Uniform application to seed is necessary to ensure best disease protection and optimum performance. Seed should be sound and well-cured before treatment. Refer to the label rates in **APPLICATION INSTRUCTIONS**. Product requires dilution prior to application. Contact your local Albaugh, LLC representative or supplier for specific recommendations.

NOTE: **This product is undyed.** The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately colored with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, Part 153.155.

SEED BAG LABEL REQUIREMENTS

THE FEDERAL SEED ACT REQUIRES THAT CONTAINERS OF SEEDS TREATED WITH THIS PRODUCT MUST BE LABELED WITH THE FOLLOWING STATEMENTS:

This seed has been treated with ipconazole fungicide.

Do not use treated seed for feed, food, or oil purposes.

THE U.S. ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON CONTAINERS OF SEEDS TREATED WITH THIS PRODUCT:

Store away from feed and foodstuffs.

DO NOT allow children, pets or livestock to have access to treated seeds.

Persons opening this bag or loading/pouring the treated seed must wear a long-sleeved shirt, long pants, shoes, socks and chemical-resistant gloves.

After seeds have been planted, **do not** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. *Exception:* Once seeds are planted and covered by soil or other planting media, the Worker Protection Standard (WPS) allows workers to enter the treated area without restriction.

DO NOT graze or feed livestock on treated areas or hay grown from treated peanut seed.

Dispose of excess treated seed by burying seed away from bodies of water.

Dispose of excess treated seed by burying seed away from bodies of water.

DO NOT contaminate water bodies when disposing of planting equipment washwater.

Dispose of seed packaging in accordance with local requirements.

Excess treated seed may be used for ethanol production only if: (1) By-products are not used for livestock feed, and (2) No measurable residues of pesticides remain in ethanol by-products that are used for agronomic practice.

Treated seed must be incorporated into the soil at the recommended depth.

RESISTANCE MANAGEMENT RECOMMENDATIONS

IPCONAZOLE	GROUP	3	FUNGICIDE
------------	-------	---	-----------

For resistance management, this product contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to this product and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens. Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.

- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance. Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crop and pathogens.
- For further information or to report suspected resistance contact Albaugh, LLC. 1-800-247-8013. You can also contact your pesticide distributor or university extension specialist to report resistance.

CROP SPECIFIC USE INSTRUCTIONS

**FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT
IN CROP INJURY OR POOR DISEASE CONTROL**

WHEAT	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Cochliobolus sativus</i>) Loose Smut (<i>Ustilago tritici</i>) Common bunt (<i>Tilletia caries</i> , <i>T. foetida</i>)	0.051 - 0.085 fl oz / 100 lb seed To deliver 0.0005 - 0.0009 mg ai / seed, apply 0.004 - 0.007 fl oz / 100,000 seeds (Based on average of 13,000 seeds / lb)
¹ Diseases Partially Controlled	
Common Root Rot (<i>Cochliobolus sativus</i>) Crown and Foot Rot (<i>Fusarium</i> spp.)	
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

¹Partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial disease control.

BARLEY	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Cochliobolus sativus</i>) True Loose Smut (<i>Ustilago nuda</i>) Covered Smut (<i>U. hordei</i>) False Loose Smut (<i>U. nigra</i>)	0.051 - 0.085 fl oz / 100 lb seed To deliver 0.0005 - 0.0009 mg ai / seed, apply 0.004 - 0.007 fl oz / 100,000 seeds (Based on average of 13,000 seeds / lb)
¹ Diseases Partially Controlled	
Common Root Rot (<i>Cochliobolus sativus</i>) Crown and Foot Rot (<i>Fusarium</i> spp.) Leaf stripe (<i>Pyrenophora graminea</i>)	
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

¹Partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial disease control.

OATS	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Cochliobolus sativus</i>) Loose Smut (<i>Ustilago avenae</i>) Covered Smut (<i>U. kollerii</i>)	0.051 - 0.085 fl oz / 100 lb seed To deliver 0.0005 - 0.0009 mg ai / seed, apply 0.004 - 0.007 fl oz / 100,000 seeds (Based on average of 13,000 seeds / lb)
¹ Diseases Partially Controlled	
Common Root Rot (<i>Cochliobolus sativus</i>) Crown and Foot Rot (<i>Fusarium</i> spp.)	
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

¹Partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial disease control.

RYE, TRITICALE	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Cochliobolus sativus</i>)	0.051 - 0.085 fl oz / 100 lb seed To deliver 0.0004 - 0.0006 mg ai / seed, apply 0.003 – 0.005 fl oz / 100,000 seeds (Based on average of 18,850 seeds / lb)
¹ Diseases Partially Controlled	
Common Root Rot (<i>Cochliobolus sativus</i>) Crown and Foot Rot (<i>Fusarium</i> spp.)	
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

¹Partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial disease control.

BUCKWHEAT, MILLET	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i>)	0.051 - 0.085 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

CORN (Field, Sweet, Popcorn)	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>)	0.044 - 0.085 fl oz / 100 lb seed For field corn: To deliver 0.0035 - 0.0067 mg ai / seed, apply 0.021 – 0.040 fl oz / 80,000 seeds (Based on average of 1,680 seeds / lb) For sweet corn: To deliver 0.0019 – 0.0036 mg ai /seed, apply 0.011 – 0.022 fl oz / 80,000 seeds (Based on average of 3,150 seeds per lb)
¹Diseases Partially Controlled	
Root Rot (<i>Fusarium</i> spp.)	For popcorn: To deliver 0.0019 - 0.0037 mg ai / seed, apply 0.012 – 0.022 fl oz / 80,000 seeds (Based on average of 3,060 seeds per lb)
Diseases Controlled	Application Rate
Corn head smut (seed- and soil- borne <i>Sphacelotheca reiliana</i>)	0.34 fl oz / 100 lb seed For field corn: To deliver 0.0269 mg ai /seed, Apply 0.16 fl oz / 80,000 seeds (Based on average of 1,680 seeds / lb) For sweet corn: To deliver 0.0144 mg ai / seed, Apply 0.086 fl oz / 80,000 seeds (Based on average of 3,150 seeds / lb) For popcorn: To deliver 0.0148 mg ai / seed, apply 0.089 fl oz / 80,000 seeds (Based on average of 3,060 seeds / lb)
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

¹Partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial disease control.

SORGHUM	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i>)	0.051 - 0.17 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

SOYBEANS	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Phomopsis</i> (<i>Diaporthe</i>) Seed-borne <i>Botrytis</i> and <i>Sclerotinia</i>	0.085 fl oz / 100 lb seed To deliver 0.0040 mg ai / seed, apply 0.043 fl oz / 140,000 seeds (Based on average of 2,800 seeds per lb)
¹ Diseases Partially Controlled	
Root rot caused by <i>Fusarium</i> spp. *Red crown rot caused by <i>Cylindrocladium</i> spp. *Not registered for use in California.	
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

¹Partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial disease control.

BEANS (dried or succulent shelled and edible podded)	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>) Seedling blight (seed-borne <i>Ascochyta</i> spp.) Seed-borne <i>Botrytis</i> and <i>Sclerotinia</i>	0.085 fl oz / 100 lb seed To deliver 0.0071 mg ai per seed, apply 0.053 fl oz per 100,000 seeds (Based on average of 1,600 seeds per lb)
¹ Diseases Partially Controlled	
Root rot (<i>Fusarium</i> spp.) Anthracnose (seed-borne <i>Colletotrichum lindemuthianum</i>)	
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

¹Partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial disease control.

PEAS (dried or succulent shelled and edible podded)	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>) Seedling blight (seed-borne <i>Ascochyta</i> spp.) Seed-borne <i>Botrytis</i> and <i>Sclerotinia</i>	0.085 fl oz / 100 lb seed To deliver 0.0035 mg ai / seed, apply 0.027 fl oz / 100,000 seeds (Based on average of 3,200 seeds per lb)
Diseases Partially Controlled	
Root rot (<i>Fusarium</i> spp.)	
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

¹Partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial disease control.

PEANUTS Not Registered for Use By California	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Cylindrocladium</i> spp.)	0.085 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year. DO NOT graze or feed livestock on treated areas or on hay grown from treated peanut seed.	

ROOT VEGETABLES, EXCEPT POTATOES (seed, tubers, or propagating material) Not Registered for Use By California	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blight (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>)	0.34 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

SUNFLOWER	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>)	0.051 - 0.085 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

LEAFY VEGETABLES (Brassica) Not Registered for Use By California	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>)	0.051 - 0.34 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

LEAFY VEGETABLES (non- Brassica) Not Registered for Use By California	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> and <i>Rhizoctonia solani</i>)	0.051 - 0.34 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

SPINACH Not Registered for Use By California	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>) Seed-borne <i>Verticillium dahliae</i>	0.34 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

CUCURBITS Not Registered for Use By California	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>)	0.051 - 0.34 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

COTTON	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>)	0.051 - 0.34 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

MUSTARD, RAPE, CANOLA Not Registered for Use By California	
Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>) Phoma stem canker / blackleg (<i>Leptosphaeria maculans</i>)	0.051 - 0.34 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

ORNAMENTAL FLOWERS

Not Registered for Use By California

Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>)	0.051 - 0.34 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

CONIFERS

Not Registered for Use By California

Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>)	0.051 - 0.34 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

TURFGRASS

Not Registered for Use By California

Diseases Controlled	Application Rate
Seed rots (seed- and soil-borne <i>Aspergillus</i> spp. and <i>Penicillium</i> spp.) Seed rot, damping off and seedling blights (seed- and / or soil-borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i>)	0.051 - 0.085 fl oz / 100 lb seed
Restrictions: DO NOT apply more than 0.00545 lbs ipconazole per acre per year.	

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office.

CONTAINER HANDLING:

less than or equal to 5 gallons

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

greater than 5 gallons

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

ALMACENAMIENTO Y DESECHO

No contamine agua, comida ni forrajes mediante el almacenamiento y desecho.

ALMACENAMIENTO DE PESTICIDAS: Almacenar únicamente en contenedores originales. Mantenga el recipiente cerrado cuando no esté en uso. No almacenar cerca de alimentos o piensos. En caso de derrame en el piso o superficies pavimentadas, limpie y retire al área de almacenamiento de desechos químicos hasta que se pueda realizar la eliminación adecuada si el producto no se puede reutilizar de acuerdo con la etiqueta.

DESECHO DE PESTICIDAS: Los residuos de pesticidas son ligeramente peligrosos. La eliminación inadecuada del exceso de pesticida, mezcla de aspersión o enjuague es una violación de la ley federal. Si estos desechos no se pueden eliminar según las instrucciones de la etiqueta, comuníquese con su agencia estatal, de control ambiental o de pesticidas, o con el representante de desechos peligrosos en la oficina regional de la EPA más cercana para obtener orientación.

MANIPULACIÓN DE RECIPIENTES:

≤ 5 galones: Recipientes no recargables. No reutilice ni rellene este recipiente. Enjuague tres veces el recipiente (o equivalente) inmediatamente después de vaciarlo. Enjuague tres veces de la siguiente manera: Vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla y drene durante 10 segundos después de que el flujo comience a gotear. Llene 1/4 del recipiente con agua y vuelva a tapar. Agitar durante 10 segundos. Vierta el enjuague en el equipo de aplicación o en un tanque de mezcla o almacene el enjuague para su uso o eliminación posterior. Drene durante 10 segundos después de que el flujo comience a gotear. Repita este procedimiento dos veces más. Luego ofrézcalo para reciclarlo, si está disponible, o perforélo y deséchelo en un vertedero sanitario, o mediante incineración, o mediante otros procedimientos aprobados por las autoridades estatales y locales.

>5 galones: Recipientes no recargables. No reutilice ni rellene este recipiente. Enjuague tres veces el recipiente (o equivalente) inmediatamente después de vaciarlo. Enjuague tres veces de la siguiente manera: Vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla. Llene el recipiente hasta 1/4 de su capacidad con agua. Vuelva a tapar y apretar los cierres. Inclíne el recipiente de lado y gírelo hacia adelante y hacia atrás, asegurando al menos una revolución completa, durante 30 segundos. Coloque el recipiente sobre su extremo e inclínelo hacia adelante y hacia atrás varias veces. Vacíe el enjuague en el equipo de aplicación o en un tanque de mezcla o almacene el enjuague para su uso o eliminación posterior. Repita este procedimiento dos veces más. Luego ofrézcalo para reciclarlo, si está disponible, o perforélo y deséchelo en un vertedero sanitario, o mediante incineración, o mediante otros procedimientos aprobados por las autoridades estatales y locales.

EL RECIPIENTE NO ES SEGURO PARA ALIMENTOS, PIENSOS O AGUA POTABLE.

CONDITIONS OF SALE, WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

ALBAUGH, LLC and the Seller offer this product subject to the following Conditions of Sale, Warranty Disclaimer and Limitation of Liability, and by purchasing and using this product, Buyer and Users accept such Conditions of Sale, Warranty Disclaimer and Limitation of Liability. No employee or agent of ALBAUGH, LLC or the Seller is authorized to modify the terms of these Conditions of Sale, Warranty Disclaimer and Limitation of Liability unless in writing and signed by a duly authorized representative of ALBAUGH, LLC.

The **DIRECTIONS FOR USE** of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH, LLC or the Seller. All such risks shall be assumed by the Buyer. **WITHOUT LIMITATION, THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.**

TO THE FULLEST EXTENT PERMITTED BY APPLICABLE LAW, ALBAUGH, LLC MAKES NO EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR ANY OTHER EXPRESS OR IMPLIED WARRANTY THAT EXTENDS BEYOND THE STATEMENTS MADE ON THIS LABEL. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S AND SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OF HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. **IN NO CASE SHALL ALBAUGH, LLC OR THE SELLER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**