



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

MOVENTO® MPC

For control of Asian citrus psyllid (ACP) on citrus during bloom periods.

EPA Reg. No. 264- 1065

SLN. No. FL-140004

24(c) Special Local Need

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

ACTIVE INGREDIENT:

spirotetramat: *cis*-3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4.5]dec-3-en-4-yl-ethyl carbonate 14.5%

OTHER INGREDIENTS: 85.5%

Total 100.0%

Contains 1.25 pounds of spirotetramat per US gallon, or 150 grams per liter.

DIRECTIONS FOR USE

Apply specified dosage of MOVENTO MPC as needed for control. Treatment must be made when insect populations begin to build and before a damaging population becomes established. Rate selected for use must depend on stage of pest development at application and infestation level of those pests. **MOVENTO MPC must be tank-mixed with a spray adjuvant / additive having spreading and penetrating properties to maximize leaf uptake and systemicity of the active ingredient within treated plants; please contact your local Bayer CropScience representative or PCA for specific recommendations.**

CITRUS FRUITS

Crops of Crop Group 10-10 Including: Australian lime (desert, finger, and round), Brown River finger lime, Calamondin, Citron, Grapefruit, Japanese summer grapefruit, Kumquat, Lemon, Lime, Mediterranean mandarin, Mount White lime, New Guinea wild lime, Orange (sour and sweet), Pummelo, Russell River lime, Satsuma mandarin, Sweet lime, Tachibana orange, Tahiti lime, Tangelo, Tangerine, Tangor, Trifoliate orange, Uniq fruit, including cultivars, varieties and/or hybrids of these commodities.

PEST CONTROLLED	Rate fluid ounces/Acre	Rate lb ai/Acre
Asian citrus psyllid	16.0	0.16

Notes and Restrictions

Pre-Harvest Interval (PHI): **1 day.**

Minimum interval between applications: **21 days.**

Maximum MOVENTO MPC allowed per 12 month period: **32 fluid ounces/Acre (0.32 lb ai/Acre).**

Do not make more than one application during the primary citrus bloom period.

Do not apply non-ionic surfactants in tank mix combination with MOVENTO on white grapefruit.

For MEDICAL And TRANSPORTATION Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

24(c) Registrant:

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

MOVENTO is a registered trademark of Bayer.