

**FIFRA §2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE of  
ALABAMA, FLORIDA, GEORGIA, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA,  
TENNESSEE & VIRGINIA**



**EPA Reg. No. 10163-332  
Fungicide for Control of Listed Diseases in Rapeseed (Canola Varieties ONLY)  
Crop Subgroup 20A**

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

**Rapeseed (Canola Varieties ONLY) Crop Subgroup 20A**

**CROP LIST: (RAPESEED (CANOLA VARIETIES ONLY)): Borage; Canola, Crambe; Cuphea; Echium; Flax Seed; Gold of Pleasure; Hare's Ear Mustard; Lesquerella; Lunaria; Meadowfoam; Milkweed; Mustard Seed; Oil Radish; Poppy Seed; Rapeseed; Sesame; Sweet Rocket Cultivars, Varieties, and/or hybrids of these.**

Disease	Dosage Rate		Application Instructions
	FL OZ/A	GPA	
Cercospora Leaf Spot	10.0 to 19.0 (0.065 lbs. azoxystrobin and 0.052 lbs. tetraconazole) — (0.124 lbs. azoxystrobin and 0.099 lbs. tetraconazole)	Ground minimum: 10  Aerial minimum: 5	Begin applications as a preventative at the beginning of flower between 20% to 50% bloom (21 to 28 days prior to crop maturity (BBCH 89)) and repeat if needed 7 to 14 days after the first application. Under severe disease conditions the shorter spray intervals must be used.

**RESTRICTIONS AND LIMITATIONS**

1. **DO NOT** make more than two (2) applications per year.
2. **DO NOT** apply more than 38 fl oz of AFFIANCE per acre per year.
3. **DO NOT** apply more than 19 fl oz/A (0.124 lbs. azoxystrobin and 0.099 lbs. tetraconazole) per single application.
4. **DO NOT** reapply within 7 days (retreatment interval) of the initial application.
5. **DO NOT** apply more than 0.2 lb ai of a tetraconazole-containing product per acre per year.
6. **DO NOT** apply more than 0.45 lb ai of an azoxystrobin-containing product per acre per year.
7. **DO NOT** apply within 30 days of harvest (30 day PHI).

ID# 23-11-03

AFFIANCE® is a registered trademark of Gowan Company, LLC



Produced For:  
Gowan Company, LLC  
P.O. BOX 5569  
Yuma, AZ 85366-5569