



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)

DELARO® 325 SC FUNGICIDE

EPA Reg. No. 264-1055

For Use On: Potatoes

This supplemental label expires on 09/21/2025 and must not be used or distributed after this date.

Supplemental Label

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read this label and the product package label before using this product. This Supplemental Label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the product label for DELARO 325 SC Fungicide attached to the container.

POTATO

Disease Controlled	Application Rate	Application Instructions
Early blight (<i>Alternaria solani</i>) Brown spot (<i>Alternaria alternata</i>) Botrytis leaf spot (<i>Botrytis cinerea</i>) Black dot (<i>Colletotrichum coccodes</i>) White mold (<i>Sclerotinia sclerotiorum</i>)	8.0 to 11.4 fl oz/acre (0.093 - 0.133 lb/acre prothioconazole) (0.079 - 0.113 lb/acre trifloxystrobin)	Begin fungicide applications preventatively and continue as needed on a 7- to 14-day interval, as long as the maximum annual rate is not exceeded.

Restrictions:

- Maximum single application rate: 11.4 fl oz/acre of DELARO 325 SC Fungicide (0.133 lb/acre prothioconazole and 0.113 lb/acre trifloxystrobin).
- Maximum annual application rate: 22.8 fl oz of DELARO 325 SC Fungicide (0.266 lb/acre prothioconazole and 0.226 lb/acre trifloxystrobin).
- Maximum number of applications: 2 (at 11.4 fl oz/acre).
- Pre-Harvest Interval (PHI): **14 day(s)**
- Minimum interval between applications: **7 day(s)**
- Minimum application volumes: **10 gallons/Acre** (Ground); **5 gallons/Acre** (Aerial)
- Regardless of formulation or method of application, **DO NOT** apply more than 0.267 lbs prothioconazole, or 0.748 lbs trifloxystrobin per acre per year from all uses, including seed treatment, soil and foliar applications.
- **DO NOT** use with handheld application equipment, including mechanically pressurized spray gun, backpack or tank pressurized spray gun or handheld boom applicators.
- To limit the potential for development of disease resistance to this fungicide, **DO NOT** make more than 2 sequential applications of DELARO 325 SC or any Group 11 containing fungicide before rotating with a fungicide from a different Group.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

DELARO is a Registered trademark of Bayer.

Created on 09/21/2022