



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3557

FOR DISTRIBUTION AND USE ONLY IN OREGON

THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS
MADE AS PERMITTED UNDER FIFRA
SECTION 2(ee) AND HAS NOT BEEN
SUBMITTED TO OR APPROVED BY THE US
EPA.

This recommendation for Topguard® Fungicide is valid until April 30, 2027, or until withdrawn, canceled, or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR PREVENTION OF EASTERN FILBERT BLIGHT IN HAZELNUTS VIA DRIP CHEMIGATION

Directions For Use

When applied as directed through drip irrigation and microsprinklers, Topguard® Fungicide will provide control of Eastern filbert blight in hazelnuts.

Ensure adequate soil moisture prior to utilizing Topguard® Fungicide in a drip irrigation system. Do not apply Topguard® Fungicide through drip irrigation deeper than 12" below the soil surface. Discontinue drip irrigation application at fungicide depletion from the main feed supply tank or after 6 hours from start, whichever is shorter. Delay subsequent irrigation (water only) for at least 24 hours following drip application for best results.

Crop	Pest	Rate (fl. oz./A)	Application Directions	Retreatment Interval (Days)
Hazelnuts	Eastern Filbert Blight <i>Anisogramma anomale</i>	14	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary.	7



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

All applicable directions, restrictions, and precautions on the Topguard® Fungicide label (EPA Reg. 279-3557) must be followed. This recommendation should be in the possession of the user at the time of pesticide application.

Restrictions:

- No single application may exceed 14 fl. oz. of product (0.114 lb ai)/A
- Do not apply more than 4 applications per year, including drip chemigation applications
- Do not apply more than 56 fl. oz. of product (0.455 lb ai)/A per year
- Restricted Entry Interval (REI) is 12 hours
- Preharvest interval (PHI): May be applied the day of harvest (0 day PHI)

R-4709 032822



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000