

RESTRICTED USE PESTICIDE

Due to Ground and Surface Water Concerns

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification. This product is a restricted use herbicide due to ground and surface water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.



ACETOCHLOR	GROUP	15	HERBICIDE
ATRAZINE	GROUP	5	HERBICIDE

VOLLEY ATZ NXT®

TENKÖZ

HERBICIDE

A preemergence herbicide for control of annual grasses and broadleaf weeds in field corn, production seed corn, silage corn, sweet corn, popcorn

Active Ingredients:

acetochlor: 2-chloro- N-ethoxymethyl-N-(2-ethyl-6-methylphenyl)acetamide....	33.4%
atrazine: [2-chloro-4-(ethylamino)-6-(isopropylamino)-s-triazine] and related triazines.....	26.9%
Other Ingredients:.....	39.7%
Total	100.0%

Contains 372 grams per liter or 3.1 pounds/gallon acetochlor and 300 grams/liter or 2.5 pounds/gallon atrazine and related components.

Not for Sale, Safe Into, Distribution and/or Use in Nassau and Suffolk Counties of New York State.

Keep Out of Reach of Children CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including First Aid and Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use and

Limitation of Remedies at end of label booklet. If terms are not acceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-671-55467

EPA Est. 11773-IA-001

294592 2505

CD02-277-020



Scan for safety information in Spanish.
Escanee para la información de seguridad en español.

® Trademarks of Corteva Agriscience and its affiliated companies.

Produced for

Tenköz, Inc

1725 Windward Concourse, Suite 410

Alpharetta, GA 30005

NET CONTENTS 2.5 GAL

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful If Swallowed or Inhaled • Causes Moderate Eye Irritation • May Cause Allergic Skin Reaction

Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

Call a poison control center or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

Call a poison control center or doctor for treatment advice. Sensitized persons should avoid further contact and reuse of contaminated clothing.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 day or night, for emergency treatment information.

Personal Protective Equipment (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as polyethylene or polyvinylchloride
- Shoes plus socks
- Chemical-resistant apron when mixing/loading, cleaning up spills, or cleaning equipment, or otherwise exposed to the concentrate

See Engineering Controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls: When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(5)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Non-Target Organism Advisory

This product is toxic to plants, fish and aquatic invertebrates and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of acetochlor from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours

Groundwater Advisory

Acetochlor is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e., well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

Refer to Use Restrictions and Tile-Outletted Terraced Fields sections for additional specific information.

Directions for Use

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any

requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for use in the states of Hawaii or Alaska, or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands, and the North Mariana Islands)

Endangered Species:

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/essp/>, call 1-844-447-3813, or email ESSP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product.

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Atrazine Watershed Information Center (AWIC) to determine whether the use of this product is prohibited in your watershed. AWIC can be accessed through www.atrazine-watershed.info or 1-866-365-3014. If use of this product is prohibited in your watershed, you may return unopened product to your point of purchase or contact Tenkoz for a refund.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with vermiculite, earth, or synthetic absorbent.

Storage and Disposal (Cont.)

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Product Information

For use only on field corn, production seed corn, silage corn, sweet corn, and popcorn. Corn in this label refers to: field corn, production seed corn, silage corn, sweet corn and popcorn.

Volley® ATZ NXT herbicide may be applied to the surface or incorporated into the top 1-2 inch layer of soil. It may be used for control alone, or in tank mix combinations, for the weeds listed in the "Target Weeds" section of these use directions. Volley ATZ NXT controls weeds by interfering with normal germination and seedling development. Volley ATZ NXT does not control established or emerged weeds present at application.

Use Restrictions

- Not for use in the states of Hawaii or Alaska, or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands, and the North Mariana Islands)
- This product is restricted to impregnation of dry bulk fertilizer to 340 tons per worker per day for no more than 30 days per calendar year for use on corn.
- **Not for Sale, Sale Into, Distribution and/or Use in Nassau and Suffolk Counties of New York State.**
- This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the ground water is shallow, may result in ground water contamination.
- Do not apply Volley ATZ NXT before pre-irrigation in irrigated areas.


- Do not allow Volley ATZ NXT to contaminate feed or food.
- Do not contaminate irrigation water used for crops other than corn or water used for domestic purposes.
- Do not use Volley ATZ NXT on any crop other than field corn, production seed corn, silage corn, sweet corn, and popcorn.
- Use on Miscanthus and other non-food perennial bioenergy crops is prohibited.
- Do not apply Volley ATZ NXT using mechanically pressurized handgun to sweet corn.
- On the following soil types, do not apply this product within 50 feet of any well where the depth to groundwater is 30 feet or less: sands with less than 3% organic matter; loamy sands with less than 2% organic matter; or sandy loams with less than 1 percent organic matter. See the figure for additional clarification.

- This product must not be mixed or loaded within 50 ft. of intermittent streams and rivers, natural or impounded lakes and reservoirs. This product must not be applied by ground within 66 feet of the points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet around natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66-foot buffer or setback from runoff entry points must be planted to crop, seeded with grass or other suitable crop.
- This product must not be mixed or loaded, or used within 50 feet of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. Additional State imposed requirements regarding well-head setbacks and operational area containment must be observed.

• Tile-Outletted Fields Containing Standpipes

To ensure protection of surface water from runoff through standpipes with tile-outlets in fields, one of the following restrictions must be used in applying this product to tile-outletted fields containing standpipes:

1. Do not apply this product within 66 feet of standpipes in tile-outletted fields.
 2. Apply this product to the entire tile-outletted field and immediately incorporate it to a depth of 2-3 inches in the entire field.
 3. Apply this product to the entire tile-outletted field under a no-till practice only when high crop residue management practices are used. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during or after crop harvest.
- Do not apply Volley ATZ NXT postemergence to sweet corn.
 - **Chemigation:** Do not apply this product through any type of irrigation system, unless otherwise directed by approved supplemental labeling in possession of the user at the time of application.
 - Do not use flood irrigation to apply or incorporate this product.
 - Product must be used in a manner that will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.



- Do not use tailwater from the first flood or furrow irrigation of treated fields to treat non-target crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.
- **Aerial Application:** Do not apply this product using aerial application equipment unless otherwise directed by approved supplemental labeling in possession of the user at the time of application.
- **Maximum Atrazine Application Rates Per Calendar Year:** Maximum annual atrazine broadcast application rates for corn must be as follows:
 - If no atrazine was applied prior to corn emergence, apply a maximum of 2.0 pounds active ingredient (3.2 quarts Volley ATZ NXT) per acre. If postemergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.5 pounds active ingredient per acre per calendar year. Note: One quart per acre Volley ATZ NXT delivers 0.625 pound active ingredient atrazine per acre.
 - Apply a maximum of 2.0 pounds active ingredient (3.2 quarts Volley ATZ NXT) per acre if a single preemergence application is made on soils that are not highly erodible or on highly erodible soil if at least 30% of the soil is covered with plant residues.
 - Apply a maximum of 1.6 pounds active ingredient (2.5 quarts Volley ATZ NXT) per acre as a single preemergence application on highly erodible soils if less than 30% of the soil is covered with plant residues; or 2.0 pounds active ingredient (3.2 quarts Volley ATZ NXT) per acre if only applied postemergence.
- **Maximum Acetochlor Application Rates Per Calendar Year:** Maximum annual acetochlor broadcast application rates for corn must not exceed 3.0 pounds active ingredient (3.8 quarts Volley ATZ NXT) per acre. Note: One quart per acre Volley ATZ NXT delivers 0.77 pound active ingredient acetochlor per acre.
- **Preharvest Interval:** Do not apply Volley ATZ NXT within 60 days of harvest of field corn for field corn forage uses or 45 days for sweet corn forage uses.
- Postemergence applications of Volley ATZ NXT to corn must be made before the crop reaches 11 inches in height.
- Volley ATZ NXT should not be stored near seeds, fertilizers, or foodstuffs.
- All containers of Volley ATZ NXT should be kept tightly closed when not in use.
- Applied according to directions and under normal growing conditions, Volley ATZ NXT will not harm the treated crop. During germination and early stages of growth, extended periods of unusually cold and wet or hot and dry weather, insect or plant disease attack, carryover pesticide residues, the use of certain soil applied systemic insecticides, improperly placed fertilizers or soil insecticides may create abnormal conditions that weaken crop seedlings. Volley ATZ NXT used under these abnormal conditions could result in crop injury.

Rotational Crop Restrictions:

When tank mixing Volley ATZ NXT with other herbicides, follow the most restrictive crop rotation guidelines on the label of each product used.

1. If a crop treated with this product is lost, field corn, seed corn, silage corn, popcorn or sweet corn may be replanted immediately. Do not exceed a total of 3.0 pounds per acre of acetochlor if, additional product is applied.
2. If applied after June 10, do not rotate to crops other than corn or sorghum the next year, or crop injury may occur.
3. Rotate the next season to the following crops: corn (all types), cotton, sorghum or soybeans. Injury from atrazine may occur to soybeans planted the year following application on soils having a calcareous subsurface layer.
4. In the High Plains and Intermountain regions of the West where rainfall is sparse and erratic or irrigation is required, use only when corn or sorghum is to follow corn.
5. In Eastern parts of the Dakotas, Kansas, western Minnesota and Nebraska, do not rotate to soybeans if the rate applied to corn was more than 2.0 pounds active ingredient equivalent of atrazine or soybean injury may occur.
6. Do not plant sugar beets, sunflower, potatoes, tobacco, dry beans or peas, spring-seeded small grains or small-seeded legumes the year following application, or injury from atrazine may occur.

Rotation to Non-food Winter Cover Crops

Following harvest of food crops treated with Volley ATZ NXT, only non-food or non-feed winter cover crops (with the exception of wheat) may be planted. Do not graze or harvest rotational cover crops for food or animal feed for 18 months following the last application of Volley ATZ NXT. This prohibition does not apply to wheat, which may be planted 4 months following the last application of Volley ATZ NXT, or to nongrass animal feeds, which may be planted 9 months after the last application of Volley ATZ NXT.

Aerial Application

Do not apply this product using aerial application equipment unless otherwise directed by approved supplemental labeling in possession of the user at the time of application.

Use Precautions

- Failure to strictly follow label directions may result in exceeding the maximum annual atrazine use rates as stipulated by the Environmental Protection Agency.
- **Note:** This product contains atrazine and thus may not control weeds that are known or suspected to be triazine resistant. Following many years of continuous use of atrazine and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by atrazine and related herbicides. Where this is known or suspected and weeds controlled by atrazine are expected to be present along with resistant biotypes, it is recommended that atrazine be used in combinations or in sequence with other registered herbicides which are not triazines. If only resistant biotypes are expected to be present, use a registered non-triazine herbicide.
- Volley ATZ NXT should not be used on corn seed stock such as Breeders, Foundation, or Increase.

Mandatory Spray Drift

Ground Boom Applications:

- Applicators are required to select a nozzle and pressure that deliver coarse or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.
- User must maintain a 15 foot (4.6 m) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.

Spray Drift Advisories

- **THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.**
- **BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.**
- **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.
- **BOOM HEIGHT - Ground Boom**
Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

• SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

• TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

• TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft

smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

• WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Weed Resistance Management

Volley ATZ NXT contains the active ingredients Acetochlor (Group 15) and Atrazine (Group 5) herbicides, based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before and after application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- If using post-emergence herbicides or tank mixes, control weeds early when they are relatively small.
- Apply full rates of Volley ATZ NXT for the most difficult to control weed in the field at the specified time to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your local company representative, local retailer, or county extension agent.
- Contact your local company representative, crop advisor, or extension agent to find out if suspected resistant weeds to these MOAs have been found in your region. **DO NOT** assume that each listed weed is being controlled by multiple modes of action. Products with multiple active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredients in this product.
- If resistance is suspected, treat weed escapes with an herbicide having a mode of action other than Group 15 or Group 5 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;

- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum herbicide with another mode of action as a foundation in a weed control program, if appropriate.
- Utilize sequential applications of herbicides with alternative modes of action.
- Rotate the use of this product with non-Group 15 and 5 herbicides.
- Avoid making more than two sequential applications of Volley ATZ NXT and any other Group 15 or Group 5 herbicides within a single growing season unless mixed with an herbicide with a different mode of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, for example, mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields to reduce weed seed production.

General principles of herbicide resistance management:

1. Apply integrated weed management practices. Use multiple herbicide modes-of-action with overlapping weed spectrums in rotation, sequences, or mixtures.
2. Use the full recommended herbicide rate and proper application timing for the hardest to control weed species present in the field.
3. Scout fields after herbicide application to ensure control has been achieved. Avoid allowing weeds to reproduce by seed or to proliferate vegetatively.
4. Monitor site and clean equipment between sites.

For annual cropping situations also consider the following:

- Start with a clean field and control weeds early by using a burndown herbicide treatment or tillage in combination with a soil-applied residual herbicide as appropriate.
- Use cultural practices such as cultivation and crop rotation, where appropriate.
- Use good agronomic principles that enhance crop competitiveness.
- Use new commercial seed that is as free of weed seed as possible.

Report any incidence of repeated non-performance of this product on a particular weed to your Tenkoz representative, local retailer, or county extension agent.

Application Directions - Corn

Carriers

Liquids: Either water or liquid fertilizers such as solutions, slurries or suspensions may be used as liquid carriers. If fluid fertilizers are used, a physical compatibility test with these must be done **before combining** in the spray tank. See Appendix I for details

of the compatibility testing procedure. Even if Volley ATZ NXT is physically compatible with a fluid fertilizer, constant agitation is necessary to maintain a uniform mixture during application.

Dry Bulk Fertilizer: Volley ATZ NXT may be impregnated on dry bulk fertilizer and applied as the fertilizer is spread. See Appendix II for directions and restrictions including which fertilizers are compatible.

- This product is restricted to impregnation of dry bulk fertilizer to 340 tons per worker per day for no more than 30 days per calendar year for use on corn.

Adding to Spray Tank

The spray tank must be clean, thoroughly rinsed and decontaminated before adding either Volley ATZ NXT alone or with tank mix combinations. If water is used as the carrier, use clean water.

Used Alone: When Volley ATZ NXT is used alone, add the specified amount to the spray tank when the tank is half filled with carrier, then add the rest of the water or fluid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

Tank Mixed: If a tank mixture is used, it is recommended that a compatibility test be done before actual tank mixing. See Appendix I for details on the procedure for such a test.

Once compatibility is confirmed for the tank mix, fill the tank half full of carrier. Start and continue agitation throughout mixing. All return lines to the spray tank must discharge below the liquid level. Add components in the following order of formulation:

- If a wettable powder or dry flowable formulation is used, make a slurry with water and add it slowly through the screen into the tank. Agitate during the procedure.
- If a flowable formulation is used, add slowly through screen into the tank. Mixing and compatibility may be improved when the flowable is diluted with water before adding to the tank.
- Add Volley ATZ NXT next.
- Add ammonium sulfate then Durango® DMA® herbicide, 2,4-D herbicide, and a nonionic surfactant last, if needed.
- Complete filling the sprayer tank and continue agitation.
- Batches should be mixed and applied the same day.

Note: For all tank mixtures, maintain agitation during mixing and throughout application to ensure spray mixture remains uniformly suspended. If spray mixture is allowed to settle at any time, thorough agitation is required to resuspend the mixture before spraying is resumed.

Volume

Liquid: Use a minimum of 10 gallons per acre in broadcast boom equipment for ground applications.

Dry Bulk Fertilizer: Use a minimum of 200 pounds of dry bulk fertilizer per acre. See Appendix II for directions and restrictions.

Pressure

If liquid carriers are used, the pressure at the nozzle should be 15 to 40 psi to ensure good distribution in the spray pattern. Use appropriate nozzles and 50-mesh or coarser screens, if

needed. Maintain sufficient agitation to ensure the mixture is suspended in the spray tank.

Application Timing and Methods

For the optimum period of effective weed control during the time most critical to corn production, preplant applications of Volley ATZ NXT should occur as close as possible to planting. Preemergence applications should occur as close as possible to planting, but prior to weed emergence; this product will not control emerged weeds present at application.

Early Preplant Surface: On medium and fine textured soils (see Table 1), Volley ATZ NXT may be applied up to 45 days prior to planting field corn or silage corn. Split applications can be made 30 to 45 days prior to planting with 60 percent of the specified broadcast rate applied initially and the remaining 40 percent applied at planting. Applications made less than 30 days prior to planting can be made either as a split or as a single application. If weeds are present at the time of application, apply this product in a tank mixture with an appropriate contact herbicide. Observe directions for use, precautions, and restrictions on the label of the contact herbicide. During the planting operation, be careful not to move untreated soil to the surface or move treated soil out of the row, as weed control may be reduced.

Preplant Incorporation: Volley ATZ NXT and certain tank mixes may be mechanically incorporated in the top 2 inches of the soil with field cultivators, discs, or spring tooth harrows at any time within 14 days prior to planting. Improper incorporation, excessive crop residues, or poor soil tilth may result in erratic, streaked or otherwise unsatisfactory weed control. Do not mix Volley ATZ NXT deeper than 2 inches into the soil and avoid moving or shaping soil after incorporation, as weed control may be reduced.

Preemergence Surface: Volley ATZ NXT and certain tank mixes may be applied to the soil surface as a broadcast or banded application. Precipitation or sprinkler irrigation of at least 0.25 inch is required to bring Volley ATZ NXT into contact with germinating seeds. If rain or sprinkler irrigation does not occur within 7 days after application, weed control may be improved by using a rotary hoe, or similar device, to incorporate the herbicide. The device used should be run at a shallow depth to prevent disturbing the corn seed. Do not remove Volley ATZ NXT from the weed control zone or dilute it with untreated soil. Erratic weed control resulting from exposure of untreated soil may occur if surface soil is moved or reshaped during incorporation.

Postplant-Preemergence: Volley ATZ NXT may be applied immediately after planting but prior to corn emergence. If rain or sprinkler irrigation does not occur within 7 days after application, weed control may be improved by using a rotary hoe, or similar device, to shallowly incorporate the herbicide. The device used should be run at a shallow depth to prevent disturbing the corn seed. Do not remove Volley ATZ NXT from the weed control zone or dilute it with untreated soil. Erratic weed control resulting from exposure of untreated soil may occur if surface soil is moved or reshaped during incorporation.

Banding-Preemergence: Volley ATZ NXT may be applied in a 10- to 14-inch band after corn planting but prior to corn emergence. If rain or sprinkler irrigation does not occur within 7 days after application, weed control may be improved by using a rotary hoe or similar device to incorporate the herbicide. The device used should be run at a shallow depth to prevent disturbing the corn seed. Do not remove Volley ATZ NXT from the weed control zone or dilute it with untreated soil. Erratic weed control resulting from exposure of untreated soil may occur if surface soil is moved or reshaped during incorporation.

Early Postemergence: Volley ATZ NXT may be applied early postemergence to corn up to 11" tall. Applications must be made prior to weed seedling emergence or in a tank mixture with a herbicide that controls the emerged weeds. Read and follow restrictions and directions on tank mix product labels.

Note: Do not make postemergence applications using sprayable liquid fertilizer as the carrier because severe crop injury may occur.

Note: Do not apply Volley ATZ NXT postemergence to sweet corn.

Sprinkler Irrigation: Do not apply Volley ATZ NXT through sprinkler irrigation systems, unless otherwise directed by approved supplemental labeling in possession of the user at the time of application. A sprinkler system may be used to incorporate Volley ATZ NXT after application. After Volley ATZ NXT has been applied, a sprinkler irrigation system set to deliver 0.25 to 0.75 inch of water per acre may be used to incorporate the product. Using more than 0.75 inch of water could result in reduced performance. On sandy soils low in organic matter, use no more than 0.5 inch of water. Do not use flood irrigation to apply or incorporate Volley ATZ NXT.

Planting

Planting should be done as close to the time of application of Volley ATZ NXT as possible. This allows Volley ATZ NXT to provide effective weed control during the time it is most critical in the production of corn.

Cultivation

Cultivation should be delayed as long as possible. If weeds emerge, a shallow cultivation or rotary hoeing will generally result in improved weed control. If Volley ATZ NXT was incorporated, cultivate to a depth of less than half the depth of incorporation.

If cultivation is necessary due to soil crusting, compaction, or escaped weeds, adjust equipment to run shallow and minimize soil movement. This will decrease the possibility of diluting or moving the herbicide from the weed control zone.

Soil Texture and Organic Matter

The specified use rate of Volley ATZ NXT is determined by a combination of two factors, soil texture and organic matter, which must be determined prior to application. Soils are grouped into three textural classes (coarse, medium and fine), as outlined in Table 1.

Table 1: Soil Texture Groupings for Volley ATZ NXT Use Rate Selection

Coarse	Medium	Fine
Sand Loamy Sand Sandy Loam	Loam Silt Loam Silt Sandy Clay Loam	Silty Clay Loam Clay Loam Sandy Clay Silty Clay Clay

Use Rates for Conventional Tillage Systems

Table 2: Volley ATZ NXT Use Rates by Soil Texture and Organic Matter Content in Conventional Tillage Systems

The following use rates are for preplant incorporated, preemergence, and early postemergence applications (see Application Timing and Methods). Apply this product before weeds reach the 2-leaf stage and the corn is no more than 11 inches in height. Consult Table 3 if reduced- or no-till applications are made or application is made more than 14 days prior to planting under conventional tillage.

Soil Texture	Broadcast Rate Per Acre (Quarts)*	
	Soil Organic Matter Content	
	Less than 3%	3% or Greater
Coarse	1.4	1.7
Medium	1.7 – 2.4	2.3 – 2.6†
Fine	2.3 – 2.6†	2.3 – 3.0†

*Use the higher rate in the rate range in areas of heavy weed infestation.

NOTE: In areas of heavy weed infestations, use up to 2.3 quarts per acre on coarse-textured soils and 2.3 – 3.0 quarts per acre on medium- and fine-textured soils.

†On highly erodible soils with less than 30% plant residue, do not apply more than 2.5 quarts of Volley ATZ NXT per acre in a single preemergence application.

Use Rates for Reduced Tillage Systems or Early Preplant Applications in Conventional Tillage Systems

Volley ATZ NXT may be used in reduced- and no-till systems and in early preplant applications in conventional tillage systems. Single applications may be made up to 30 days prior to planting or after planting but before crop emergence. Optimal weed control will be obtained when applications are made as close to planting as possible but before crop emergence. If weeds are emerged at time of application, apply a labeled burndown herbicide such as Durango DMA, Gramoxone, or 2,4-D with Volley ATZ NXT.

Table 3: Volley ATZ NXT Use Rates by Soil Texture in Reduced and No-till Systems or Conventional Tillage Systems when Applications are made more than 14 days Prior to Planting

Soil Texture	Broadcast Rate Per Acre (Quarts)*
Coarse**	1.7 – 2.3
Medium	2.3 – 3.0†
Fine	2.3 – 3.0†

*Use the higher rate in the rate range in areas of heavy weed infestation. Rates are for single applications. Split applications may be used; apply 60% of the specified rate up to 45 days before planting and the remaining 40% at or immediately following planting but before crop emergence.

**Do not apply more than 14 days prior to planting on coarse textured soils.

† On highly erodible soils with less than 30% plant residue, do not apply more than 2.5 quarts of Volley ATZ NXT per acre in a single preemergence application.

This product may also be tank mixed with 1.0-1.25 quarts per acre of Princep 4L (1.1-1.4 pounds per acre of Princep Caliper 90) to provide improved control of crabgrass and fall panicum.

Band Applications

For band applications, use row and band width measurements (inches) to calculate the amount of Volley ATZ NXT to be applied per acre as follows:

$$\begin{array}{l} \text{Band width} \\ \text{in inches} \end{array} \times \begin{array}{l} \text{Rate} \\ \text{per acre for a} \\ \text{broadcast} \\ \text{treatment} \end{array} = \begin{array}{l} \text{Amount of} \\ \text{Volley ATZ NXT} \\ \text{to apply} \\ \text{per acre} \end{array}$$

Weeds Controlled

Volley ATZ NXT applied as directed in this label will provide control or partial control of the weeds listed in Table 4. Additional weeds may be controlled with tank mixes. See the "Tank Mix Combinations" section of this label for tank mix directions. Always consult the tank mix product labels for specific use rates and directions. Always follow the most restrictive label when tank mixing Volley ATZ NXT with another product. Volley ATZ NXT may be tank mixed with any other registered corn product as long as compatibility is verified and tank mixing is not prohibited by the tank mix product label. Note: This product contains atrazine and may not control weeds that are known or suspected to be triazine resistant.

Table 4: Weeds Controlled or Partially Controlled by Volley ATZ NXT at Specified Use Rates

Grasses and Sedges	C = Control PC = Partial Control	Broadleaves	C = Control PC = Partial Control
barnyardgrass	C	beggarweed, Florida	C
crabgrass spp.	C	carpetweed	C
crowfootgrass	C	cocklebur (2)	C
cupgrass, southwestern	C	galinsoga	C
cupgrass, woolly	PC	groundcherry, annual	C
foxtail, giant	C	groundcherry, cutleaf	C
foxtail, green	C	henbit	C
foxtail, robust (purple, white)	C	jimsonweed	C
foxtail, yellow	C	kochia	C
goosegrass	C	lambsquarters, common	C
johnsongrass, seedling	PC	morningglory spp. (2)	C
millet, foxtail	C	mustard spp.	C
millet, wild proso	PC	nightshade, black	C
nutsedge, yellow (1,2)	C	nightshade, hairy	C
oat, wild	C	pigweed, spp.	C
panicum, browntop	C	purslane, common	C
panicum, fall	C	pusley, Florida	C
panicum, Texas (3)	PC	ragweed, common	C
rice, red	C	ragweed, giant	PC
sandbur, field	PC	sicklepod	PC
shattercane	PC	sida, prickly	C
signalgrass, broadleaf (3)	C	smartweed spp.	C
sprangletop, red	C	sunflower, common	PC
wheat, volunteer	C	velvetleaf (2)	C
witchgrass	C	waterhemp, tall	C

(1) Preplant incorporate for improved control.

(2) Use the higher rate in the specified application rate range. Control of these weeds can be erratic, especially under dry conditions. Additional atrazine and/or sequential herbicide applications may be needed for complete control.

(3) Best control is achieved when Volley ATZ NXT is applied within 5 days of planting and rainfall occurs shortly after application or mechanical incorporation is used to activate the herbicide. If rainfall does not occur within 7 days after application, shallow cultivation will enhance activity. Excessive rainfall after application may reduce control. Under adverse weather conditions and/or heavy infestations, a cultivation or follow-up herbicide application may be needed.

Volley ATZ NXT Tank Mix Combinations

When tank mixing or sequentially applying atrazine or simazine or products containing either a.i. to corn, the total pounds of simazine and/or atrazine applied (lb ai/A) must not exceed 2.5 pounds of active ingredient per year.

For all applications, do not exceed the maximum rate of acetochlor as specified in the Maximum Acetochlor Application Rates per Calendar Year section of this label.

Use of Spray Adjuvants

Volley ATZ NXT is a preemergence herbicide for which spray adjuvants have little or no effect on performance. However, several herbicides used in tank mixtures with Volley ATZ NXT require use of adjuvants to aid in the burndown of emerged weeds. Use only those adjuvants specified on tank mix product labels and approved for agricultural crop use. Adjuvants and/or low rate liquid fertilizers (28%, 30% or 32% UAN) or ammonium sulfate (AMS) may be used with tank mixes applied preplant or preemergence to the crop. NOTE: Do not use liquid fertilizer as

the carrier when Volley ATZ NXT is applied postemergence to corn as severe injury may result. The addition of liquid fertilizers used as adjuvants with Volley ATZ NXT tank mixes applied postemergence to corn under environmental stress conditions may result in significant crop injury and should be avoided if the risk of crop injury is unacceptable.

Preemergence Tank Mix Combinations

Tank mix combinations may be used in either conventional, reduced or no-till systems and be applied by the same methods and at the same timings as Volley ATZ NXT unless otherwise specified in the tank mix product label.

Volley ATZ NXT may be tank-mixed with Aim EC, atrazine, Balance Pro, Balance Flexx, Banvel, Callisto, Callisto Xtra, Clarity, Distinct, Durango DMA, Hornet WDG, Linex 4L, Lorox DF, Marksman, Princep, Python WDG, Resource, or 2,4-D for preemergence use in corn. Ensure that specific product being used in the tank mixture is registered for preemergence application to corn. Read and follow label directions of all products in the tank mixture; the most restrictive label directions apply.

When tank mixing Volley ATZ NXT with atrazine, do not exceed the maximum allowable rate of atrazine in your county or state. In some atrazine management areas, atrazine is more restricted. Consult your county extension office or state university for further information.

Conventional Tillage Corn (Volley ATZ NXT plus):

Tank Mix Herbicide †	Comments
Atrazine 4L	<ul style="list-style-type: none"> • Preplant surface, preplant incorporated, preemergence. If emerged weeds are greater than 1.5 inches tall at the time of application, add an appropriate postemergence herbicide. • Longer growing season areas • High rainfall areas • Heavy broadleaf weed pressure
Balance Pro	<ul style="list-style-type: none"> • Not labeled in all states; refer to the Balance Pro label for precautionary statements, use directions, and geographic and other restrictions. • For use in field corn only. • Refer to the use rates section of the Volley ATZ NXT label for minimum use rates.
Hornet WDG	<ul style="list-style-type: none"> • Tank mixing 3.0 – 4.0 oz/acre Hornet® WDG herbicide provides consistent control of velvetleaf, lambsquarters, pigweed species, waterhemp and triazine resistant varieties of these species. Will also provide improved control of cocklebur, common ragweed, giant ragweed, common sunflower and jimsonweed.
Princep 4L	<ul style="list-style-type: none"> • Provides improved crabgrass or fall panicum control.
Python WDG	<ul style="list-style-type: none"> • Tank mixing 0.8 – 1.0 oz/acre Python® WDG herbicide provides consistent control of velvetleaf, lambsquarters, pigweed species, waterhemp and triazine resistant varieties of these species.
Surpass NXT	<ul style="list-style-type: none"> • Tank mix 1 pt/acre for enhanced grass and nutsedge control.

† Formulations that are not listed may be used: Perform a compatibility test and check the label of the tank mix product label for application rates, applicable use directions, precautions and limitations.

Reduced or No-Tillage Corn (Volley ATZ NXT plus):

Tank Mix Herbicide †	Comments
Atrazine 4L	<ul style="list-style-type: none"> • Longer growing season areas • High rainfall areas • Heavy broadleaf weed pressure • If emerged weeds are greater than 1.5 inches tall at the time of application, add an appropriate postemergence herbicide.
Banvel/Clarity Marksman	<ul style="list-style-type: none"> • Apply preplant or preemergence in reduced/no-till systems for burndown of existing weeds. • Preemergence on all soils; medium and fine textured with >2% OM
Durango DMA, Roundup WeatherMax, Touchdown	<ul style="list-style-type: none"> • Burndown existing weeds
Gramoxone Inteon	<ul style="list-style-type: none"> • Control annuals, suppress perennials
Pendimax® / Prowl	<ul style="list-style-type: none"> • Preemergence to early postemergence (up to 3" tall corn) but before weeds are more than 1" tall

Postemergence Tank Mixes (Volley ATZ NXT plus): (Cont.)

Tank Mix Herbicide †	Comments
Princep 4L	<ul style="list-style-type: none"> Provides improved crabgrass or fall panicum control
Surpass NXT	<ul style="list-style-type: none"> Enhanced grass and nutsedge control
2,4-D	<ul style="list-style-type: none"> Burndown existing weeds

† Formulations that are not listed may be used: Perform a compatibility test and check the product label for application rates, applicable use directions, precautions and limitations.

Volley ATZ NXT plus Burndown Herbicide Tank Mixtures

In reduced or no-tillage corn, Durango DMA, Roundup WeatherMax, Gramoxone Inteon, Touchdown, and/or 2,4-D can be tank mixed with Volley ATZ NXT to burn down existing weeds. Burndown herbicides should be applied to emerged weeds when they are small; weeds less than 6 inches in height are easiest to control. Consult the burndown product labels for further information on weeds controlled.

Postemergence Tank Mix Combinations

Volley ATZ NXT may be applied before, with, or following the use of one or more of the following herbicides for postemergence use in corn: Accent, Aim EC, atrazine, Balance Flexx, Banvel, Basis, Basis Gold, Beacon, Butcrl, Butcrl/atrazine, Callisto, Callisto Xtra, Capreno, Clarity, Distinct, Hornet WDG, Impact, Laudis, Liberty, Lightning, Linex 4L, Lorox DF, Marksman, Peak, Permit, Prowl, Pendimox, Pursuit, Resource, Shotgun, Spirit, Status, Steadfast, or 2,4-D. Refer to the tank mix product

label(s) regarding use directions, precautions and restrictions, and the list of weeds controlled. Volley ATZ NXT may be tank mixed with any product approved for use on corn unless it is prohibited on the tank mix product label. Ensure that specific product being used in the tank mixture is registered for postemergence application to corn. Read and follow label directions of all products in the tank mixture; the most restrictive label directions apply. **Note:** DO NOT use liquid fertilizer as the carrier when Volley ATZ NXT is applied postemergence to corn as severe injury may result. The addition of liquid fertilizers used as adjuvants with Volley ATZ NXT tank mixes applied postemergence to corn under environmental stress conditions may result in significant crop injury and should be avoided if the risk of crop injury is unacceptable.

When tank mixing, refer to the tank mix product label and follow the additional use directions given in this table. Volley ATZ NXT can be applied to corn up to 11" tall.

Postemergence Tank Mixes (Volley ATZ NXT plus):

Tank Mix Herbicide	Rate	Comments
Hornet WDG	2-5 oz/acre	<ul style="list-style-type: none"> Always add NIS at 0.25% v/v or COC at 1% v/v.
Aim herbicide	0.3 oz/acre	<ul style="list-style-type: none"> Always add NIS at 0.25% v/v.
Banvel Clarity Marksman	0.5 - 1.0 pt/acre 0.5 - 1.0 pt/acre 2 - 3.5 pt/acre	<ul style="list-style-type: none"> Early postemergence up to 8" tall corn on all soils. If grasses are more than 2- leaf stage, combine with another herbicide to control these weeds. The maximum atrazine application rate for corn is 2.5 pounds active ingredient per calendar year or 2.0 pounds atrazine active ingredient for postemergence application if no atrazine was applied preemergence.
Butcrl Butcrl/atrazine Shotgun herbicide	1.5 pt/acre 2.0 pt/acre 2 - 3 pt/acre	<ul style="list-style-type: none"> Refer to product label for use directions. Refer to label for Shotgun herbicide for timing and use directions.
Atrazine	0.5 - 2.0 lb ai/acre	<ul style="list-style-type: none"> Preplant surface, preplant incorporated, preemergence or early postemergence (up to 8" tall corn). If emerged weeds are greater than 1.5 inches tall at the time of application, add an appropriate postemergence herbicide. Note: The maximum atrazine application rate for corn is 2.5 pounds atrazine active ingredient per acre per calendar year.
Distinct	4.0 - 6.0 oz/acre	<ul style="list-style-type: none"> Always add NIS at 0.25% v/v and 1.25% UAN. Can be applied up to 10-inch corn.
Liberty	16 - 28 oz/acre	<ul style="list-style-type: none"> For use on Liberty tolerant corn only. Apply to grass and broadleaves up to 6" tall. Do not add additional surfactant.
Lightning	1.28 oz/acre	<ul style="list-style-type: none"> For use on Clearfield corn only. Use NIS at 25% v/v and a liquid nitrogen fertilizer at 1 - 2 qt per acre or ammonium sulfate at 2.5 lb per acre.
Pendimox / Prowl	1.8 - 3.6 pt/acre	<ul style="list-style-type: none"> Preemergence to early postemergence (up to 3" tall corn) but before weeds are more than 1" tall.

Postemergence Tank Mixes (Volley ATZ NXT plus): (Cont.)

Tank Mix Herbicide	Rate	Comments												
Pursuit 2.5L Pursuit 70DG	4.0 fl oz/acre 1.4 fl oz/acre	<ul style="list-style-type: none"> Use only on Clearfield varieties. Apply preplant surface, preplant incorporated, preemergence or early postemergence (up to 3" tall weeds). 												
Resource	4.0 - 6.0 oz/acre	<ul style="list-style-type: none"> Apply to weeds less than 5" tall. Add a crop oil concentrate at 1 - 2 pt/acre and either 28% nitrogen at 2% v/v or ammonium sulfate at 2.5 lb/acre. May cause some burn or spotting to corn leaves. 												
Spirit	1.0 oz/acre	<ul style="list-style-type: none"> Always add crop oil concentrate at 1% v/v. See label for geographic restrictions. 												
2,4-D Ester	See Label	<ul style="list-style-type: none"> Apply preplant surface or preemergence to control emerged broadleaf weeds in corn. 												
Accent 75WDG Beacon 75WDG Basis Steadfast	1/4 - 2/3 oz/acre 0.76 oz/acre 1/4 - 2/3 oz/acre 0.75 oz/acre	<ul style="list-style-type: none"> Minimum Volley ATZ NXT use rates (qts/acre): <table border="1"> <tr> <td>Soil</td> <td><3%OM</td> <td>3% or moreOM</td> </tr> <tr> <td>Coarse</td> <td>1.7</td> <td>1.7</td> </tr> <tr> <td>Medium</td> <td>1.7</td> <td>1.7- 2.3</td> </tr> <tr> <td>Fine</td> <td>1.7</td> <td>1.7-2.3</td> </tr> </table> Always add NIS at .25% (v/v); and in addition if applied in dry conditions, add 4% (v/v) clear liquid fertilizer. Banvel, Clarity, Marksman, Buctril, Buctril/ atrazine may be added to this mixture to provide burndown and residual control of broadleaf weeds. 	Soil	<3%OM	3% or moreOM	Coarse	1.7	1.7	Medium	1.7	1.7- 2.3	Fine	1.7	1.7-2.3
Soil	<3%OM	3% or moreOM												
Coarse	1.7	1.7												
Medium	1.7	1.7- 2.3												
Fine	1.7	1.7-2.3												
Basis Gold	14.0 oz/acre	<ul style="list-style-type: none"> Minimum Volley ATZ NXT use rates (qts/acre): <table border="1"> <tr> <td>Soil</td> <td><3%OM</td> <td>3% or moreOM</td> </tr> <tr> <td>Coarse</td> <td>1.7</td> <td>1.7</td> </tr> <tr> <td>Medium</td> <td>1.7</td> <td>1.7 - 2.3</td> </tr> <tr> <td>Fine</td> <td>1.7</td> <td>1.7-2.3</td> </tr> </table> Always add crop oil concentrate at 1.0% v/v or under dry arid conditions, 2.0% v/v and 28% liquid nitrogen at 2 qt/acre or ammonium sulfate at 2 lb/acre. Banvel, Clarity, Marksman, Buctril, or Tough herbicide may be added to this mixture to provide burndown and residual control of broadleaf weeds. The maximum atrazine application rate for corn is 2.5 pounds active ingredient per calendar year or 2.0 pounds atrazine active ingredient for postemergence application if no atrazine was applied preemergence. 	Soil	<3%OM	3% or moreOM	Coarse	1.7	1.7	Medium	1.7	1.7 - 2.3	Fine	1.7	1.7-2.3
Soil	<3%OM	3% or moreOM												
Coarse	1.7	1.7												
Medium	1.7	1.7 - 2.3												
Fine	1.7	1.7-2.3												

Appendix I

Procedure for Testing the Compatibility of Volley ATZ NXT and Tank Mixes with Fluid Fertilizers

Since fluid fertilizers vary, the following procedure is suggested for determining whether Volley ATZ NXT may be combined with a specific fluid fertilizer for spray tank application.

Materials Needed:

- Volley ATZ NXT and any tank mix products
- Fluid fertilizer to be used
- Adjuvant for fertilizer tank mix: Use any adjuvant cleared for use on growing crops under 40 CFR 180.1001 to improve the compatibility of Volley ATZ NXT with fluid fertilizers. The adjuvant that provides the best emulsification depends on the specific fertilizer under consideration.
- Two 1 quart, wide mouth glass jars with lid or stopper
- Measuring spoons (a 25-ml pipette or graduated cylinder provides more accurate measurement)
- Measuring cup, 8 ounces (257 ml)

Procedure:

- Pour a pint (about 473 ml) of the fluid fertilizer into each of the quart jars.
- Add Volley ATZ NXT and any tank mix combination to the jars. The order of addition is wettable powders first with mixing, followed by flowables with mixing and the EC's last. The rate of wettable powders and dry flowables is 1 1/2 teaspoon per pound of product per acre to be applied. EC's should be added at the rate of 1/2 teaspoon for each pint per acre to be applied. Premixing the wettable powders in 1 ounce of water before adding to the pint of fluid fertilizer will improve the compatibility of the final mixture.
- Add 1/2 teaspoon (2 ml) adjuvant to one of the jars, label it as "with", and mix. The rate of 1/2 teaspoon per pint is equal to 3 pints of adjuvant per 100 gallons of fluid fertilizer.
- Close both jars with lids or stoppers and mix the contents by turning the jars upside down ten times.
- Inspect the surface and body of the mixtures:
 - Immediately after completing the jar inversions
 - After allowing the jars to stand quietly for 30 minutes
 - And then again after turning the jars upside down 10 times after the 30 minute inspection

Evaluation:

If either mixture remains uniform for 30 minutes, the combination may be used. Should either mixture separate after 30 minutes, but readily remix uniformly with 10 jar inversions, the mixture can be used if adequate agitation is maintained in the tank. If the mixture with adjuvant is satisfactory but the mixture without adjuvant is not, be sure to use the adjuvant in the spray tank. Add the adjuvant first at a rate of 3 pints per 100 gallons of fluid fertilizer. Foaming may be minimized by using only moderate agitation. **If non-dispersible oil, sludge, or clumps of solids form in the mixtures, the combination should not be used.**

Appendix II

Dry Bulk Fertilizer Impregnation

Impregnation of bulk fertilizer is restricted to commercial facilities. On-farm fertilizer impregnation is prohibited. No more than 500 tons of bulk fertilizer can be impregnated per day. No single facility may impregnate fertilizer with this product for more than 30 days per calendar year.

The commercial facility impregnating the dry bulk fertilizer must inform, in writing, the user (applicator) of the dry bulk fertilizer that:

- Applicator must wear long-sleeved shirt, long pants, shoes, and socks
- The restricted entry interval is 12 hours.

All individual state regulations relating to dry bulk fertilizer blending, registration, labeling and application are the responsibility of the individual and/or company selling the Volley ATZ NXT.

Dry bulk fertilizers (Table 5) may be impregnated with this product or the tank mixtures of this product plus atrazine on corn. This product and these tank mixtures must be applied with 200 to 450 pounds of dry bulk fertilizer per acre and shallowly incorporated within 14 days prior to planting. On medium- and fine-textured soils in areas where incorporation is not planned (i.e., reduced tillage situations or in some conventional tillage situations), applications can be made up to 30 days before planting to allow moisture to move the herbicide-fertilizer mixture into the soil. On coarse-textured soils, applications can be made up to 14 days prior to planting. When applying Volley ATZ NXT alone or in tank mixes with dry bulk fertilizers, follow all directions for use and precautions on the respective tank mix product labels regarding rates, soil type, application methods and rotational restrictions. Refer to the table for broadcast rate per acre to determine the application rate per acre for the herbicide treatment to be applied.

Table 5: Approved Dry Fertilizer Ingredients for Use with Volley ATZ NXT

Fertilizer	N	P	K
Ammonium Phosphate-Sulfate	16	20	0
Ammonium Sulfate	21	0	0
Diammonium Phosphate	18	46	0
Monoammonium Phosphate	11	56	0

Table 5: Approved Dry Fertilizer Ingredients for Use with Volley ATZ NXT (Cont.)

Fertilizer	N	P	K
Potassium Chloride	0	0	60
Potassium Sulfate	0	0	52
Urea †	45	0	0

† Some ureas may be phytotoxic when high rates are applied to corn. Use only urea rates known to be safe for corn application.

For impregnating the pesticides on dry fertilizers, use an appropriate mixer equipped with suitable spraying equipment. The spray nozzles should be positioned inside the mixer to provide uniform spray coverage of the tumbling fertilizer. The Volley ATZ NXT should be sprayed uniformly onto the fertilizer using a fine spray pattern. Tank mix components may be applied as separate ingredients with powders and dry flowables added first or they may be mixed in a slurry in the proper ratio and added jointly. Volley ATZ NXT may also be impregnated on the go and applied with pneumatic applicators.

The following table provides a reference to determine the amount of Volley ATZ NXT to be mixed per ton of dry bulk fertilizer for a range of herbicide and fertilizer rates per acre.

Table 6. Volley ATZ NXT Fertilizer Impregnation Rate Conversions

Fertilizer Rate (lbs/acre)	Acres Covered (per ton)	Quarts of Volley ATZ NXT per Ton of Fertilizer to Deliver:		
		1.4 qts/ acre	1.8 qts/ acre	2.3 qts/ acre
200	10.0	14.0	18.0	23.0
250	8.0	11.2	14.4	18.4
300	6.7	9.4	12.0	15.4
350	5.7	8.0	10.3	13.1
400	5.0	7.0	9.0	11.5
450	4.5	6.2	8.0	10.4

To determine the amount of Volley ATZ NXT needed for other fertilizer rates, use the following formula:

$$\text{Volley ATZ NXT rate (quarts/acre)} \times \frac{\text{X 2000}}{\text{Pounds of fertilizer/acre}} = \frac{\text{Quarts of Volley ATZ NXT}}{\text{per ton of fertilizer}}$$

If the herbicide/fertilizer mixture is too wet, use of a drying agent is required to provide a dry, free-flowing mixture. For mixtures to be used in spinning-disc applicators, Micro-Cel E calcium silicate powder (Manville, Filtration & Minerals) is recommended for use as a drying agent. Mixtures to be used in pneumatic applicators should use Micro-Cel E or Agsorb 16/30 RVM-MS granular clay (Oil-Dri Corporation). The drying agents should be added separately and uniformly to the prepared pesticide/fertilizer mixture, in a quantity that is sufficient to provide a suitable free-flowing mixture. Generally, less than 2% Micro-Cel E or 5% Agsorb 16/30 RVM-MS by weight is required.

Precaution: To avoid potential for explosion, do not impregnate Volley ATZ NXT on ammonium sorbate nitrate, potassium nitrate, or sodium nitrate fertilizer or fertilizer blends. Do not impregnate on single (0-20-0) or triple (0-46-0) super phosphate. Do not impregnate on agricultural limestone because Volley ATZ NXT will not be absorbed.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Tenkoz warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Tenkoz MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the

control of Tenkoz or the seller. Tenkoz will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by Tenkoz. To the extent permitted by law, all such risks associated with non-directed use shall be assumed by buyer and/or user.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, tort, strict liability, or other legal theories), shall be limited to, at Tenkoz's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Tenkoz shall not be liable for losses or damages resulting from handling or use of this product unless Tenkoz is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Tenkoz be liable for consequential, incidental or special damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Tenkoz or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™® Trademarks of Corteva Agriscience and its affiliated companies.
EPA accepted 11/10/21