



TEBUCONAZOLE	GROUP	3	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

ADAMENT™ 50 WG Fungicide

EPA Reg. No. 264-1052

For Control of Red Leaf Blotch in Almonds

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

ALMOND		
Disease Control	Rate (oz./acre)	Application Timing
Red Leaf Blotch (RLB) (<i>Polystigma amygdalinum</i>)	6.0 to 8.0	Begin applications when conditions are favorable for disease but before infection. Apply on a 7- to 14-day spray schedule. Refer to University and/or extension guidelines for best application timings.
Restrictions: Do not apply more than 32 oz. of ADAMENT 50 WG FUNGICIDE per acre per season. Do not make more than 4 total applications of ADAMENT 50 WG FUNGICIDE per season. Do not apply ADAMENT 50 WG FUNGICIDE within 35 days of harvest. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. ADAMENT 50 WG FUNGICIDE is a mixture of a Group 11 and a Group 3 fungicide. To limit the potential for the development of resistance, do not apply more than 2 sequential applications of ADAMENT 50 WG FUNGICIDE or other Group 11-containing fungicides.		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167