



2ee RECOMMENDATION

FIFRA SECTION 2(ee) RECOMMENDATION

ZOLERA FX FUNGICIDE EPA REGISTRATION NO. 66330-424

USE DIRECTIONS FOR TAR SPOT CONTROL IN CORN

FOR USE AND DISTRIBUTION IN THE FOLLOWING STATES:

IA, IL, IN, KS, KY, MI, MN, MO, NE, ND, OH, PA, SD, VA, WI, WV

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by US EPA. Use of this product according to this bulletin has not been reviewed or endorsed by any of the state departments of agriculture.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

CROP	DISEASE	USE RATE FL OZ PER ACRE	USE DIRECTIONS	RESTRICTIONS
Corn (field, seed)	Tar spot (<i>Phyllachora maydis</i>)	5.0 – 6.8	For optimum results, begin applications preventively. Use the higher specified rates when disease pressure is high. Apply by ground in a minimum of 10 GPA, by air in a minimum	-Do not exceed a total of 6.8 fl oz (0.09 lb tetraconazole + 0.09 lb fluoxastrobin) of ZOLERA FX FUNGICIDE per acre per calendar year. -Do not make more than 1 application of ZOLERA FX FUNGICIDE per acre per calendar year. -Do not apply more than 0.09 lb tetraconazole per acre per year (total of all applications of tetraconazole products and application methods).

			of 2 GPA, or through chemigation in sufficient water to obtain thorough coverage of plants.	<ul style="list-style-type: none"> -Do not apply more than 0.36 lb fluoxastrobin per acre per year (total of all applications of fluoxastrobin products and application methods). -Do not apply after corn growth stage R3 (brown silk/milk). -Do not use adjuvants in sprays made between V8 (8 leaf collar) and VT (lowest branch of the tassel visible but silks have not emerged) growth stage. An adjuvant may be used at any other growth stage. -Pre-Harvest Interval (PHI): 30 days
--	--	--	---	---

March 2021

Arysta Lifescience c/o UPL NA INC.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406