



Supplemental Label

BIFENTURE® EC Agricultural Insecticide

(EPA Reg. No. 70506-57)

DIRECTIONS FOR USE ON SOYBEANS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label must be in the possession of the user at the time of pesticide application.

PESTS CONTROLLED	RATE BIFENTURE EC/ACRE	APPLICATION INSTRUCTIONS	PHI (days)
Alfalfa Caterpillar Aphids Aster Leafhopper Bean Leaf Beetle Beet Armyworm* Cloverworm Corn Earworm Corn Rootworm Adult Cucumber Beetles Cutworms European Corn Borer Fall Armyworm Flea Beetle Grasshoppers Imported Cabbageworm Japanese Beetle Adult Pea Leaf Weevil Pea Weevil Plant Bug Saltmarsh Caterpillar Sap Beetle Southern Armyworm Stink Bugs Tarnished Plant Bug Thrips Tobacco Budworm* Webworms Western Bean Cutworm Whitefly Yellowstriped Armyworm Soybean Aphid	0.033 - 0.10 lb. ai (2.1 – 6.4 fl. oz. product)	Apply in a minimum of 10 gallons/A by ground, or in 2 gallons/A by aircraft. Do not apply more than once every 30 days. Do not apply more than 0.3 lb. ai/A per season. *Because pyrethroid resistance is common for Beet Armyworm and Tobacco Budworm, consult your local or state agricultural authorities to find out if resistant pest populations are in the treatment area. If so, refer to the resistance information in the Directions for Use section of this label.	18
Lygus spp. Whitefly Two-spotted Spider Mite	0.08 – 0.10 lb. ai (5.12 – 6.4 fl. oz. product)		

Rev. 5/09
CA appr. 10/28/09

United Phosphorus, Inc.
630 Freedom Business Center
King of Prussia, PA 19406
1-800-438-6071 • www.upi-usa.com