

FLUTRIAFOL	Group	3	Fungicide
AZOXYSTROBIN	Group	11	Fungicide

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**



EPA Reg. No.: 279-3596

## 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY IN  
THE STATE OF TEXAS**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A  
MANNER INCONSISTENT WITH IT LABELING**

**For Fall and Spring Disease Management in Spring and Winter Wheat and Barley**

*THIS 2(ee) RECOMMENDATION EXPIRES ON DECEMBER 31, 2028, AND MUST NOT  
BE USED AFTER THIS DATE*

CROP	PEST	RATE OF APPLICATION
<b>Wheat</b> (Spring and Winter)  <b>Barley</b>	<b>Leaf Rust</b> ( <i>Puccinia</i> spp.) <b>Stem Rust</b> ( <i>Puccinia graminis</i> ) <b>Stripe Rust</b> ( <i>Puccinia striiformis</i> ) <b>Powdery Mildew</b> ( <i>Blumeria graminis tritici</i> ) <b>Septoria Leaf and Glume Blotch</b> ( <i>Septoria</i> spp.) <b>Tan Spot</b> ( <i>Pyrenophora tritici-repentis</i> )	<b>3 - 4 fl. oz./A</b>

### DIRECTIONS FOR USE:

- Apply Topguard® EQ fungicide in the fall after emergence and/or in early spring from tillering to jointing growth stages (Feekes stages 2-7) to provide protection against listed early season leaf diseases.
- Topguard EQ fungicide may be tank-mixed with adjuvants, fertilizer, labeled insecticides (such as **Mustang® MAXX** insecticide) or herbicides (such as **Affinity®** brand herbicides with TotalSol® soluble granules, **Harmony® Extra SG** herbicide with TotalSol soluble granules or **Finesse®** Cereal and Fallow herbicide) to implement disease, insect and weed management.
- An application of Topguard EQ fungicide may be required at flag leaf (Feekes 8 - 9) for control of foliar diseases as specified by product label.
- Apply in a minimum of 2 gallons per acre for aerial application and 10 gallons per acre by ground application.

### RESTRICTIONS:

- Do not apply more than 2 applications per year.
- Do not apply more than 0.228 lb. ai of flutriafol/A per year.
- Do not apply more than 0.4 lb. ai of azoxystrobin per acre per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Intervals (PHI): Do not apply within 30 days of harvest when grown for grain, 15 days of harvest when grown for hay, 7 days of harvest when grown for forage.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED TOPGUARD EQ FUNGICIDE, EPA REG.NO. 279-3596, LABEL MUST BE FOLLOWED. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

R-4967 081825 08-18-25



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

FMC, Topguard, Harmony, TotalSol, Mustang  
Finesse and Affinity are trademarks of FMC Corporation  
©2025 FMC Corporation. All rights reserved