

PRODUCT BULLETIN



Thiolux® **EPA Reg. No. 34704-1079**

2(ee) Recommendation[†]

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

ATTENTION

- **[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA or state lead pesticide agency.**
- Read and follow all applicable directions for use, precautions and limitations on the federally registered label attached to the product container.

Crop	Pest	Rate of Application
Strawberries	Control of Powdery Mildew	For mildew control, apply 5.0 to 10.0 lb Thiolux in 50.0 to 200 gal/A finished spray

Directions for Use:

Apply 5.0 to 10.0 pounds of Thiolux per acre in 50.0 to 200 gallons of water per acre by ground. Make applications prior to disease development in conjunction with good cultural management practices.

Restrictions:

- REI is 24 hours
- Thiolux may be applied up to the date of harvest
- Apply by ground equipment only
- Allow foliage to completely dry after application
- **Do not** apply when conditions favor wet tissue for prolonged periods (> 4 hours)
- **Do not** use on sulfur sensitive varieties

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This recommendation should be in the possession of the user at the time of pesticide application.