

Bayer Environmental Science

Bayer Environmental Science
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-800-331-2867

TEMPO® SC ULTRA Insecticide

For use on citrus and ornamentals to control
Asian citrus psyllid

EPA Reg. No. 432-1363

SLN No. CA-080015

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA

It is a violation of federal law to use this product in a manner inconsistent with its label. This label must be in the possession of the user at the time of application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered Section 3 label.

THIS LABEL IS VALID UNTIL AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

DIRECTIONS FOR USE

Only for use in and around residential areas, industrial and commercial buildings.

CITRUS: Crops of group 10 including: calamondin, Citrus citron, citrus hybrids (includes chironja, tangelo, and tango), grapefruit, kumquat, lemon, lime, mandarin (tangerine), pummelo, orange (sweet and sour), Satsuma mandarin, white sapote (Casimiroa spp.), and other cultivars and hybrids of these.

ORNAMENTALS: *Murraya* spp. (Curry Tree, Orange Jessamine)

Pest Controlled	Rate of TEMPO SC ULTRA Insecticide	Remarks
Asian citrus psyllid	65 mL (2.2 fl oz) per 100 gallons of water or 2.0 mL per 3 gallons of water	Foliar Applications: Applications should be started prior to establishment of high pest populations and reapplication made as necessary. Good spray coverage is necessary to provide the most effective level of control. Avoid excess application that could result in runoff from plant foliage. Addition of a spreader/sticker at recommended rates may enhance control of insects on certain species of ornamentals having hard-to-wet foliage.

Notes and Restrictions:

Preharvest interval (PHI): 0 days

Do not apply this product by aerial application or by any type of chemigation system. Do not apply when the average windspeed is greater than 15 mph.

Do not enter treated areas until the spray has dried.

Do not apply more than 0.15 fl oz (4.4 ml) per 1000 ft² per year.

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

As with any plant-protection product, always read and follow label directions.