THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

2(ee) Recommendation



IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

This Product Bulletin Contains New or Additional Directions For Use Which Are Recommended By Sym-Agro, Inc. Under FIFRA Section 2(ee)

FOR DISTRIBUTION AND USE ONLY IN Oregon, Washington, Idaho, Montana, Utah

EPA REG. No. 84876-2

CROP	PEST	RATE OF APPLICATION
Stone Fruit	Alternaria, Alternaria late blight (Alternaria alternata) Leaf spots (Alternaria spp.) Gray Mold Botrytis blossom blight (Botrytis cinerea) Powdery mildew (Oidium spp., Podosphaera leucotricha, Sphaerotheca pannosa) Ripe fruit rot (Monilinia fruticola, Monilinia laxa, Botrytis cinerea, Rhizopus spp.) Peach Leaf Curl (Taphrina deformans) Rust (Tranzschelia discolor) Scab (Venturia carpophila) Shot Hole (Wilsonomyces carpophilus)	20.4 – 45.7 fl oz/A (0.34 – 0.75 lb ai/A)

DIRECTIONS FOR USE:

Botrytis: For control of Botrytis gray mold apply ProBlad® Verde prior to onset of disease development when conditions favor Botrytis development up to the harvest.

Powdery mildew: Begin applications of ProBlad® Verde preventively when conditions are favorable for disease development and continue on a 7 to 10-day interval if conditions continue to favor disease development. Under conditions of severe disease pressure, use the higher labeled rate.

General: Increase spray volume as growth increases in order to provide thorough coverage of foliage and fruit for optimum disease control.

RESTRICTIONS:

Do not make more than two sequential applications of ProBlad[®] Verde Fungicide before alternating to a labeled fungicide with a different mode of action.

Do not make more than 5 applications of ProBlad® Verde Fungicide per harvest cycle.

PHI: 1 Day

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED PROBLAD VERDE® FUNGICIDE (EPA REG. No. 84876-2) LABEL MUST BE FOLLOWED. THIS RECCOMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

ProBlad is a trademark of CEV S.A. ProBlad Verde_2ee_Stone Fruit Crop Expanded Diseases_11/1/2020

