

FIFRA Sec. 2(ee) Recommendation

Nurizma™

Insecticide

For nozzle or drop-tube in-furrow applications at bed formation in potatoes grown in Idaho, Michigan, Oregon, and Washington

This Section 2(ee) Recommendation expires on December 31, 2027.

EPA Reg. No. 7969-423

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

All applicable directions, restrictions, precautions, and Conditions of Sale and Warranty on the EPA registered container label are to be followed.

Information contained in this Section 2(ee) Recommendation is not intended to replace or amend any product labeling. Always read and follow all label directions when using any pesticide alone or in tank mix combinations.

The user must have this recommendation in their possession at the time of use.

Directions For Use

Apply as a pre-plant in-furrow treatment at bed formation.

Apply into the furrow using a nozzle or drop-tube directed into the open furrow. Close the furrow completely and immediately using suitable cultivation equipment (e.g., ripper bedder, bed conditioner).

For rates based on 30", 34", 36", 38" and 40" row spacing, refer to expanded conversion table below.

Conversion Table

Nurizma Required Per Acre for Various Row Spacings in Potato					
Rate per 1000 Row Feet (fl oz product)	Rate per Acre (fl ozs product)				
	30" rows	34" rows	36" rows	38" rows	40" rows
Row Feet/Acre	17,424	15,374	14,520	13,756	13,068
0.08	1.3	1.2	1.1	1.1	1.0
0.16	2.3 ^a	2.3 ^b	2.3	2.2	2.0
^a For 2.3 fl ozs/A rate in 30-inch row spacing, use 0.13 fl oz/1000 ft					
^b For 2.3 fl ozs/A rate in 34-inch row spacing, use 0.15 fl oz/1000 ft					

Nurizma is a trademark of BASF.

© 2024 BASF Agricultural Solutions US LLC
All rights reserved.

007969-00423.20240829.NVA 2024-04-0585-0027