

# **FIFRA §2(ee) Recommendation**

## **Propicure 3.6F**

Fungicide  
EPA Reg. No. 83222-18

For The Control of Rust and Scab on Almonds in California Only

### **DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the Propicure 3.6F Fungicide container label and read and follow all directions for use, restrictions, and precautions found on that label.
- Users should have this recommendation in their possession at the time of use.
- This recommendation is permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US EPA.

### **CROP: ALMONDS**

<b>PESTS CONTROLLED</b>	<b>RATE OF APPLICATION PER ACRE</b>	<b>APPLICATION INSTRUCTIONS</b>
Rust ( <i>Tranzschelia discolor</i> )  Scab ( <i>Cladosporium carpophilum</i> )	6 to 8 fl. oz.	Apply 6 to 8 fl. oz (0.18- 0.225 lbs active) per acre. Begin application approximately 2-5 weeks after petal fall when environmental conditions are favorable for disease development. For optimum disease control, apply on a 7- to 14-day schedule. For disease management, make no more than two consecutive applications before rotation to a fungicide with a different mode of action (not a Group 3)
<ul style="list-style-type: none"><li>• Do not apply Propicure 3.6F within 60 days of harvest.</li><li>• Do not apply more than 32 fl. oz. (0.90 lb. ai) of Propicure 3.6F per acre per season.</li><li>• Do not apply more than 0.90 lbs active of any propiconazole product per acre per season.</li><li>• Do not graze livestock in treated areas or cut treated cover crop for feed.</li></ul>		

**Effective Date: 1-7-13**

Direct Ag Source, LLC  
30473 260<sup>th</sup> Street, Eldora, IA 50627