

FIFRA Section 24(c) Special Local Need



FLUMIOXAZIN GROUP 14 HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WYOMING



EPA Reg. No. 59639-221
EPA SLN No. WY-230002

FOR WEED CONTROL IN ESTABLISHED DORMANT ALFALFA GROWN FOR SEED ONLY

This label for Chateau® EZ Herbicide is valid until December 31, 2028 or until otherwise amended, withdrawn, canceled, or suspended.

| | |
|-------------------------|--------------|
| Active Ingredient | By Wt. |
| Flumioxazin* | 41.4% |
| Other Ingredients | <u>58.6%</u> |
| Total | 100.0% |

*2-[7-fluoro-3,4-dihydro-3-oxo-4-(20propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione

Chateau EZ is a suspension concentrate containing 41.4% active ingredient.

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this supplemental label and the main EPA-registered label. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

This label and the federal label for this product must be present with the user at the time of pesticide application. Read the label affixed to the container for *Chateau* EZ Herbicide and this Section 24c label before applying. Do not use in manner inconsistent with applicable directions, restrictions, and precautions provided on the label affixed to the container for *Chateau* EZ Herbicide and this Section 24c label. Use of *Chateau* EZ Herbicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for *Chateau* EZ Herbicide.

READ ENTIRE CONTAINER LABEL AND SLN LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

APPLICATION DIRECTIONS

APPLICATION AND TIMING:

- Apply *Chateau* EZ Herbicide at 4 fl oz/A to **established, dormant** seed alfalfa during late fall or winter before alfalfa shoot growth has started or between cuttings prior to six inches of regrowth. *Chateau* EZ

Herbicide may be applied to established, dormant alfalfa with a maximum amount of growth of 6 inches or less for the preemergence control of the weed listed in Table A (Weeds Controlled by Residual Activity of *Chateau* EZ Herbicide).

- Applications should be made as soon as possible after cutting and removing alfalfa to minimize injury to alfalfa growth. Applications to alfalfa greater than 6 inches of growth may result in unacceptable crop injury.
- Tank-mix *Chateau* EZ Herbicide with paraquat to control emerged seedling weeds as a dormant season application. Use a nonionic surfactant at 0.25% v/v or a crop oil concentrate at 1% v/v per the paraquat label. Apply by ground in 20-50 GPA water volume.
- To activate *Chateau* EZ Herbicide, apply at least 0.25" sprinkler irrigation within 5 days of *Chateau* EZ Herbicide application if the equivalent amount of rainfall has not occurred. Furrow irrigation to activate the herbicide is not recommended.
- Cultivation, renovation or field burning should be done **before** *Chateau* EZ Herbicide is applied. Do not cultivate after application, to maintain the herbicide barrier and residual weed control.

METHODS OF APPLICATION:

Chateau EZ Herbicide can be applied:

1. Using ground equipment at 20 to 50 gallons of waters per acre (GPA)
2. By air at 10 GPA. When making applications, observe drift management restrictions and precautions listed under AERIAL APPLICATION on the container label
3. By chemigation. Refer to the CHEMIGATION section of the *Chateau* EZ Herbicide container label for use, restrictions and precautions when applying through an irrigation system

RESTRICTIONS AND LIMITATIONS:

- Do not apply more than 4 fl oz of *Chateau* EZ Herbicide per acre during a single application.
- Do not apply more than 8 fl oz of *Chateau* EZ Herbicide per acre during a single growing season.
- Do not make a sequential *Chateau* EZ Herbicide application within 60 days of the first *Chateau* EZ Herbicide application.
- Application will result in burning of treated leaves and stems. **Users should understand and accept this risk before using *Chateau* EZ Herbicide on alfalfa.**
- Do not apply within 25 days of harvest or grazing.
- Do not use on intended mixed alfalfa-grass stands.
- When making aerial applications: (1) do not apply this product within 40 ft. of non-target plants including non-target crops, and (2) do not apply this product within 40 ft. of streams, wetlands, marshes, ponds, lakes or reservoirs.
- Do not apply within 300 yards of non-dormant pears.
- Follow Rotational Restrictions indicated on container label.
- Do not apply with any adjuvant or tank mix with any products formulated as an emulsifiable concentrate "EC", unless making application following last cutting of the season.
- When tank mixing *Chateau* EZ Herbicide with other products registered for alfalfa grown for seed, the most restrictive labeling of any tank mix product must be followed.

SEED CONDITIONING REQUIREMENTS:

- For use only on fields in production of alfalfa seed.
- Not for use on fields producing alfalfa for livestock feed.
- Do not graze current year's treated alfalfa seed crop.
- Do not cut current year's treated alfalfa seed crop for hay or forge.
- Treated alfalfa seed is not to be used for sprouting. All alfalfa seed treated with *Chateau* EZ Herbicide is to be tagged at the conditioning plant, "NOT FOR HUMAN OR ANIMAL CONSUMPTION". It shall be the grower's responsibility to notify the conditioning plant of any alfalfa seed crop treated with *Chateau* EZ Herbicide.
- Screenings and seed from seed conditioning are prohibited from feed channels. All *Chateau* EZ Herbicide treated alfalfa seed screenings must be removed from the feed market.

Table A. Weeds Controlled by Residual Activity of CHATEAU EZ HERBICIDE

| BROADLEAF WEED SPECIES | | | | |
|------------------------------------|----------------------------------|-----------------------|------------------|-------------|
| COMMON NAME | SCIENTIFIC NAME | ORGANIC MATTER | SOIL TYPE | RATE |
| Bristly Starbur | <i>Acanthospermum hispidum</i> | Up to 5% | All Soil Types | 4 fl oz/A |
| Carpetweed | <i>Mollugo verticillata</i> | | | |
| Chickweeds | | | | |
| Common | <i>Stellaria media</i> | | | |
| Mouseear | <i>Cerastium vulgatum</i> | | | |
| Coffee Senna | <i>Cassia occidentalis</i> | | | |
| Copperleaf, Hophornbeam | <i>Acalypha ostryifolia</i> | | | |
| Dandelion | <i>Taraxacum officinale</i> | | | |
| Eclipta | <i>Eclipta prostrate</i> | | | |
| Eveningprimrose, Cutleaf | <i>Oenothera laciniata</i> | | | |
| False Chamomile | <i>Tripleurospermum maritima</i> | | | |
| Flixweed | <i>Descurainia spophia</i> | | | |
| Florida Beggarweed | <i>Desmodium tortuosum</i> | | | |
| Florida Pusley | <i>Richardia scabra</i> | | | |
| Golden Crownbeard | <i>Verbesina encelioides</i> | | | |
| Groundsel, Common | <i>Senecio vulgaris</i> | | | |
| Hairy Indigo | <i>Indigofera hirsuta</i> | | | |
| Hemp Sesbania | <i>Sesbania exaltata</i> | | | |
| Henbit | <i>Lamium amplexicaule</i> | | | |
| Jimsonweed | <i>Datura stramonium</i> | | | |
| Kochia | <i>Kochia scoparia</i> | | | |
| Lambsquarters, Common | <i>Chenopodium album</i> | | | |
| Little Mallow | <i>Malva parviflora</i> | | | |
| Marestail/Horseweed | <i>Conyza canadensis</i> | | | |
| Morningglories | | | | |
| Entireleaf | <i>Ipomoea hederacea</i> var. | | | |
| Ivyleaf | <i>Ipomoea hederacea</i> | | | |
| Red/Scarlet | <i>Ipomoea coccinea</i> | | | |
| Smallflower | <i>Jacquemontia tamnifolia</i> | | | |
| Tall | <i>Ipomoea purpurea</i> | | | |
| Mustard | | | | |
| Tansy | <i>Descurainia pinnata</i> | | | |
| Tumble | <i>Sisymbrium altissimum</i> | | | |
| Wild | <i>Brassica kaber</i> | | | |
| Nightshades | | | | |
| Black | <i>Solanum nigrum</i> | | | |
| Eastern Black | <i>Solanum ptycanthum</i> | | | |
| Hairy | <i>Solanum sarrachoides</i> | | | |
| Pigweeds | | | | |
| Palmer Amaranth | <i>Amaranthus palmeri</i> | | | |
| Redroot | <i>Amaranthus retroflexus</i> | | | |
| Smooth | <i>Amaranthus hybridus</i> | | | |
| Spiny Amaranth | <i>Amaranthus spinosus</i> | | | |
| Tumble | <i>Amaranthus albus</i> | | | |
| Prickly Lettuce (China Lettuce) | <i>Lactuca serriola</i> | | | |

Table A. Weeds Controlled by Residual Activity of CHATEAU EZ HERBICIDE

| BROADLEAF WEED SPECIES | | | | |
|------------------------|--|----------------|----------------|-----------|
| COMMON NAME | SCIENTIFIC NAME | ORGANIC MATTER | SOIL TYPE | RATE |
| Prickly Sida (Teaweed) | <i>Sida spinosa</i> | Up to 5% | All Soil Types | 4 fl oz/A |
| Puncturevine | <i>Tribulus terrestris</i> | | | |
| Purslane | | | | |
| Common | <i>Portulaca oleracea</i> | | | |
| Horse | <i>Trianthema portulacastrum</i> | | | |
| Radish, Wild | <i>Raphanus raphanistrum</i> | | | |
| Ragweed, Common | <i>Ambrosia artemisiifolia</i> | | | |
| Redmaids | <i>Calandrinia ciliata</i> var. <i>menziesii</i> | | | |
| Russian Thistle | <i>Salsola iberica</i> | | | |
| Shepherd's-purse | <i>Capsella bursa-pastoris</i> | | | |
| Smartweeds | | | | |
| Ladysthumb | <i>Polygonum persicaria</i> | | | |
| Pennsylvania | <i>Polygonum pensylvanicum</i> | | | |
| Smellmelon | <i>Cucumis melo</i> | | | |
| Spotted Spurge | <i>Euphorbia maculate</i> | | | |
| Spurred Anoda | <i>Anoda cristata</i> | | | |
| Tropic Croton | <i>Croton glandulosus</i> | | | |
| Velvetleaf | <i>Abutilon theophrasti</i> | | | |
| Venice Mallow | <i>Hibiscus trionum</i> | | | |
| Waterhemp | | | | |
| Common | <i>Amaranthus rudis</i> | | | |
| Tall | <i>Amaranthus tuberculatus</i> | | | |
| Wild Poinsettia | <i>Euphorbia heterophylla</i> | | | |
| Wormwood, Biennial | <i>Artemisia biennis</i> | | | |
| GRASS WEED SPECIES | | | | |
| Barnyardgrass | <i>Echinochloa crus-galli</i> | | | |
| Bluegrass, Annual | <i>Poa annua</i> | | | |
| Crabgrass, Large | <i>Digitaria sanguinalis</i> | | | |
| Foxtail, Giant | <i>Setaria faberi</i> | | | |
| Goosegrass | <i>Eleusine indica</i> | | | |
| Lovegrass, California | <i>Eragrostis diffusa</i> | | | |
| Panicums | | | | |
| Fall | <i>Panicum dichotomiflorum</i> | | | |
| Texas | <i>Panicum texanum</i> | | | |
| Ryegrass, Italian | <i>Lolium multiflorum</i> | | | |
| Signalgrass, Broadleaf | <i>Brachiaria platyphylla</i> | | | |

FIFRA §24(c) Registrant: Valent U.S.A. LLC
P.O. Box 5075
San Ramon, CA 94583

Chateau is a registered trademark of Valent U.S.A. LLC