



Supplemental Label

BIFENTURE® EC Agricultural Insecticide (EPA Reg. No. 70506-57)

DIRECTIONS FOR USE ON FRUITING VEGETABLES

Eggplant, bell and non-bell pepper, pepino, groundcherry, tomato, tomatillo

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label must be in the possession of the user at the time of pesticide application.

CROP	PESTS CONTROLLED	RATE BIFENTURE EC/ACRE	APPLICATION INSTRUCTIONS	PHI (days)
Eggplant Pepper (bell and non- bell) Pepino Groundcherry	Armyworms (including: Beet Armyworm, Fall Armyworm, Southern Yellowstriped Armyworm) Cabbage Looper Colorado Potato Beetle Corn Earworm Cucumber Beetle Cutworms European Corn Borer Flea Beetle Leafhoppers Leafminers Loopers Pepper Weevil Plant Bug Stink Bug Thrips Tomato Pinworm Tomato Hornworm Vegetable Leafminer Whitefly	0.033-0.10 lb. ai (2.1-6.4 fl. oz. product)	Apply in at least 2 gallons finished spray per acre by air or at least 10 gallons per acre by ground. When application is made by air, 1-2 qts. emulsified oil may replace 1-2 qts of water in the finished spray. For best control thorough coverage is necessary. Do not make applications less than 7 days apart. Do not apply more than 0.2 lb. a.i. (12.8 fl. oz. product) per acre per season.	7
	Banks Grass Mite Broad Mite Carmine Mite Lygus spp. Pacific Spider Mite Twospotted Spider Mite	0.08-0.10 lb. ai (5.12-6.4 fl. oz. product)		

CROP	PESTS CONTROLLED	RATE BIFENTURE EC/ACRE	APPLICATION INSTRUCTIONS	PHI (days)
Tomato Tomatillo	Aphids Armyworms (including: Beet Armyworm, Fall Armyworm, Southern Yellowstriped Armyworm) Bean Leaf Beetle Cabbageworm Carmine Mite Cloverworm Corn Earworm Corn Rootworm Cucumber Beetle Cutworms Diamondback Moth European Corn Borer Flea Beetle Fleahopper Grasshopper Japanese Beetle (adult) Leafhoppers Loopers Lygus spp. Melonworm Pea Weevil Pea Leaf Weevil Pickleworm Plant Bug Rindworm Salt Marsh Caterpillar Sap Beetle Seedpod Weevil Squash Bugs Stink Bug spp. Tobacco Budworm Tarnished Plant Bug Thrips Whitefly	0.033-0.08 lb. ai (2.1-5.2 fl. oz. product)	Apply in at least 3 gallons finished spray/A by air or at least 15 gallons/A by ground. For best control thorough coverage is necessary. Do not make applications less than 7 days apart. Do not make more than 4 applications per season.	1
	Two Spotted Spider Mite	0.08-0.10 lb ai (5.12-6.4 fl. oz product)		

Rev. 5/09
CA appr. 10/28/09

United Phosphorus, Inc.
630 Freedom Business Center
King of Prussia, PA 19406
1-800-438-6071 • www.upi-usa.com