

INSECTICIDAL VIRUS

Biological Insecticide for Control of the Codling Moth and Oriental Fruit Moth

FOR ORGANIC PRODUCTION



ACTIVE INGREDIENT*:

Cydia pomonella (granulovirus is	solate V22)	 	0.06%
OTHER INGREDIENTS			99.4%
TOTAL:			100.0%

^{*}Contains a minimum of 3×10¹³ viral occlusion bodies per liter of product.

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE INSIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

FIRST AID			
IF SWALLOWED:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. 		
IF ON SKIN OR CLOTHING:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 		
IF INHALED:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. 		
IF IN EYES:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice 		

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For general information on product use, etc., call the National Pesticide Information Center at 1-800-858-7378. For emergencies, call the National Poison Control Hotline at 1-800-222-1222.

MANUFACTURED FOR:

Andermatt Group AG Stahlermatten 6 CH-6146 Grossdietwil Switzerland

EPA Reg No. 69553-1-70051 EPA Est. No.

DISTRIBUTED IN THE US BY:

Certis USA LLC 9145 Guilford Road, Suite 175 Columbia, MD 21046

ESL20210825 Ver. 20211202 (Name)

This is a Specimen Label. It may not reflect the most-recent approved label for use in your state. Always refer to the label on the product packaging for approved use instructions. Please contact your Certis sales representative for more information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, absorbed through skin, or inhaled. Avoid contact with skin, eyes or clothing, and breathing spray mist. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE): Applicators and other handlers must wear:

- · long-sleeved shirt and long pants
- · waterproof gloves or chemical-resistant gloves
- shoes plus socks

Mixers/loaders and applicators must wear a NIOSH-approved particulate respirator with any R or P filter with NIOSH approval number prefix TC-84A or a NIOSH-approved powered air purifying respirator with an HE filter with NIOSH approval number prefix TC-21C. Repeated exposures to high concentrations of microbial proteins can cause allergic sensitization.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

Environmental Hazards

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, wear:

- Coveralls
- Waterproof gloves
- Shoes plus socks

PRODUCT INFORMATION

MADEX HP acts very specifically against larvae of the apple codling moth, *Cydia pomonella*, and the Oriental fruit moth, *Grapholita molesta*, as a pre-harvest biological insecticide on pome and stone fruit trees including apple, quince, pear, peach, nectarine, apricot, plum, cherry, walnut, almond, pluot, and aprium. After ingestion by the larva, the virus multiplies within the cells of the host insect causing a fatal infection. Rapid virus multiplication within the host cells results in cell destruction and death of the organism, typically within 3-7 days.

USE INFORMATION

Start spraying shortly before larvae hatch from the eggs and before the larvae feed their way into the fruit. In general, one to two sprays per generation are sufficient. Due to unfavorable weather conditions, flight, egg laying, and hatching may extend over a longer period of time and up to 6 applications may be needed. Frequent applications at low rates may be more effective than one or two applications at high rates.

USE RESTRICTION:

Do not apply this product through any type of irrigation system.

MIXING AND APPLICATION INSTRUCTIONS

Mix MADEX HP with water and apply as a spray to fruit and foliage of trees in orchards and nurseries. Use water at or near neutral pH (7).

CROP	RATE	COMMENTS
Pome and Stone fruit trees, including apple, quince, pear, peach, nectarine, apricot, plum, prunes, cherry, walnut, almond, pluot and aprium	0.5 – 3.0 fluid ounces per acre, depending on pest pressure and application frequency.	
		Consult your local Cooperative Extension Service if you need advice about how to attain good spray coverage in your orchard.
		Make at least two (2) applications of MADEX HP per codling moth or Oriental fruit moth larval generation. Target small larvae early in their life cycle. Repeat as necessary to maintain control. MADEX HP breaks down
		under direct exposure to sunlight. If moths are still flying and laying eggs, reapply after 6-8 sunny days (counting 2 partially sunny days as 1 sunny day).

MADEX HP can be applied up to and including the day of harvest.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store this product in original sealed container in a cool, dry place inaccessible to children and pets. Store at temperatures below 41°F (5°C). Bioactivity may be impaired by storage above 86°F (30°C).

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by State and local ordinances. If burned, stay out of smoke.

WARRANTY: Certis USA, LLC warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. To the extent consistent with applicable law, NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

ESL20210825 (Name) Ver. 20211202