

Supplemental Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Grasp[®] SC

EPA Reg. No. 62719-500

For Distribution and Use Only in the State of Arkansas

Aerial Application of Tank Mixes of Grasp SC plus Command 3ME and Grasp SC plus Command 3ME and Other Herbicides

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Grasp[®] SC herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Grasp SC according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Grasp SC.

Directions for Use

For aerial application requirements, the user must refer to Arkansas Pesticide Use and Application Act, ACT 389 of 1975 as amended, and the regulations written pursuant thereto. Tank mixes including Facet or Quinstar must comply with the buffer zones required by the state for application of these products.

Aerially apply tank mixes of Grasp SC plus Command 3ME or Grasp SC plus Command 3ME and other herbicides for weed management in rice. The following herbicides may be tank mixed with Grasp SC plus Command 3ME for additional weed spectrum: Grandstand[®] R herbicide; Facet; Quinstar; Permit; SuperWham; Facet + Permit; Quinstar + Permit.

The addition of a crop oil concentrate or methylated seed oil at 1 quart per acre must be included.

Uniformly apply Grasp SC or tank mixes of Grasp SC by aircraft in no less than 10 gallons of water per acre total spray volume. Apply postemergence to the rice at the 1 to 2 leaf stage of growth and when weeds and grasses are small. Weeds and grasses should be actively growing at the time of application. If not, flush field in sufficient time so that rice and weeds are actively growing at time of treatment. Any factor, such as reduced spray volume, which adversely affects coverage and canopy penetration will have a negative effect on the performance of Grasp SC. Use nozzle types and arrangements that provide maximum coverage and minimize the potential for off target movement of spray particles.

Use Rates

Grasp SC - 2 to 2.3 fl oz per acre

Command 3ME – Refer to the 24(c) Special Local Need registration for Command 3ME for rates, aerial use directions, and precautions.

Permit, Grandstand R, Facet, Quinstar, and SuperWham – Refer to the product label for rate, use directions, and precautions.

Weeds Controlled**Common Name****Scientific Name****Residual control**

barnyardgrass	(<i>Echinochloa crus-galli</i>) ¹
broadleaf signalgrass	(<i>Brachiaria platyphylla</i>)
fall panicum	(<i>Panicum dichotomiflorum</i>)
sprangletop	(<i>Leptochloa</i> spp.)

Postemergence control

alligatorweed	(<i>Alternanthera philoxeroides</i>) ³
cocklebur	(<i>Xanthium strumarium</i>)
dayflower	(<i>Commelina communis</i>)
ducksalad	(<i>Heteranthera limosa</i>)
eclipta	(<i>Eclipta alba</i>)
hemp sesbania	(<i>Sesbania exaltata</i>)
Indian/northern jointvetch	(<i>Aeschynomene</i> spp.)
junglerice	(<i>Echinochloa colona</i>)
moringglory spp.	(<i>Ipomea</i> spp.) ³
pigweed	(<i>Amaranthus</i> spp.) ²
redstem	(<i>Ammania</i> spp.) ³
rice flatsedge	(<i>Cyperus iria</i>)
smartweed spp, annual	(<i>Polygonum</i> spp.)
Texas/Mexicanweed	(<i>Caperonia</i> spp.)
yellow nutsedge	(<i>Cyperus esculentus</i>) ³

¹Including propanil and quinclorac resistant barnyardgrass.

²Will not control ALS resistant pigweed.

³Suppression only.

Specific Use Precautions

Precautions must be taken to minimize off-target drift of pesticide applications. Off-target movement of Command 3ME can cause foliar whitening or yellowing of some plants. Do not make applications containing Command 3ME when spray particles may be carried by air currents to areas where sensitive crops and plants are growing or if temperature inversions exist. Prior to application, adjacent properties must be checked and applications within 300 feet of desirable plants must be avoided.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. This labeling must be in the possession of the user at the time of pesticide application.

™® Trademarks of Corteva Agriscience and its affiliated companies

R194-022

Accepted: 06/04/2019

Replaces: R194-017