

Drexel®

Beau-Ron® L

Micronutrient

GUARANTEED ANALYSIS:

Boron (B) 10.0%
Derived from Boric acid.
Contains 1.1 pounds elemental Boron per gallon.
Density, Lbs./Gallon @ 68°F: 11.1 Lbs. (5.035 kg)

KEEP OUT OF REACH OF CHILDREN

WARNING

Causes skin irritation.
Causes serious eye irritation.
Harmful if swallowed.
Suspected of damaging fertility or the unborn child.



See FIRST AID and Additional Precautionary Statements Below

Read Entire Label Before Using This Product

SHAKE WELL BEFORE USING (Recirculate Contents Before Use)

Net Weight: 27.75 lb (12.59 kg) **Net Content: 2.5 gal (9.46 l)**

F1743

Information about the components of this lot of fertilizer material may be obtained by writing to Drexel Chemical Company, P.O. Box 13327, Memphis, TN 38113-0327, and giving the lot number found on the container.

PRECAUTIONARY STATEMENTS

WARNING: Causes skin and eye irritation. Harmful if swallowed. Suspected of damaging fertility or the unborn child. Do not get on skin, in eyes or on clothing. Take off contaminated clothing and wash before reuse. If swallowed, call poison center, doctor/physician if you feel unwell. Do not eat, drink or smoke when using this product. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTION EQUIPMENT (PPE)

Wear eye protection, face protection, protective clothing and protective gloves.

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- If skin irritation occurs, get medical advice.

IF IN EYES:

- Rinse cautiously with water for 15 to 20 minutes.
- Remove contact lenses, if present and easy to do so. Continue rinsing.
- If eye irritation persists, get medical attention.

IF SWALLOWED:

- Call a poison control center or doctor if you feel unwell.
- Rinse mouth.

IF INHALED:

- Move person to fresh air and keep comfortable.
- Call poison control center or doctor for treatment advice if you feel unwell.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency call CHEMTREC at 1-800-424-9300.

GENERAL INFORMATION

WARNING: Contains Boron. Do not use on plants sensitive to Boron. The use of this product on any crop(s) other than those recommended may result in serious injury to the crop(s).

BEAU-RON® L is for use as a nutrient fertilizer supplement. This product is a liquid formulation containing 1.1 pounds per gallon of elemental Boron. The total amount of this product applied over the growing season should not exceed the total recommended for your area (typically between 1 to 2 pounds of elemental Boron per acre) and should be based on soil tests and/or plant analysis and yield goals along with previous experience and growing season conditions. Do not exceed 0.5 pound per acre of this product per application.

DIRECTIONS FOR USE

This product is recommended for aerial, concentrate and dilute application to Field, Nut, Tree and Vegetable crops.

Note: When using products packaged in a dissolvable PVA bag, always put these in the tank-mix first and dissolve; add this product after this has been done.

Rates and timing of applications are dependent on local conditions, and should be made only in conjunction with the recommendation of the local authority. Soil and/or plant analysis should be obtained and the results used as a guide. This product may be applied by ground or aerial equipment. Thorough coverage of this product is necessary for best results and sufficient water should be used to assure uniform application.

SOIL APPLICATION: This product may be applied with starter, side dress and drip irrigation fertilizer suspension except aqua ammonia.

Mixing Sequence: 1) One-third liquid fertilizer, 2) This product, 3) Balance of liquid fertilizer while spray tank is under agitation. This product may be applied with most herbicides.

FOLIAR APPLICATION: Foliar fertilization is intended as a supplement to a regular fertilization program and may not, by itself, provide all the nutrients normally required by crops or other intended plants. This product, when applied foliarly, may be mixed with most insecticides, fungicides or foliar fertilizers.

Mixing Sequence: 1) Water, 2) This product, 3) Pesticide while spray tank is under agitation.

While this product is compatible with most pesticides and liquid fertilizers, a compatibility test is recommended if the desired combination has not previously been used. When using pesticides, be sure to carefully read and follow all use directions on the product label.

Split applications are recommended if the crop requires more than 2 quarts per acre. Use adequate spray volumes to assure good coverage. Refer to the crop and rate recommendation chart to determine crops to which split application would apply.

Manufactured By:

Drexel Chemical Company
P.O. Box 13327, Memphis, TN 38113-0327
(901) 774-4370
SINCE 1972

BEAU-RON and the DREXEL logo are registered trademarks of Drexel Chemical Company.

AERIAL APPLICATION: This product may be applied by air to the labeled crops. Thorough coverage is essential. Do not apply less than 5 gallons of spray volume per acre when applying by air. Follow all local and state guidelines for aerial application of agricultural products. See the crop and rate recommendation chart for the specific rate of this product to be used per crop.

Mixing Sequence: 1) Water, 2) This product, 3) Pesticide while spray tank is under agitation. Note the preceding dissolvable PVA bag products statement.

USE PRECAUTION: Rates and timing of Boron applications are dependent on local conditions and should always be made as a result of soil and/or plant tissue analysis and in conjunction with your local agricultural authority. When used as directed, this product does not supply all the nutrients required by plants and is to supplement a soil fertility program based on soil tests. Please refer to your local dealer representative or extension agent for use guidelines. Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to, a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

CROP AND RATE RECOMMENDATIONS

FIELD CROPS ⁴	RATE (Qts./Ac.)
Alfalfa ³	2 to 4
Clover ³	2 to 4
Coffee	Consult Local Usage
Corn	1 to 2
Cotton	1 to 2
Lentils	1 to 2
Peanuts	1 to 2
Peas (Field)	1 to 2
Rape (Canola) ³	1 to 2
Small grains: Barley, Oats, Wheat	1 to 2
Sorghum	1 to 2
Soybeans	1 to 2
Sugarbeets ¹	2 to 4
Sunflowers ^{1,3}	2 to 4

FRUIT AND NUT CROPS ⁴	RATE (Qts./Ac.)
Almonds	2 to 4
Apples ³	2 to 4
Apricots ²	6 to 8
Avocados	Soil and/or Plant Analysis
Cherries	Soil and/or Plant Analysis
Citrus ²	1 to 3
Figs	Soil and/or Plant Analysis
Grapes ²	3 to 5
Peaches	2 to 4
Pears ²	2 to 4
Pecans	Soil and/or Plant Analysis
Plums, Prunes ²	2 to 8
Strawberries	2 to 4
Walnuts ²	6 to 8

VEGETABLE CROPS ⁴	RATE (Qts./Ac.)
Artichokes	Soil and/or Plant Analysis
Beans	Soil and/or Plant Analysis
Beets (Red table) ³	2 to 4
Broccoli ³	3 to 8
Brussels sprouts ³	1 to 6
Cabbage ³	1 to 6
Carrots ³	1 to 2
Cauliflower ³	2 to 8
Celery ³	1 to 6
Lettuce ³	1 to 3
Onions	1 to 3
Potatoes	1 to 4
Pumpkin	Soil and/or Plant Analysis
Radishes	1 to 3
Spinach ³	1 to 3
Sweet potatoes	Soil and/or Plant Analysis
Tomatoes ³	1 to 3
Turnips	1 to 3

¹ Apply up to 2.5 quarts when Sugarbeets and Sunflowers are at least 6 inches high. Apply up to 2.5 quarts approximately 30 days later.

² Care should be taken when applying this product to these crops. Excessive amounts of Boron may cause crop damage.

³ **FOR USE IN WISCONSIN:** Beau-Ron L can be applied to these crops commonly grown in Wisconsin and requiring a medium to high level of Boron (B). Foliar fertilization can be an effective remedy for diagnosed plant deficiencies of micronutrients, but may cause plant damage if applied at more than recommended rates or concentrations. Use of this product is recommended only as a supplement to a regular fertilization program and only on plants with confirmed micronutrient deficiencies.

⁴ **FOR USE IN PENNSYLVANIA:**

- The product may not be applied near water, storm drains or drainage ditches.
- The product may not be applied if heavy rain is expected.
- The product may only be applied to the intended application site.

THE FOLLOWING CHART CAN BE USED AS A GUIDE FOR APPLYING THE CORRECT RATES OF THIS PRODUCT.

CONVERSION CHART	
LBS. OF BORON (B) TO BE APPLIED PER ACRE	AMOUNT OF THIS PRODUCT NEEDED PER ACRE
0.10	11.0 fl.ozs.
0.20	1.4 pts.
0.50	1.8 qts.
0.75	2.7 qts.
1.00	3.6 qts.
1.50	5.4 qts.
2.00	7.1 qts.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow all governmental requirements for the disposal of agricultural spray solutions. Triple rinse container. Empty rinsate into spray tank. Dispose of empty container according to approved local practices. Contact County Agricultural Commissioner for disposal information.

WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

To the extent consistent with applicable law, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

BEAU-RON and the DREXEL logo are registered trademarks of Drexel Chemical Company.