

TECHNICAL INFORMATION BULLETIN FOR THE USE OF MAJESTENE® BIONEMATICIDE

On Peanuts (all varieties)

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

MAJESTENE®

EPA Reg. No. 84059-14

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Target Nematodes	Application Method	Product Use Rate per Application	Application Instructions
Peanuts – all varieties	Suppression of plant parasitic nematodes including root knot, sting and lesion nematodes	Pre-plant, in furrow and T- band applications	1 - 2 gallons per acre	Apply Majestene into the seed furrow and cover with soil during planting at a rate of 1-2 gallons per acre. Adjust rates per 1000 foot of row according to Table 1 below.
				Or apply Majestene in a 4-6" T-band spray over an open seed furrow andn immediately cover with soil during planting at a rate of 1-2 gallons per acre. Adjust rates per 1000 foot of row according to Table 1 below.
		Post-emergence applications		Apply Majestene in a 12-18" band over the row. Irrigate immediately after application and complete within 24 hours unless natural rainfall is received.
	has a ma hawyat intowy			Or apply Majestene as a side dress application during cultivation at a rate of 1-2 gallons per acre.

MAJESTENE® has a **pre-harvest interval (PHI)** of **0 days**.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.



Table 1.

Rate	In-Furrow and T-band Application Rates Product per 1000 ft. row.						
. 13.10	30" Rows	32" Rows	34" Rows	36" Rows	38" Rows	40" Rows	
1-2 gallons MAJESTENE per acre	7.3-14.7 fluid ounces	7.8-15.7 fluid ounces	8.3-16.7 fluid ounces	8.8-17.6 fluid ounces	9.3-18.6 fluid ounces	9.8-19.6 fluid ounces	

Nematode management considerations - To obtain maximum results in a nematode management program, it is important to take soil samples prior to application to assess nematode populations and determination of an appropriate management strategy. Utilization of the services of a fieldperson or crop consultant who is knowledgeable in nematode management is advised. When nematode populations are moderate to heavy, use the higher label rate.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

MAJESTENE® is a registered trademark of Marrone Bio Innovations, Inc. © Copyright Marrone Bio Innovations, Inc., 2014
Marrone Bio Innovations, Inc.
1540 Drew Ave.
Davis, California 95618
Telephone (530) 750-2800
www.marronebioinnovations.com

Phone: 530-750-2800