



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN MONTANA, NORTH DAKOTA, SOUTH DAKOTA

BATALIUM® AMPED™ Herbicide (EPA Reg. No. 70506-615)

REDUCED RATE OF BATALIUM AMPED TANK-MIXED WITH FLUROXYPYR AND BROMOXYNIL FOR USE IN CERTAIN SOIL CONDITIONS ON SPRING, WINTER, OR DURUM WHEAT ONLY

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

As a tank-mix with fluroxypyr and bromoxynil to maintain control of broadleaf weeds listed on the BATALIUM AMPED HERBICIDE label:

Apply BATALIUM AMPED HERBICIDE at the reduced rate of 12 fl oz/A in winter wheat and spring wheat, including durum wheat, for soils with a pH of 7.8 or greater and if organic matter is less than 3%. See table below for tank mix rates.

Follow all applicable directions (including mixing instructions), restrictions, and precautions on the labels of all products used in this tank-mix.

BATALIUM AMPED HERBICIDE USE RATE and RESPECTIVE TANK-MIX RATES TO BE ADDED

BATALIUM AMPED HERBICIDE Rate (fl oz/A)	Fluroxypyr¹ Minimum Tank-Mix Rate (lb ae/A)	Bromoxynil² Minimum Tank-Mix Rate (lb ae/A)
12	0.024	0.065

¹Do not cumulatively apply more than 0.25 lb ae/A fluroxypyr per year.

²Do not cumulatively apply more than 0.50 lb ae/A bromoxynil per year.

Rev. 2/27/23