Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

PastureGard® HL

EPA Reg. No. 62719-637

2(ee) Recommendation[†]

Control of Sumac with herbicide wick applicators

ATTENTION

- [†] This recommendation is permitted under FIFRA 2(ee).
- It as a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions and limitations on the product label attached to the container for PastureGard[®] HL herbicide.

Directions for Use

Apply PastureGard HL in a 10% v/v solution with water in a wick applicator to control sumac. Wick applications should target the leaf canopy. Therefore, to optimize coverage, position the wick applicator 6-12 inches below the target plants' height. Given that individual plants within a sumac population may vary in height, some plants may not receive adequate coverage the first application and require follow up applications in subsequent years in order to achieve acceptable control. Do not apply more than 0.5 lb ae/acre of fluroxypyr per annual growing season.

Plants sensitive to the active ingredients in PastureGard HL that come into contact with the wick will also be affected. Best practices with wicking are to place the herbicide onto plants that are taller than the surrounding vegetation so as to target control of only the undesirable plant.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R404-006 Issued: 7/18/16 Initial printing