

TECHNICAL INFORMATION BULLETIN

THIOPHANATE-METHYL

GROUP

1

FUNGICIDE

TALARIS[™] 70 WSB

EPA Reg. No. 87373-6-91234

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF FLORIDA,
GEORGIA, OREGON & WASHINGTON**

Crown Rot* (*Colletotrichum spp.*) Suppression Only

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Talaris[™] 70 WSB** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

STRAWBERRIES			
PESTS	LBS. PRODUCT per ACRE	LBS. AI per ACRE	APPLICATION INSTRUCTIONS
Crown Rot* (<i>Colletotrichum spp.</i>) Suppression only *Not for use in California	0.75 to 1.0 lb. Use highest rate under severe conditions	0.525-0.7 lb.	Begin applications after establishment of transplants and continue through first bloom at 10 to 14-day intervals. Use the high rate if the field have a history of <i>Colletotrichum</i> crown rot and/or conditions are favorable for development of disease. Will not control <i>Phytophthora</i> species.
Restrictions: Do not apply more than 4 lbs. of product (2.8 lbs. a.i.)/A/year. Do not apply more than 1.0 lb. of product (0.7 lb a.i.) / A / application. Do not make more than 4 applications per year. Minimum Retreatment Interval = 7 days REI = 1 day PHI = 1 day			



Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513